

CORRECTED VERSION

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
10 January 2002 (10.01.2002)

PCT

(10) International Publication Number
WO 02/002506 A3(51) International Patent Classification⁷: C07C 243/22,
243/28, 239/20, A61K 31/165, A61P 25/28LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN,
MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI,
SK (utility model), SK, SL, TJ, TM, TR, TT, TZ, UA, UG,
UZ, VN, YU, ZA, ZW.

(21) International Application Number: PCT/US01/20930


(22) International Filing Date: 29 June 2001 (29.06.2001)

(25) Filing Language: English

(26) Publication Language: English

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).(30) Priority Data:
60/215,323 30 June 2000 (30.06.2000) USPublished:
— with international search report(71) Applicant: ELAN PHARMACEUTICALS, INC.
[US/US]; 800 Gateway Boulevard, So. San Francisco, CA
94080 (US).(88) Date of publication of the international search report:
29 August 2002(72) Inventors: FANG, Lawrence, Y.; 1193 Beach Park Boule-
vard, Foster City, CA 94404 (US). JOHN, Varghese; 1772
18th Avenue, San Francisco, CA 94122 (US).(48) Date of publication of this corrected version:
20 November 2003(81) Designated States (*national*): AE, AG, AL, AM, AT (util-
ity model), AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA,
CH, CN, CO, CR, CU, CZ (utility model), CZ, DE (util-
ity model), DE, DK (utility model), DK, DM, DZ, EC, EE
(utility model), EE, ES, FI (utility model), FI, GB, GD, GE,
GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ,(15) Information about Correction:
see PCT Gazette No. 47/2003 of 20 November 2003, Sec-
tion IIFor two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: COMPOUNDS TO TREAT ALZHEIMER'S DISEASE



(XV)

(57) Abstract: The present invention
is substituted amines of formula (XV)
useful in treating Alzheimer's disease
and other similar diseases.

WO 02/002506 A3

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
10 January 2002 (10.01.2002)

PCT

(10) International Publication Number
WO 02/02506 A2

(51) International Patent Classification⁷: C07C 215/00

(21) International Application Number: PCT/US01/20930

(22) International Filing Date: 29 June 2001 (29.06.2001)

(25) **Filing Language:** English

(26) **Publication Language:** English

(30) Priority Data:
60/215,323 30 June 2000 (30.06.2000) US

(71) Applicant: ELAN PHARMACEUTICALS, INC.
[US/US]; 800 Gateway Boulevard, So. San Francisco, CA
94080 (US).

(81) Designated States (national): AE, AG, AL, AM, AT, AT (utility model), AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, CZ (utility model), DE, DE (utility model), DK, DK (utility model), DM, DZ, EC, EE, FF (utility model), ES, FI, FI (utility model), GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SK (utility model), SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

— *without international search report and to be republished upon receipt of that report*


For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(72) **Inventors:** FANG, Lawrence, Y.; 1193 Beach Park Boulevard, Foster City, CA 94404 (US). JOHN, Varghese; 1772 18th Avenue, San Francisco, CA 94122 (US).

(74) Agent: DAIGNAULT, Ronald, A.; Merchant & Gould
P.C., P.O. Box 2903, Minneapolis, MN 55402-0903 (US).

WO 02/02506 A2

(54) Title: COMPOUNDS TO TREAT ALZHEIMER'S DISEASE



(57) Abstract: The present invention is substituted amines of formula (XV) useful in treating Alzheimer's disease and other similar diseases.

COMPOUNDS TO TREAT ALZHEIMER'S DISEASE

This application is being filed as a PCT International Patent Application in the name of Elan Pharmaceuticals, Inc., a U.S. national corporation and resident,
5 (Applicant for all countries), on 29 June 2001, designating all countries except US.

Background of the Invention**Field of the Invention**

10 The present invention is directed to compounds useful in treatment of Alzheimer's disease and similar diseases.

Description of the Related Art

Alzheimer's disease (AD) is a progressive degenerative disease of the brain
15 primarily associated with aging. Clinical presentation of AD is characterized by loss of memory, cognition, reasoning, judgment, and orientation. As the disease progresses, motor, sensory, and linguistic abilities are also affected until there is global impairment of multiple cognitive functions. These cognitive losses occur gradually, but typically lead to severe impairment and eventual death in the range of
20 four to twelve years.

Alzheimer's disease is characterized by two major pathologic observations in the brain: neurofibrillary tangles and beta amyloid (or neuritic) plaques, comprised predominantly of an aggregate of a peptide fragment known as A beta. Individuals with AD exhibit characteristic beta-amyloid deposits in the brain (beta amyloid
25 plaques) and in cerebral blood vessels (beta amyloid angiopathy) as well as neurofibrillary tangles. Neurofibrillary tangles occur not only in Alzheimer's disease but also in other dementia-inducing disorders. On autopsy, large numbers of these lesions are generally found in areas of the human brain important for memory and cognition.

30 Smaller numbers of these lesions in a more restricted anatomical distribution are found in the brains of most aged humans who do not have clinical AD. Amyloidogenic plaques and vascular amyloid angiopathy also characterize the brains of individuals with Trisomy 21 (Down's Syndrome), Hereditary Cerebral Hemorrhage with Amyloidosis of the Dutch-Type (HCHWA-D), and other

neurogenerative disorders. Beta-amyloid is a defining feature of AD, now believed to be a causative precursor or factor in the development of disease. Deposition of A beta in areas of the brain responsible for cognitive activities is a major factor in the development of AD. Beta-amyloid plaques are predominantly composed of amyloid beta peptide (A beta, also sometimes designated betaA4). A beta peptide is derived by proteolysis of the amyloid precursor protein (APP) and is comprised of 39-42 amino acids. Several proteases called secretases are involved in the processing of APP.

Cleavage of APP at the N-terminus of the A beta peptide by beta-secretase and at the C-terminus by one or more gamma-secretases constitutes the beta-amyloidogenic pathway, i.e. the pathway by which A beta is formed. Cleavage of APP by alpha-secretase produces alpha-sAPP, a secreted form of APP that does not result in beta-amyloid plaque formation. This alternate pathway precludes the formation of A beta peptide. A description of the proteolytic processing fragments of APP is found, for example, in U.S. Patent Nos. 5,441,870; 5,721,130; and 5,942,400.

An aspartyl protease has been identified as the enzyme responsible for processing of APP at the beta-secretase cleavage site. The beta-secretase enzyme has been disclosed using varied nomenclature, including BACE, Asp, and Memapsin. See, for example, Sinha et.al., 1999, *Nature* 402:537-554 (p501) and published PCT application WO00/17369.

Several lines of evidence indicate that progressive cerebral deposition of beta-amyloid peptide (A beta) plays a seminal role in the pathogenesis of AD and can precede cognitive symptoms by years or decades. See, for example, Selkoe, 1991, *Neuron* 6:487. Release of A beta from neuronal cells grown in culture and the presence of A beta in cerebrospinal fluid (CSF) of both normal individuals and AD patients has been demonstrated. See, for example, Seubert et al., 1992, *Nature* 359:325-327.

It has been proposed that A beta peptide accumulates as a result of APP processing by beta-secretase, thus inhibition of this enzyme's activity is desirable for the treatment of AD. *In vivo* processing of APP at the beta-secretase cleavage site is thought to be a rate-limiting step in A beta production, and is thus a

therapeutic target for the treatment of AD. See for example, Sabbagh, M., et al., 1997, *Alz. Dis. Rev.* 3, 1-19.

BACE1 knockout mice fail to produce A beta, and present a normal phenotype. When crossed with transgenic mice that overexpress APP, the progeny
5 show reduced amounts of A beta in brain extracts as compared with control animals (Luo et.al., 2001 *Nature Neuroscience* 4:231-232). This evidence further supports the proposal that inhibition of beta-secretase activity and reduction of A beta in the brain provides a therapeutic method for the treatment of AD and other beta amyloid disorders.

10 Published PCT application WO00/47618 entitled "Beta-Secretase Enzyme Compositions and Methods" identifies the beta-secretase enzyme and methods of its use. This publication also discloses oligopeptide inhibitors that bind the enzyme's active site and are useful in affinity column purification of the enzyme. In addition, WO00/77030 discloses tetrapeptide inhibitors of beta-secretase activity that are
15 based on a statine molecule.

Various pharmaceutical agents have been proposed for the treatment of Alzheimer's disease but without any real success. US Patent 5,175,281 discloses 21-aminosteroids as being useful for treating Alzheimer's disease. US Patent 5,502,187 discloses bicyclic heterocyclic amines as being useful for treating
20 Alzheimer's disease.

US Patents 4,616,088 and 4,665,193 discloses hydroxyethylamine compounds as anti-hypertensive agents due to their ability to inhibit renin.

US Patent 4, 636,491 discloses various tetrapeptides which are useful as renin inhibitors.

25 US Patent 4,749,792 discloses amino compounds useful as analgesics because of their ability to inhibit an enkephalin-degrading aminopeptidase.

US Patent 5,142,056 discloses peptide derivatives with a C₂-symmetric dihydroxyethylene core as retroviral protease inhibitors.

US Patents 5,461,067 and 5,753,652 disclose the synthesis of retroviral
30 protease inhibitors.

US Patent 5,475,138 and 5,631,405 disclose processes and various intermediates useful in the synthesis of selected protease inhibitors.

US Patent 5,502,061 discloses HIV protease inhibitors containing an unsaturated carbocycle or heterocycle at the C-terminus.

US Patent 5,545,640 discloses compounds which inhibit HIV protease activity.

US Patent 5,516,784 discloses compounds active against retroviruses, including HIV.

5 US Patent 5,602,175 discloses hydroxyethylamine compounds as retroviral protease inhibitors.

US Patent 5,631,405 discloses a process for the formation of intermediates useful in the synthesis of selected protease inhibitors.

10 US Patent 5,733,882 and International Publications WO 93/02057 and WO 93/17003 disclose dipeptide analogs as retroviral protease inhibitors.

US Patent 5,760,076 discloses hydroxyethylamino sulfonamide compounds as retrovirus protease inhibitors.

US Patent 5,807,870 discloses hydroxyethylamine compounds for the inhibition of HIV protease.

15 US Patent 5,827,891 discloses HIV protease inhibitors.

US Patent 5,830,897 discloses hydroxyethylamino sulfonamide compounds as retrovirus protease inhibitors.

US Patent 5,831,117 discloses a process and intermediates useful in retroviral protease inhibitor intermediates.

20 US Patent 5,847,169 discloses a process for preparing aminoepoxides involving the activation of the terminal hydroxyl of an aminodiol.

US Patent 5,849,911 discloses hydroxyethylamine HIV protease inhibitors which form hydrazines with one of the amino groups; this amino group must also be alkylated.

25 US Patent 5,922,770 discloses peptide derivatives which are useful in treating disorders resulting from a deficiency in growth hormone.

US Patent 6,013,658 discloses peptide derivatives which are useful in treating disorders resulting from a deficiency in growth hormone.

30 US Patent 6,022,872 discloses hydroxyethylamino sulfonyl urea compounds as HIV protease inhibitors.

US Patent 6,060,476 discloses hydroxyethylamino sulfonamide compounds as HIV protease inhibitors.

International Publication WO 89/01488 discloses renin inhibiting peptides with a hydroxyethylene or dihydroxyethylene isostere in the 10,11-position of the renin substrate angiotensinogen.

International Publication WO92/00750 discloses retroviral protease inhibitors.

International Publication WO 94/04492 discloses hydroxyethylamine intermediates useful for the treatment of retroviral diseases such as HIV. This disclosure also presents epoxides as intermediates for the retroviral inhibitors.

International Publication WO 95/06030 discloses epoxides, chloromethyl ketones, and alcohols prepared as intermediates for HIV protease inhibitors, with a single protecting group on the amine and arylalkyl side chain substituted with alkyl, nitro, nitrile, alkoxy, and thioalkoxy; a preferred side chain is 4-fluorophenylmethyl.

International publication WO98/29401 discloses a method for the preparation of aminoepoxides from aminoaldehydes by which the aminoaldehyde continuously flows into a mixing zone containing an in situ generated halomethyl organometallic reagent.

International Publication WO98/33795 discloses non-peptide inhibitors of cathepsin D.

International Publication WO98/50342 discloses bis aminomethyl carbonyl compounds as inhibitors of cysteine and serine proteases.

International Publication WO00/056335 discloses non-peptide inhibitors of aspartyl proteases. These compounds influence processing of the amyloid precursor protein APP.

EP 0 609 625 discloses HIV protease inhibitors with only one noncyclized nitrogen atom.

Bioorganic & Medicinal Chemistry Letters, 5, 721-726 (1995) describes the synthesis of compounds useful for the inhibition of HIV protease in which the C-terminal nitrogen of the hydroxyethylamine compound is incorporated into a ring system such that a piperidine ring, with a amide substituent next to the nitrogen, is formed.

The hydroxyethylamine "nucleus" or isostere, which is present in the compounds of the present invention has been employed with success in the area of HIV protease inhibition. Many of these hydroxyethylamine compounds are known as well as how to make them. See for example, *J. Am. Chem. Soc.*, 93, 288-291

(1993), *Tetrahedron Letters*, 28(45) 5569-5572 (1987), *J. Med. Chem.*, 38(4), 581-584 (1994), *Tetrahedron Letters*, 38(4), 619-620 (1997).

US Patent 5,648,511 discloses a diprotected aralkyl epoxide.

US Patents 5,482,947, 5,508,294, 5,510,349, 5,510,388, 5,521,219,
5 5,610,190, 5,639,769, 5,760,064 and 5,965,588 disclose monoprotected (substituted) aralkyl epoxides.

Tetrahedron Lett., 30(40),5425-5428 (1989) discloses a process in which doubly protected alpha-amino aldehydes are transformed into the corresponding aminoalkyl epoxides.

10 *J. Med. Chem.*, 36, 2300 (1993) discloses an azide substituted benzyl epoxide.

Tetrahedron Lett., 38, 3175 (1997) discloses a process for the preparation of N-BOC protected epoxides from protected amino acid esters.

15 *J. Med. Chem.*, 35, 2525 (1992) discloses hydroxyethylamine inhibitors of HIV protease.

US Patent 5,481,011 discloses arylalkyl amino epoxides in which the amino group is protected by a carbamate functionality.

Synlett, 6, 902 (2000) discloses the preparation of alpha-chloroketones of aminoprotected-(substituted)benzyl esters.

20 US Patent 5,648,511 discloses a diprotected aralkyl alcohol.

US Patents 5,482,947, 5,508,294, 5,510,349, 5,510,388, 5,521,219,
5,610,190, 5,639,769, 5,760,064 and 5,965,588 disclose monoprotected (substituted) aralkyl alcohols.

US Patents 5,482,947, 5,508,294, 5,510,349, 5,510,388, 5,521,219,
25 5,610,190, 5,639,769, 5,760,064 and 5,965,588 disclose a process for removing the protecting group of the monoprotected (substituted) aralkyl alcohols to give the free amino alcohol product as the amine salt.

US Patent 5,648,511 discloses the removal of the amino protecting group of a protected amino-alcohol to give a free amino-alcohol.

30 US Patent 6,150,344 discloses phosphate containing compounds useful in treating Alzheimer's disease.

EP 652 009 A1 discloses inhibitors of aspartyl protease which inhibit beta-amyloid peptide production in cell culture and *in vivo*. The compounds which

inhibit intracellular beta-amyloid peptide production are useful in treating Alzheimer's Disease.

WO00/69262 discloses a new beta-secretase and its use in assays to screen for potential drug candidates against Alzheimer's disease.

5 WO01/00663 discloses memapsin 2 (human beta-secretase) as well as catalytically active recombinant enzyme. In addition, a method of identifying inhibitors of memapsin 2, as well as two inhibitors are disclosed. Both inhibitors that are disclosed are peptides.

10 WO01/00665 discloses inhibitors of memapsin 2 that are useful in treating Alzheimer's disease.

WO01/19797 discloses lactams of the formula -C-C-CO-N-lactam-W-X-Y-Z which are useful in treating Alzheimer's disease.

EP 98/14450 and *J. Med. Chem.*, 41(18), 3387-3401 (1998) disclose aza analogs of HIV inhibitors.


15 At present there are no effective treatments for halting, preventing, or reversing the progression of Alzheimer's disease. Therefore, there is an urgent need for pharmaceutical agents capable of slowing the progression of Alzheimer's disease and/or preventing it in the first place.

20 Compounds that are effective inhibitors of beta-secretase, that inhibit beta-secretase-mediated cleavage of APP, that are effective inhibitors of A beta production, and/or are effective to reduce amyloid beta deposits or plaques, are needed for the treatment and prevention of disease characterized by amyloid beta deposits or plaques, such as AD.

25

Summary of the Invention

Disclosed is a substituted amine of formula (XV)



30

where R_1 is:

- (I) C_1-C_6 alkyl, optionally substituted with one, two or three substituents selected from the group consisting of C_1-C_3 alkyl, C_1-C_7 alkyl (optionally substituted with C_1-C_3 alkyl and C_1-C_3 alkoxy), $-F$, $-Cl$, $-Br$, $-I$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, C_1-C_3 alkoxy, $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are $-H$ or C_1-C_6 alkyl, and $-OC(=O)NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,
- (II) $-CH_2-S(O)_{0-2}-(C_1-C_6 \text{ alkyl})$,
- (III) $-CH_2-CH_2-S(O)_{0-2}-(C_1-C_6 \text{ alkyl})$,
- (IV) C_2-C_6 alkenyl with one or two double bonds, optionally substituted with one, two or three substituents selected from the group consisting of $-F$, $-Cl$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, C_1-C_3 alkoxy, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are $-H$ or C_1-C_6 alkyl,
- (V) C_2-C_6 alkynyl with one or two triple bonds, optionally substituted with one, two or three substituents selected from the group consisting of $-F$, $-Cl$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, C_1-C_3 alkoxy, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are $-H$ or C_1-C_6 alkyl,
- (VI) $-(CH_2)_{n1}-(R_{1-aryl})$ where n_1 is zero or one and where R_{1-aryl} is phenyl, 1-naphthyl, 2-naphthyl and indanyl, indenyl, dihydronaphthalyl, or tetralinyl optionally substituted with one, two, three or four of the following substituents on the aryl ring:
- (A) C_1-C_6 alkyl optionally substituted with one, two or three substituents selected from the group consisting of C_1-C_3 alkyl, $-F$, $-Cl$, $-Br$, $-I$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, and C_1-C_3 alkoxy, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,
- (B) C_2-C_6 alkenyl with one or two double bonds, optionally substituted with one, two or three substituents selected from the group consisting of $-F$, $-Cl$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, C_1-C_3 alkoxy, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are $-H$ or C_1-C_6 alkyl,
- (C) C_2-C_6 alkynyl with one or two triple bonds, optionally substituted with one, two or three substituents selected from the group consisting of $-F$, $-Cl$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, C_1-C_3 alkoxy, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are $-H$ or C_1-C_6 alkyl,

(D) -F, Cl, -Br or -I,

(F) -C₁-C₆ alkoxy optionally substituted with one, two or three of - F,

(G) -NR_{N-2}R_{N-3} where R_{N-2} and R_{N-3} are as defined below,

5 (H) -OH,

(I) -C≡N,

(J) C₃-C₇ cycloalkyl, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, -C≡N, -CF₃, C₁-C₃ alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are -H or C₁-C₆ alkyl,

10 (K) -CO-(C₁-C₄ alkyl),

(L) -SO₂-NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(M) -CO-NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

or

(N) -SO₂-(C₁-C₄ alkyl),

15 (VII) -(CH₂)_{n1}-(R_{1-heteroaryl}) where n₁ is as defined above and where R_{1-heteroaryl} is selected from the group consisting of:

pyridinyl,

pyrimidinyl,

quinolinyl,

20 benzothienyl,

indolyl,

indolinyl,

pyridazinyl,

pyrazinyl,

25 isoquinolyl,

quinazolinyl,

quinoxalinyl,

phthalazinyl,

imidazolyl,

30 isoxazolyl,

pyrazolyl,

oxazolyl,

thiazolyl,

	indoliziny1,
	indazoly1,
	benzothiazoly1,
	benzimidazoly1,
5	benzofurany1,
	furany1,
	thieny1,
	pyrroly1,
	oxadiazoly1,
10	thiadiazoly1,
	triazoly1,
	tetrazoly1,
	oxazolopyridiny1,
	imidazopyridiny1,
15	isothiazoly1,
	naphthyridiny1,
	cinnoliny1,
	carbazoly1,
	beta-carboliny1,
20	isochromany1,
	chromany1,
	tetrahydroisoquinoliny1,
	isoindoliny1,
	isobenzotetrahydrofurany1,
25	isobenzotetrahydrothieny1,
	isobenzothiény1,
	benzoxazoly1,
	pyridopyridiny1,
	benzotetrahydrofurany1,
30	benzotetrahydrothieny1,
	puriny1,
	benzodioxoly1,
	triaziny1,
	phenoxaziny1,

	phenothiazinyl,
	pteridinyl,
	benzothiazolyl,
	imidazopyridinyl,
5	imidazothiazolyl,
	dihydrobenzisoaxazinyl,
	benzisoaxazinyl,
	benzoxazinyl,
	dihydrobenzisothiazinyl,
10	benzopyranyl,
	benzothiopyranyl,
	coumarinyl,
	isocoumarinyl,
	chromonyl,
15	chromanonyl,
	pyridinyl-N-oxide,
	tetrahydroquinolinyl
	dihydroquinolinyl
	dihydroquinolinonyl
20	dihydroisoquinolinonyl
	dihydrocoumarinyl
	dihydroisocoumarinyl
	isoindolinonyl
	benzodioxanyl
25	benzoxazolinonyl
	pyrrolyl N-oxide,
	pyrimidinyl N-oxide,
	pyridazinyl N-oxide,
	pyrazinyl N-oxide,
30	quinolinyl N-oxide,
	indolyl N-oxide,
	indolinyl N-oxide,
	isoquinolyl N-oxide,
	quinazolinyl N-oxide,

quinoxaliny N-oxide,
 phthalaziny N-oxide,
 imidazolyl N-oxide,
 isoxazolyl N-oxide,
 5 oxazolyl N-oxide,
 thiazolyl N-oxide,
 indoliziny N-oxide,
 indazolyl N-oxide,
 benzothiazolyl N-oxide,
 10 benzimidazolyl N-oxide,
 pyrrolyl N-oxide,
 oxadiazolyl N-oxide,
 thiadiazolyl N-oxide,
 triazolyl N-oxide,
 15 tetrazolyl N-oxide,
 benzothiopyranyl S-oxide, and
 benzothiopyranyl S,S-dioxide,

where the $R_{1\text{-heteroaryl}}$ group is bonded to $-(CH_2)_{n1}-$ by any ring atom
 of the parent $R_{1\text{-heteroaryl}}$ group substituted by hydrogen such that the new bond to the
 20 $R_{1\text{-heteroaryl}}$ group replaces the hydrogen atom and its bond, where heteroaryl is
 optionally substituted with one, two, three or four of:

(1) C_1-C_6 alkyl optionally substituted with one, two or
 three substituents selected from the group consisting of C_1-C_3 alkyl, -F, -Cl, -Br, -I,
 -OH, -SH, $-C\equiv N$, $-CF_3$, C_1-C_3 alkoxy, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as
 25 defined above,

(2) C_2-C_6 alkenyl with one or two double bonds,
 optionally substituted with one, two or three substituents selected from the group
 consisting of -F, -Cl, -OH, -SH, $-C\equiv N$, $-CF_3$, C_1-C_3 alkoxy, and $-NR_{1-a}R_{1-b}$ where R_{1-a}
 and R_{1-b} are -H or C_1-C_6 alkyl,

(3) C_2-C_6 alkynyl with one or two triple bonds,
 optionally substituted with one, two or three substituents selected from the group
 consisting of -F, -Cl, -OH, -SH, $-C\equiv N$, $-CF_3$, C_1-C_3 alkoxy, and $-NR_{1-a}R_{1-b}$ where R_{1-a}
 and R_{1-b} are -H or C_1-C_6 alkyl,

- (4) -F, Cl, -Br or -I,
- (6) -C₁-C₆ alkoxy optionally substituted with one, two, or three of -F,
- (7) -NR_{N-2}R_{N-3} where R_{N-2} and R_{N-3} are as defined
- 5 below,
- (8) -OH,
- (9) -C≡N,
- (10) C₃-C₇ cycloalkyl, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH,
- 10 -C≡N, -CF₃, C₁-C₃ alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are -H or C₁-C₆ alkyl,
- (11) -CO-(C₁-C₄ alkyl),
- (12) -SO₂-NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,
- (13) -CO-NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined
- 15 above, or
- (14) -SO₂-(C₁-C₄ alkyl), with the proviso that when n₁ is zero R_{1-heteroaryl} is not bonded to the carbon chain by nitrogen; or
- (VIII) -(CH₂)_{n1}-(R_{1-heterocycle}) where n₁ is as defined above and R_{1-heterocycle} is selected from the group consisting of:
- 20 morpholinyl,
- thiomorpholinyl,
- thiomorpholinyl S-oxide,
- thiomorpholinyl S,S-dioxide,
- piperazinyl,
- 25 homopiperazinyl,
- pyrrolidinyl,
- pyrrolinyl,
- tetrahydropyranyl,
- piperidinyl,
- 30 tetrahydrofuranyl,
- tetrahydrothienyl,
- homopiperidinyl,
- homomorpholinyl,

homothiomorpholinyl,
 homothiomorpholinyl S,S-dioxide,
 oxazolidinonyl,
 dihydropyrazolyl,
 5 dihydropyrrolyl,
 dihydropyrazinyl,
 dihydropyridinyl,
 dihydropyrimidinyl,
 dihydrofuryl,
 10 dihydropyranyl,
 tetrahydrothienyl S-oxide,
 tetrahydrothienyl S,S-dioxide, and
 homothiomorpholinyl S-oxide,

where the $R_{1\text{-heterocycle}}$ group is bonded by any atom of the parent R_1 -
 15 heterocycle group substituted by hydrogen such that the new bond to the $R_{1\text{-heterocycle}}$
 group replaces the hydrogen atom and its bond, where heterocycle is optionally
 substituted with one, two, three or four:

(1) C_1 - C_6 alkyl optionally substituted with one, two or
 three substituents selected from the group consisting of C_1 - C_3 alkyl, -F, -Cl, -Br, -I,
 20 -OH, -SH, $-C\equiv N$, $-CF_3$, C_1 - C_3 alkoxy, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as
 defined above,

(2) C_2 - C_6 alkenyl with one or two double bonds,
 optionally substituted with one, two or three substituents selected from the group
 consisting of -F, -Cl, -OH, -SH, $-C\equiv N$, $-CF_3$, C_1 - C_3 alkoxy, and $-NR_{1-a}R_{1-b}$ where R_{1-}
 25 a and R_{1-b} are -H or C_1 - C_6 alkyl,

(3) C_2 - C_6 alkynyl with one or two triple bonds,
 optionally substituted with one, two or three substituents selected from the group
 consisting of -F, -Cl, -OH, -SH, $-C\equiv N$, $-CF_3$, C_1 - C_3 alkoxy, and $-NR_{1-a}R_{1-b}$ where R_{1-}
 a and R_{1-b} are -H or C_1 - C_6 alkyl,

30 (4) -F, Cl, -Br or -I,

(5) C_1 - C_6 alkoxy,

(6) $-C_1$ - C_6 alkoxy optionally substituted with one,

two, or three -F,

(7) $-NR_{N-2}R_{N-3}$ where R_{N-2} and R_{N-3} are as defined

below,

(8) $-OH$,

(9) $-C\equiv N$,

5 (10) C_3-C_7 cycloalkyl, optionally substituted with one, two or three substituents selected from the group consisting of $-F$, $-Cl$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, C_1-C_3 alkoxy, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are $-H$ or C_1-C_6 alkyl,

(11) $-CO-(C_1-C_4 \text{ alkyl})$,

(12) $-SO_2-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined

10 above,

(13) $-CO-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined

above,

(14) $-SO_2-(C_1-C_4 \text{ alkyl})$, or

(15) $=O$, with the proviso that when n_1 is zero R_1 -

15 heterocycle is not bonded to the carbon chain by nitrogen;

where R_2 is:

(I) $-H$,

(II) C_1-C_6 alkyl, optionally substituted with one, two or three
20 substituents selected from the group consisting of C_1-C_3 alkyl, $-F$, $-Cl$, $-Br$, $-I$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, C_1-C_3 alkoxy, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

(III) $-(CH_2)_{0-4}-R_{2-1}$ where R_{2-1} is R_{1-aryl} or $R_{1-heteroaryl}$ where R_{1-aryl} and $R_{1-heteroaryl}$ are as defined above;

25 (IV) C_2-C_6 alkenyl with one or two double bonds, optionally substituted with one, two or three substituents selected from the group consisting of $-F$, $-Cl$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, C_1-C_3 alkoxy, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are $-H$ or C_1-C_6 alkyl,

(V) C_2-C_6 alkynyl with one or two triple bonds, optionally substituted
30 with one, two or three substituents selected from the group consisting of $-F$, $-Cl$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, C_1-C_3 alkoxy, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are $-H$ or C_1-C_6 alkyl, or

(VI) $-(CH_2)_{0-4}-C_3-C_7$ cycloalkyl, optionally substituted with one, two or three substituents selected from the group consisting of $-F$, $-Cl$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, C_1-C_3 alkoxy, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are $-H$ or C_1-C_6 alkyl;

5 where R_3 is selected from the group consisting of:

(I) $-H$,

(II) C_1-C_6 alkyl, optionally substituted with one, two or three substituents selected from the group consisting of C_1-C_3 alkyl, $-F$, $-Cl$, $-Br$, $-I$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, C_1-C_3 alkoxy, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined

10 above,

(III) $-(CH_2)_{0-4}-R_{2-1}$ where R_{2-1} is R_{1-aryl} or $R_{1-heteroaryl}$ where R_{1-aryl} and $R_{1-heteroaryl}$ are as defined above

(IV) C_2-C_6 alkenyl with one or two double bonds,

(V) C_2-C_6 alkynyl with one or two triple bonds; or

15 (VI) $-(CH_2)_{0-4}-C_3-C_7$ cycloalkyl, optionally substituted with one, two or three substituents selected from the group consisting of $-F$, $-Cl$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, C_1-C_3 alkoxy, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are $-H$ or C_1-C_6 alkyl,

and where R_2 and R_3 are taken together with the carbon to which they are attached to form a carbocycle of three, four, five, six, or seven carbon atoms,

20 optionally where one carbon atom is replaced by a heteroatom selected from the group consisting of $-O-$, $-S-$, $-SO_2-$, and $-NR_{N-2}-$, where R_{N-2} is as defined below;

where R_N is:

(I) $R_{N-1}-X_N-$ where X_N is selected from the group consisting of:

25 (A) $-CO-$,

(B) $-SO_2-$,

(C) $-(CR'R'')_{1-6}$ where R' and R'' are the same or different and are $-H$ and C_1-C_4 alkyl,

(D) $-CO-(CR'R'')_{1-6}-X_{N-1}$ where X_{N-1} is selected from the group consisting of $-O-$, $-S-$ and $-NR'-$ and where R' and R'' are as defined above, and

(E) a single bond;

where R_{N-1} is selected from the group consisting of:

(A) $R_{N\text{-aryl}}$ where $R_{N\text{-aryl}}$ is phenyl, 1-naphthyl, 2-naphthyl, tetralinyl, indanyl, 6,7,8,9-tetrahydro-5H-benzo[a]cycloheptenyl, or dihydronaphthyl optionally substituted with one, two or three of the following substituents which can be the same or different and are:

- 5 (1) $C_1\text{-}C_6$ alkyl, optionally substituted with one, two or three substituents selected from the group consisting of $C_1\text{-}C_3$ alkyl, -F, -Cl, -Br, -I, -OH, -SH, $-C\equiv N$, $-CF_3$, $C_1\text{-}C_3$ alkoxy, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,
- (2) -OH,
- 10 (3) $-NO_2$,
- (4) -F, -Cl, -Br, -I,
- (5) -CO-OH,
- (6) $-C\equiv N$,
- (7) $-(CH_2)_{0-4}\text{-CO-NR}_{N-2}R_{N-3}$ where R_{N-2} and R_{N-3} are
- 15 the same or different and are selected from the group consisting of:
 - (a) -H,
 - (b) $-C_1\text{-}C_6$ alkyl optionally substituted with one substituent selected from the group consisting of:
 - (i) -OH, and
 - 20 (ii) $-NH_2$,
 - (c) $-C_1\text{-}C_6$ alkyl optionally substituted with one, two, or three -F, -Cl, -Br, or -I,
 - (d) $-C_3\text{-}C_7$ cycloalkyl,
 - (e) $-(C_1\text{-}C_2\text{ alkyl})\text{-}(C_3\text{-}C_7\text{ cycloalkyl})$,
 - 25 (f) $-(C_1\text{-}C_6\text{ alkyl})\text{-O}\text{-}(C_1\text{-}C_3\text{ alkyl})$,
 - (g) $-C_2\text{-}C_6$ alkenyl with one or two double bonds,
 - (h) $-C_2\text{-}C_6$ alkynyl with one or two triple bonds,
 - 30 (i) $-C_1\text{-}C_6$ alkyl chain with one double bond and one triple bond,
 - (j) $-R_{1\text{-aryl}}$ where $R_{1\text{-aryl}}$ is as defined above, and

(k) $-R_{1\text{-heteroaryl}}$ where $R_{1\text{-heteroaryl}}$ is as defined

above,

(8) $-(CH_2)_{0-4}-CO-(C_1-C_{12} \text{ alkyl})$,

(9) $-(CH_2)_{0-4}-CO-(C_2-C_{12} \text{ alkenyl with one, two or}$

5 three double bonds),

(10) $-(CH_2)_{0-4}-CO-(C_2-C_{12} \text{ alkynyl with one, two or}$

three triple bonds),

(11) $-(CH_2)_{0-4}-CO-(C_3-C_7 \text{ cycloalkyl})$,

(12) $-(CH_2)_{0-4}-CO-R_{1\text{-aryl}}$ where $R_{1\text{-aryl}}$ is as defined

10 above,

(13) $-(CH_2)_{0-4}-CO-R_{1\text{-heteroaryl}}$ where $R_{1\text{-heteroaryl}}$ is as

defined above,

(14) $-(CH_2)_{0-4}-CO-R_{1\text{-heterocycle}}$ where $R_{1\text{-heterocycle}}$ is as

defined above,

15 (15) $-(CH_2)_{0-4}-CO-R_{N-4}$ where R_{N-4} is selected from

the group consisting of morpholinyl, thiomorpholinyl, piperazinyl, piperidinyl, homomorpholinyl, homothiomorpholinyl, homothiomorpholinyl S-oxide, homothiomorpholinyl S,S-dioxide, pyrrolinyl and pyrrolidinyl where each group is optionally substituted with one, two, three, or four of C_1-C_6 alkyl,

20 (16) $-(CH_2)_{0-4}-CO-O-R_{N-5}$ where R_{N-5} is

selected from the group consisting of:

(a) C_1-C_6 alkyl,

(b) $-(CH_2)_{0-2}-(R_{1\text{-aryl}})$ where $R_{1\text{-aryl}}$ is as defined

above,

25 (c) C_2-C_6 alkenyl containing one or two double

bonds,

(d) C_2-C_6 alkynyl containing one or two triple

bonds,

(e) C_3-C_7 cycloalkyl, and

30 (f) $-(CH_2)_{0-2}-(R_{1\text{-heteroaryl}})$ where $R_{1\text{-heteroaryl}}$ is as

defined above,

(17) $-(CH_2)_{0-4}-SO_2-NR_{N-2}R_{N-3}$ where R_{N-2} and R_{N-3} are

as defined above,

(18) $-(CH_2)_{0-4}-SO-(C_1-C_8 \text{ alkyl})$,

(19) $-(\text{CH}_2)_{0-4}-\text{SO}_2-(\text{C}_1-\text{C}_{12} \text{ alkyl})$,

(20) $-(\text{CH}_2)_{0-4}-\text{SO}_2-(\text{C}_3-\text{C}_7 \text{ cycloalkyl})$,

(21) $-(\text{CH}_2)_{0-4}-\text{N}(\text{H or } \text{R}_{\text{N-5}})-\text{CO}-\text{O}-\text{R}_{\text{N-5}}$ where $\text{R}_{\text{N-5}}$

can be the same or different and is as defined above,

5 (22) $-(\text{CH}_2)_{0-4}-\text{N}(\text{H or } \text{R}_{\text{N-5}})-\text{CO}-\text{N}(\text{R}_{\text{N-5}})_2$, where $\text{R}_{\text{N-5}}$

can be the same or different and is as defined above,

(23) $-(\text{CH}_2)_{0-4}-\text{N}-\text{CS}-\text{N}(\text{R}_{\text{N-5}})_2$, where $\text{R}_{\text{N-5}}$ can be the

same or different and is as defined above,

(24) $-(\text{CH}_2)_{0-4}-\text{N}(-\text{H or } \text{R}_{\text{N-5}})-\text{CO}-\text{R}_{\text{N-2}}$ where $\text{R}_{\text{N-5}}$ and

10 $\text{R}_{\text{N-2}}$ can be the same or different and are as defined above,

(25) $-(\text{CH}_2)_{0-4}-\text{NR}_{\text{N-2}}\text{R}_{\text{N-3}}$ where $\text{R}_{\text{N-2}}$ and $\text{R}_{\text{N-3}}$ can be

the same or different and are as defined above,

(26) $-(\text{CH}_2)_{0-4}-\text{R}_{\text{N-4}}$ where $\text{R}_{\text{N-4}}$ is as defined above,

(27) $-(\text{CH}_2)_{0-4}-\text{O}-\text{CO}-(\text{C}_1-\text{C}_6 \text{ alkyl})$,

15 (28) $-(\text{CH}_2)_{0-4}-\text{O}-\text{P}(\text{O})-(\text{OR}_{\text{N-aryl-1}})_2$ where $\text{R}_{\text{N-aryl-1}}$ is –

H or $\text{C}_1-\text{C}_4 \text{ alkyl}$,

(29) $-(\text{CH}_2)_{0-4}-\text{O}-\text{CO}-\text{N}(\text{R}_{\text{N-5}})_2$ where $\text{R}_{\text{N-5}}$ is as

defined above,

(30) $-(\text{CH}_2)_{0-4}-\text{O}-\text{CS}-\text{N}(\text{R}_{\text{N-5}})_2$ where $\text{R}_{\text{N-5}}$ is as defined

20 above,

(31) $-(\text{CH}_2)_{0-4}-\text{O}-(\text{R}_{\text{N-5}})_2$ where $\text{R}_{\text{N-5}}$ is as defined

above,

(32) $-(\text{CH}_2)_{0-4}-\text{O}-(\text{R}_{\text{N-5}})_2-\text{COOH}$ where $\text{R}_{\text{N-5}}$ is as

defined above,

25 (33) $-(\text{CH}_2)_{0-4}-\text{S}-(\text{R}_{\text{N-5}})_2$ where $\text{R}_{\text{N-5}}$ is as defined

above,

(34) $-(\text{CH}_2)_{0-4}-\text{O}-(\text{C}_1-\text{C}_6 \text{ alkyl optionally substituted}$

with one, two, three, four, or five of $-\text{F}$),

(35) $\text{C}_3-\text{C}_7 \text{ cycloalkyl}$,

30 (36) $\text{C}_2-\text{C}_6 \text{ alkenyl}$ with one or two double bonds

optionally substituted with $\text{C}_1-\text{C}_3 \text{ alkyl}$, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, $-\text{OH}$, $-\text{SH}$, $-\text{C}\equiv\text{N}$, $-\text{CF}_3$, C_1-

$\text{C}_3 \text{ alkoxy}$, or $-\text{NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

(37) C₂-C₆ alkynyl with one or two triple bonds

optionally substituted with C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₃ alkoxy, or -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(38) -(CH₂)₀₋₄-N(-H or R_{N-5})-SO₂-R_{N-2} where R_{N-5} and
5 R_{N-2} can be the same or different and are as described above, or

(39) -(CH₂)₀₋₄-C₃-C₇ cycloalkyl,

(B) -R_{N-heteroaryl} where R_{N-heteroaryl} is selected from the group
consisting of:

10 pyridinyl,
pyrimidinyl,
quinolinyl,
benzothienyl,
indolyl,
indolinyl,
15 pyridazinyl,
pyrazinyl,
isoindolyl,
isoquinolyl,
quinazolinyl,
20 quinoxalinyl,
phthalazinyl,
imidazolyl,
isoxazolyl,
pyrazolyl,
25 oxazolyl,
thiazolyl,
indolizinyl,
indazolyl,
benzothiazolyl,
30 benzimidazolyl,
benzofuranyl,
furanyl,
thienyl,

	pyrrolyl,
	oxadiazolyl,
	thiadiazolyl,
	triazolyl,
5	tetrazolyl,
	oxazolopyridinyl,
	imidazopyridinyl,
	isothiazolyl,
	naphthyridinyl,
10	cinolyl,
	carbazolyl,
	beta-carbolinyl,
	isochromanyl,
	chromanyl,
15	tetrahydroisoquinolyl,
	isoindolyl,
	isobenzotetrahydrofuranyl,
	isobenzotetrahydrothienyl,
	isobenzothienyl,
20	benzoxazolyl,
	pyridopyridinyl,
	benzotetrahydrofuranyl,
	benzotetrahydrothienyl,
	purinyl,
25	benzodioxolyl,
	triazinyl,
	phenoxazinyl,
	phenothiazinyl,
	pteridinyl,
30	benzothiazolyl,
	imidazopyridinyl,
	imidazothiazolyl,
	dihydrobenzisoxazinyl,
	benzisoxazinyl,

	benzoxazinylyl,
	dihydrobenziso-thiazinylyl,
	benzopyranylyl,
	benzothiopyranylyl,
5	coumarinylyl,
	isocoumarinylyl,
	chromonylyl,
	chromanonylyl,
	pyridinylyl-N-oxide,
10	tetrahydroquinolinylyl,
	dihydroquinolinylyl,
	dihydroquinolinonylyl,
	dihydroisoquinolinonylyl,
	dihydrocoumarinylyl,
15	dihydroisocoumarinylyl,
	isoindolinonylyl,
	benzodioxanylyl,
	benzoxazolinonylyl,
	pyrrolylyl N-oxide,
20	pyrimidinyl N-oxide,
	pyridazinylyl N-oxide,
	pyrazinylyl N-oxide,
	quinolinylyl N-oxide,
	indolyl N-oxide,
25	indolinylyl N-oxide,
	isoquinolyl N-oxide,
	quinazolinyl N-oxide,
	quinoxalinylyl N-oxide,
	phthalazinylyl N-oxide,
30	imidazolyl N-oxide,
	isoxazolyl N-oxide,
	oxazolyl N-oxide,
	thiazolyl N-oxide,
	indolizinylyl N-oxide,

indazolyl N-oxide,
 benzothiazolyl N-oxide,
 benzimidazolyl N-oxide,
 pyrrolyl N-oxide,
 5 oxadiazolyl N-oxide,
 thiadiazolyl N-oxide,
 triazolyl N-oxide,
 tetrazolyl N-oxide,
 benzothiopyranyl S-oxide, and
 10 benzothiopyranyl S,S-dioxide

where the $R_{N\text{-heteroaryl}}$ group is bonded by any atom of the parent $R_{N\text{-heteroaryl}}$ group substituted by hydrogen such that the new bond to the $R_{N\text{-heteroaryl}}$ group replaces the hydrogen atom and its bond, where heteroaryl is optionally substituted with one, two, three, or four of:

15 (1) $C_1\text{-}C_6$ alkyl, optionally substituted with one, two or three substituents selected from the group consisting of $C_1\text{-}C_3$ alkyl, -F, -Cl, -Br, -I, -OH, -SH, $-C\equiv N$, $-CF_3$, $C_1\text{-}C_3$ alkoxy, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

(2) -OH,
 20 (3) $-NO_2$,
 (4) -F, -Cl, -Br, or -I,
 (5) -CO-OH,
 (6) $-C\equiv N$,
 (7) $-(CH_2)_{0-4}\text{-CO-NR}_{N-2}R_{N-3}$ where R_{N-2} and R_{N-3} are
 25 the same or different and are selected from the group consisting of:

(a) -H,
 (b) $-C_1\text{-}C_6$ alkyl optionally substituted with one substituent selected from the group consisting of:
 (i) -OH, and
 30 (ii) $-NH_2$,
 (c) $-C_1\text{-}C_6$ alkyl optionally substituted with one, two, or three -F, -Cl, -Br, -I,
 (d) $-C_3\text{-}C_7$ cycloalkyl,

- (e) $-(C_1-C_2 \text{ alkyl})-(C_3-C_7 \text{ cycloalkyl})$,
 (f) $-(C_1-C_6 \text{ alkyl})-O-(C_1-C_3 \text{ alkyl})$,
 (g) $-C_2-C_6 \text{ alkenyl}$ with one or two double bonds,
 5 (h) $-C_2-C_6 \text{ alkynyl}$ with one or two triple bonds,
 (i) $-C_1-C_6 \text{ alkyl chain}$ with one double bond and one triple bond,
 (j) $-R_{1-aryl}$ where R_{1-aryl} is as defined above,
 10 (k) $-R_{1-heteroaryl}$ where $R_{1-heteroaryl}$ is as defined above,
 (8) $-(CH_2)_{0-4}-CO-(C_1-C_{12} \text{ alkyl})$,
 (9) $-(CH_2)_{0-4}-CO-(C_2-C_{12} \text{ alkenyl}$ with one, two or three double bonds),
 15 (10) $-(CH_2)_{0-4}-CO-(C_2-C_{12} \text{ alkynyl}$ with one, two or three triple bonds),
 (11) $-(CH_2)_{0-4}-CO-(C_3-C_7 \text{ cycloalkyl})$,
 (12) $-(CH_2)_{0-4}-CO-R_{1-aryl}$ where R_{1-aryl} is as defined above,
 20 (13) $-(CH_2)_{0-4}-CO-R_{1-heteroaryl}$ where $R_{1-heteroaryl}$ is as defined above,
 (14) $-(CH_2)_{0-4}-CO-R_{1-heterocycle}$ where $R_{1-heterocycle}$ is as defined above,
 (15) $-(CH_2)_{0-4}-CO-R_{N-4}$ where R_{N-4} is selected from
 25 the group consisting of morpholinyl, thiomorpholinyl, piperazinyl, piperidinyl, homomorpholinyl, homothiomorpholinyl, homothiomorpholinyl S-oxide, homothiomorpholinyl S,S-dioxide, pyrrolinyl and pyrrolidinyl where each group is optionally substituted with one, two, three, or four of $C_1-C_6 \text{ alkyl}$,
 (16) $-(CH_2)_{0-4}-CO-O-R_{N-5}$ where R_{N-5} is selected from
 30 the group consisting of:
 (a) $C_1-C_6 \text{ alkyl}$,
 (b) $-(CH_2)_{0-2}-(R_{1-aryl})$ where R_{1-aryl} is as defined above,

- bonds,
- (c) C₂-C₆ alkenyl containing one or two double bonds,
- (d) C₂-C₆ alkynyl containing one or two triple bonds,
- 5 (e) C₃-C₇ cycloalkyl, and
- (f) -(CH₂)₀₋₂-(R₁-heteroaryl) where R₁-heteroaryl is as defined above,
- (17) -(CH₂)₀₋₄-SO₂-NR_{N-2}R_{N-3} where R_{N-2} and R_{N-3} are as defined above,
- 10 (18) -(CH₂)₀₋₄-SO-(C₁-C₈ alkyl),
- (19) -(CH₂)₀₋₄-SO₂-(C₁-C₁₂ alkyl),
- (20) -(CH₂)₀₋₄-SO₂-(C₃-C₇ cycloalkyl),
- (21) -(CH₂)₀₋₄-N(H or R_{N-5})-CO-O-R_{N-5} where R_{N-5} can be the same or different and is as defined above,
- 15 (22) -(CH₂)₀₋₄-N(H or R_{N-5})-CO-N(R_{N-5})₂, where R_{N-5} can be the same or different and is as defined above,
- (23) -(CH₂)₀₋₄-N-CS-N(R_{N-5})₂, where R_{N-5} can be the same or different and is as defined above,
- (24) -(CH₂)₀₋₄-N(-H or R_{N-5})-CO-R_{N-2} where R_{N-5} and R_{N-2} can be the same or different and are as defined above,
- 20 (25) -(CH₂)₀₋₄-NR_{N-2}R_{N-3} where R_{N-2} and R_{N-3} can be the same or different and are as defined above,
- (26) -(CH₂)₀₋₄-R_{N-4} where R_{N-4} is as defined above,
- (27) -(CH₂)₀₋₄-O-CO-(C₁-C₆ alkyl),
- 25 (28) -(CH₂)₀₋₄-O-P(O)-(OR_{N-aryl-1})₂ where R_{N-aryl-1} is -H or C₁-C₄ alkyl,
- (29) -(CH₂)₀₋₄-O-CO-N(R_{N-5})₂ where R_{N-5} is as defined above,
- (30) -(CH₂)₀₋₄-O-CS-N(R_{N-5})₂ where R_{N-5} is as defined above,
- 30 (31) -(CH₂)₀₋₄-O-(R_{N-5})₂ where R_{N-5} is as defined above,
- (32) -(CH₂)₀₋₄-O-(R_{N-5})₂-COOH where R_{N-5} is as defined above,

(33) $-(\text{CH}_2)_{0-4}-\text{S}-(\text{R}_{\text{N}-5})_2$ where $\text{R}_{\text{N}-5}$ is as defined

above,

(34) $-(\text{CH}_2)_{0-4}-\text{O}-(\text{C}_1-\text{C}_6 \text{ alkyl optionally substituted with one, two, three, four, or five of } -\text{F}),$

5 (35) $\text{C}_3-\text{C}_7 \text{ cycloalkyl},$

(36) $\text{C}_2-\text{C}_6 \text{ alkenyl with one or two double bonds optionally substituted with } \text{C}_1-\text{C}_3 \text{ alkyl, } -\text{F, } -\text{Cl, } -\text{Br, } -\text{I, } -\text{OH, } -\text{SH, } -\text{C}\equiv\text{N, } -\text{CF}_3, \text{ C}_1-\text{C}_3 \text{ alkoxy, or } -\text{NR}_{1-a}\text{R}_{1-b} \text{ where } \text{R}_{1-a} \text{ and } \text{R}_{1-b} \text{ are as defined above,}$

(37) $\text{C}_2-\text{C}_6 \text{ alkynyl with one or two triple bonds optionally substituted with } \text{C}_1-\text{C}_3 \text{ alkyl, } -\text{F, } -\text{Cl, } -\text{Br, } -\text{I, } -\text{OH, } -\text{SH, } -\text{C}\equiv\text{N, } -\text{CF}_3, \text{ C}_1-\text{C}_3 \text{ alkoxy, or } -\text{NR}_{1-a}\text{R}_{1-b} \text{ where } \text{R}_{1-a} \text{ and } \text{R}_{1-b} \text{ are as defined above,}$

(38) $-(\text{CH}_2)_{0-4}-\text{N}(-\text{H or } \text{R}_{\text{N}-5})-\text{SO}_2-\text{R}_{\text{N}-2}$ where $\text{R}_{\text{N}-5}$ and $\text{R}_{\text{N}-2}$ can be the same or different and are as defined above, or

(39) $-(\text{CH}_2)_{0-4}-\text{C}_3-\text{C}_7 \text{ cycloalkyl},$

15 (C) $\text{R}_{\text{N-aryl}}-\text{W}-\text{R}_{\text{N-aryl}},$ where $\text{R}_{\text{N-aryl}},$ and $\text{R}_{\text{N-aryl}}$ are as defined above,

(D) $\text{R}_{\text{N-aryl}}-\text{W}-\text{R}_{\text{N-heteroaryl}},$ where $\text{R}_{\text{N-aryl}},$ and $\text{R}_{\text{N-heteroaryl}},$ are as defined above,

(E) $\text{R}_{\text{N-aryl}}-\text{W}-\text{R}_{\text{N-1-heterocycle}},$ wherein $\text{R}_{\text{N-1-heterocycle}}$ is the same as $\text{R}_{1-\text{heterocycle}},$ and $\text{R}_{1-\text{heterocycle}}$ is as defined above

(F) $\text{R}_{\text{N-heteroaryl}}-\text{W}-\text{R}_{\text{N-aryl}},$ where $\text{R}_{\text{N-aryl}},$ and $\text{R}_{\text{N-heteroaryl}},$ are as defined above,

(G) $\text{R}_{\text{N-heteroaryl}}-\text{W}-\text{R}_{\text{N-heteroaryl}},$ where $\text{R}_{\text{N-heteroaryl}}$ is as defined above,

25 (H) $\text{R}_{\text{N-heteroaryl}}-\text{W}-\text{R}_{\text{N-1-heterocycle}},$ where $\text{R}_{\text{N-heteroaryl}},$ and $\text{R}_{\text{N-1-heterocycle}},$ are as defined above,

(I) $\text{R}_{\text{N-heterocycle}}-\text{W}-\text{R}_{\text{N-aryl}},$ wherein $\text{R}_{\text{N-heterocycle}}$ is the same as $\text{R}_{1-\text{heterocycle}},$ and $\text{R}_{1-\text{heterocycle}}$ is as defined above, and $\text{R}_{\text{N-aryl}}$ is as defined above,

(J) $\text{R}_{\text{N-heterocycle}}-\text{W}-\text{R}_{\text{N-heteroaryl}},$ where $\text{R}_{\text{N-heteroaryl}},$ and $\text{R}_{\text{N-heterocycle}},$ are as defined above, and

(K) $\text{R}_{\text{N-heterocycle}}-\text{W}-\text{R}_{\text{N-1-heterocycle}},$ where $\text{R}_{\text{N-heterocycle}},$ and $\text{R}_{\text{N-1-heterocycle}},$ are as defined above,

where W is

- (1) $-(\text{CH}_2)_{0-4}-$,
 (2) $-\text{O}-$,
 (3) $-\text{S}(\text{O})_{0-2}-$,
 (4) $-\text{N}(\text{R}_{\text{N-5}})-$ where $\text{R}_{\text{N-5}}$ is as defined above, or
 (5) $-\text{CO}-$;

(II) $-\text{CO}-(\text{C}_1-\text{C}_{10} \text{ alkyl})$ where alkyl is optionally substituted with one three substituents selected from the group consisting of:

- (A) $-\text{OH}$,
 (B) $-\text{C}_1-\text{C}_6$ alkoxy,
 (C) $-\text{C}_1-\text{C}_6$ thioalkoxy,
 (D) $-\text{CO}-\text{O}-\text{R}_{\text{N-8}}$ where $\text{R}_{\text{N-8}}$ is $-\text{H}$, C_1-C_6 alkyl or $-\text{phenyl}$,
 (E) $-\text{CO}-\text{NR}_{\text{N-2}}\text{R}_{\text{N-3}}$ where $\text{R}_{\text{N-2}}$ and $\text{R}_{\text{N-3}}$ are the same or different and are as defined above,
 (F) $-\text{CO}-\text{R}_{\text{N-4}}$ where $\text{R}_{\text{N-4}}$ is as defined above,
 (G) $-\text{SO}_2-(\text{C}_1-\text{C}_8 \text{ alkyl})$,
 (H) $-\text{SO}_2-\text{NR}_{\text{N-2}}\text{R}_{\text{N-3}}$ where $\text{R}_{\text{N-2}}$ and $\text{R}_{\text{N-3}}$ are the same or different and are as defined above,
 (I) $-\text{NH}-\text{CO}-(\text{C}_1-\text{C}_6 \text{ alkyl})$,
 (J) $-\text{NH}-\text{CO}-\text{O}-\text{R}_{\text{N-8}}$ where $\text{R}_{\text{N-8}}$ is as defined above,
 (K) $-\text{NR}_{\text{N-2}}\text{R}_{\text{N-3}}$ where $\text{R}_{\text{N-2}}$ and $\text{R}_{\text{N-3}}$ are the same or different and are as defined above,
 (L) $-\text{R}_{\text{N-4}}$ where $\text{R}_{\text{N-4}}$ is as defined above,
 (M) $-\text{O}-\text{CO}-(\text{C}_1-\text{C}_6 \text{ alkyl})$,
 (N) $-\text{O}-\text{CO}-\text{NR}_{\text{N-8}}\text{R}_{\text{N-8}}$ where $\text{R}_{\text{N-8}}$ are the same or different and are as defined above,
 (O) $-\text{O}-(\text{C}_1-\text{C}_5 \text{ alkyl})-\text{COOH}$,
 (P) $-\text{O}-(\text{C}_1-\text{C}_6 \text{ alkyl})$ optionally substituted with one, two, or three of $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$,
 (Q) $-\text{NH}-\text{SO}_2-(\text{C}_1-\text{C}_6 \text{ alkyl})$, and
 (R) $-\text{F}$, or $-\text{Cl}$,

(III) $-\text{CO}-(\text{C}_1-\text{C}_6 \text{ alkyl})-\text{O}-(\text{C}_1-\text{C}_6 \text{ alkyl})$ where alkyl is optionally substituted with one, two, or three of substituents selected from the group consisting of:

- (A) -OH,
(B) -C₁-C₆ alkoxy,
(C) -C₁-C₆ thioalkoxy,
(D) -CO-O-R_{N-8} where R_{N-8} is -H, C₁-C₆ alkyl or -phenyl,
5 (E) -CO-NR_{N-2}R_{N-3} where R_{N-2} and R_{N-3} are the same or different and are as defined above,
(F) -CO-R_{N-4} where R_{N-4} is as defined above,
(G) -SO₂-(C₁-C₈ alkyl),
(H) -SO₂-NR_{N-2}R_{N-3} where R_{N-2} and R_{N-3} are the same or
10 different and are as defined above,
(I) -NH-CO-(C₁-C₆ alkyl),
(J) -NH-CO-O-R_{N-8} where R_{N-8} is as defined above,
(K) -NR_{N-2}R_{N-3} where R_{N-2} and R_{N-3} are the same or different and are as defined above,
15 (L) -R_{N-4} where R_{N-4} is as defined above,
(M) -O-CO-(C₁-C₆ alkyl),
(N) -O-CO-NR_{N-8}R_{N-8} where R_{N-8} are the same or different and are as defined above,
(O) -O-(C₁-C₅ alkyl)-COOH,
20 (P) -O-(C₁-C₆ alkyl optionally substituted with one, two, or three of -F, -Cl, -Br, or -I),
(Q) -NH-SO₂-(C₁-C₆ alkyl), and
(R) -F, or -Cl,
(IV) -CO-(C₁-C₆ alkyl)-S-(C₁-C₆ alkyl) where alkyl is optionally
25 substituted with one, two, or three of substituents selected from the group consisting of:
(A) -OH,
(B) -C₁-C₆ alkoxy,
(C) -C₁-C₆ thioalkoxy,
30 (D) -CO-O-R_{N-8} where R_{N-8} is as defined above,
(E) -CO-NR_{N-2}R_{N-3} where R_{N-2} and R_{N-3} are the same or different and are as defined above,
(F) -CO-R_{N-4} where R_{N-4} is as defined above,

(G) $-\text{SO}_2-(\text{C}_1-\text{C}_8 \text{ alkyl})$,

(H) $-\text{SO}_2-\text{NR}_{\text{N}-2}\text{R}_{\text{N}-3}$ where $\text{R}_{\text{N}-2}$ and $\text{R}_{\text{N}-3}$ are the same or different and are as defined above,

(I) $-\text{NH}-\text{CO}-(\text{C}_1-\text{C}_6 \text{ alkyl})$,

5 (J) $-\text{NH}-\text{CO}-\text{O}-\text{R}_{\text{N}-8}$ where $\text{R}_{\text{N}-8}$ is as defined above,

(K) $-\text{NR}_{\text{N}-2}\text{R}_{\text{N}-3}$ where $\text{R}_{\text{N}-2}$ and $\text{R}_{\text{N}-3}$ are the same or different and are as defined above,

(L) $-\text{R}_{\text{N}-4}$ where $\text{R}_{\text{N}-4}$ is as defined above,

(M) $-\text{O}-\text{CO}-(\text{C}_1-\text{C}_6 \text{ alkyl})$,

10 (N) $-\text{O}-\text{CO}-\text{NR}_{\text{N}-8}\text{R}_{\text{N}-8}$ where $\text{R}_{\text{N}-8}$ are the same or different and are as defined above,

(O) $-\text{O}-(\text{C}_1-\text{C}_5 \text{ alkyl})-\text{COOH}$,

(P) $-\text{O}-(\text{C}_1-\text{C}_6 \text{ alkyl})$ optionally substituted with one, two, or three of $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, or $-\text{I}$,

15 (Q) $-\text{NH}-\text{SO}_2-(\text{C}_1-\text{C}_6 \text{ alkyl})$, and

(R) $-\text{F}$, or $-\text{Cl}$,

(V) $-\text{CO}-\text{CH}(-(\text{CH}_2)_{0-2}-\text{O}-\text{R}_{\text{N}-10})-(\text{CH}_2)_{0-2}-\text{R}_{\text{N-aryl}}/\text{R}_{\text{N-heteroaryl}}$ where $\text{R}_{\text{N-aryl}}$ and $\text{R}_{\text{N-heteroaryl}}$ are as defined above, where $\text{R}_{\text{N}-10}$ is selected from the group consisting of:

20 (A) $-\text{H}$,

(B) $\text{C}_1-\text{C}_6 \text{ alkyl}$,

(C) $\text{C}_3-\text{C}_7 \text{ cycloalkyl}$,

(D) $\text{C}_2-\text{C}_6 \text{ alkenyl}$ with one double bond,

(E) $\text{C}_2-\text{C}_6 \text{ alkynyl}$ with one triple bond,

25 (F) $\text{R}_{1-\text{aryl}}$ where $\text{R}_{1-\text{aryl}}$ is as defined above, and

(G) $\text{R}_{\text{N-heteroaryl}}$ where $\text{R}_{\text{N-heteroaryl}}$ is as defined above, or

(VI) $-\text{CO}-(\text{C}_3-\text{C}_8 \text{ cycloalkyl})$ where alkyl is optionally substituted with one or two substituents selected from the group consisting of:

(A) $-(\text{CH}_2)_{0-4}-\text{OH}$,

30 (B) $-(\text{CH}_2)_{0-4}-\text{C}_1-\text{C}_6 \text{ alkoxy}$,

(C) $-(\text{CH}_2)_{0-4}-\text{C}_1-\text{C}_6 \text{ thioalkoxy}$,

(D) $-(\text{CH}_2)_{0-4}-\text{CO}-\text{O}-\text{R}_{\text{N}-8}$ where $\text{R}_{\text{N}-8}$ is $-\text{H}$, $\text{C}_1-\text{C}_6 \text{ alkyl}$ or phenyl,

(E) $-(CH_2)_{0-4}-CO-NR_{N-2}R_{N-3}$ where R_{N-2} and R_{N-3} are the same or different and are as defined above,

(F) $-(CH_2)_{0-4}-CO-R_{N-4}$ where R_{N-4} is as defined above,

(G) $-(CH_2)_{0-4}-SO_2-(C_1-C_8 \text{ alkyl})$,

5 (H) $-(CH_2)_{0-4}-SO_2-NR_{N-2}R_{N-3}$ where R_{N-2} and R_{N-3} are the same or different and are as defined above,

(I) $-(CH_2)_{0-4}-NH-CO-(C_1-C_6 \text{ alkyl})$,

(J) $-NH-CO-O-R_{N-8}$ where R_{N-8} is as defined above,

(K) $-(CH_2)_{0-4}-NR_{N-2}R_{N-3}$ where R_{N-2} and R_{N-3} are the same or
10 different and are as defined above,

(L) $-(CH_2)_{0-4}-R_{N-4}$ where R_{N-4} is as defined above,

(M) $-O-CO-(C_1-C_6 \text{ alkyl})$,

(N) $-O-CO-NR_{N-8}R_{N-8}$ where R_{N-8} are the same or different and are as defined above,

15 (O) $-O-(C_1-C_5 \text{ alkyl})-COOH$,

(P) $-O-(C_1-C_6 \text{ alkyl})$ optionally substituted with one, two, or three of $-F$, $-Cl$, $-Br$, or $-I$,

(Q) $-NH-SO_2-(C_1-C_6 \text{ alkyl})$, and

(R) $-F$, or $-Cl$;

20

where R_A is:

(I) $-C_1-C_{10}$ alkyl optionally substituted with one, two or three substituents selected from the group consisting of C_1-C_3 alkyl, $-F$, $-Cl$, $-Br$, $-I$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, C_1-C_6 alkoxy, $-O$ -phenyl, $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as
25 defined above, $-OC=O NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined above, $-S(=O)_{0-2} R_{1-a}$ where R_{1-a} is as defined above, $-NR_{1-a}C=O NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined above, $-C=O NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined above, and $-S(=O)_2 NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

(II) $-(CH_2)_{0-3}-(C_3-C_8)$ cycloalkyl where cycloalkyl can be optionally
30 substituted with one, two or three substituents selected from the group consisting of C_1-C_3 alkyl, $-F$, $-Cl$, $-Br$, $-I$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, C_1-C_6 alkoxy, $-O$ -phenyl, $-$

CO-OH, -CO-O-(C₁-C₄ alkyl), and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(III) -(CR_{A-x}R_{A-y})₀₋₄-R_{A-aryl} where R_{A-x} and R_{A-y} are

(A) -H,

5 (B) C₁-C₄ alkyl optionally substituted with one or two -OH,

(C) C₁-C₄ alkoxy optionally substituted with one, two, or

three of -F,

(D) -(CH₂)₀₋₄-C₃-C₇ cycloalkyl,

(E) C₂-C₆ alkenyl containing one or two double bonds,

10 (F) C₂-C₆ alkynyl containing one or two triple bonds, or

(G) phenyl,

and where R_{A-x} and R_{A-y} are taken together with the carbon to which they are attached to form a carbocycle of three, four, five, six or seven carbon atoms, optionally where one carbon atom is replaced by a heteroatom selected from the

15 group consisting of -O-, -S-, -SO₂-, and -NR_{N-2}- and R_{A-aryl} is the same as R_{N-aryl},

(IV) -(CR_{A-x}R_{A-y})₀₋₄-R_{A-heteroaryl} where R_{A-heteroaryl} is the same as R_{N-}

heteroaryl and R_{A-x} and R_{A-y} are as defined above,

(V) -(CR_{A-x}R_{A-y})₀₋₄-R_{A-aryl}-R_{A-aryl} where R_{A-aryl}, R_{A-x} and R_{A-y} are as defined above,

20 (VI) -(CR_{A-x}R_{A-y})₀₋₄-R_{A-aryl}-R_{A-heteroaryl} where R_{A-aryl}, R_{A-heteroaryl}, R_{A-x} and R_{A-y} are as defined above,

(VII) -(CR_{A-x}R_{A-y})₀₋₄-R_{A-heteroaryl}-R_{A-aryl} where R_{A-heteroaryl}, R_{A-aryl}, R_{A-x} and R_{A-y} are as defined above,

(VIII) -(CR_{A-x}R_{A-y})₀₋₄-R_{A-heteroaryl}-R_{A-heteroaryl} where R_{A-heteroaryl}, R_{A-x}

25 and R_{A-y} are as defined above,

(IX) -(CR_{A-x}R_{A-y})₀₋₄-R_{A-aryl}-R_{A-heterocycle} where R_{A-heterocycle} is defined as R_{1-heterocycle}, and where R_{A-aryl}, R_{A-x} and R_{A-y} are as defined above,

(X) -(CR_{A-x}R_{A-y})₀₋₄-R_{A-heteroaryl}-R_{A-heterocycle} where R_{A-heteroaryl}, R_{A-heterocycle}, R_{A-x} and R_{A-y} are as defined above,

30 (XI) -(CR_{A-x}R_{A-y})₀₋₄-R_{A-heterocycle}-R_{A-aryl} where R_{A-heterocycle}, R_{A-aryl}, R_{A-x} and R_{A-y} are as defined above,

(XII) -(CR_{A-x}R_{A-y})₀₋₄-R_{A-heterocycle}-R_{A-heteroaryl} where R_{A-heterocycle}, R_{A-heteroaryl}, R_{A-x} and R_{A-y} are as defined above,

(XIII) $-(\text{CR}_{\text{A-x}}\text{R}_{\text{A-y}})_{0-4}-\text{R}_{\text{A-heterocycle}}-\text{R}_{\text{A-heterocycle}}$ where $\text{R}_{\text{A-heterocycle}}$, $\text{R}_{\text{A-x}}$ and $\text{R}_{\text{A-y}}$ are as defined above,

(XIV) $-(\text{CR}_{\text{A-x}}\text{R}_{\text{A-y}})_{0-4}-\text{R}_{\text{A-heterocycle}}$ where $\text{R}_{\text{A-heterocycle}}$, $\text{R}_{\text{A-x}}$ and $\text{R}_{\text{A-y}}$ are as defined above,

5 (XV) $-[\text{C}(\text{R}_{\text{A-1}})(\text{R}_{\text{A-2}})]_{1-3}-\text{CO}-\text{N}-(\text{R}_{\text{A-3}})_2$ where $\text{R}_{\text{A-1}}$ and $\text{R}_{\text{A-2}}$ are the same or different and are selected from the group consisting of:

(A) -H,

(B) $\text{C}_1\text{-C}_6$ alkyl, optionally substituted with one, two or three substituents selected from the group consisting of $\text{C}_1\text{-C}_3$ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C \equiv N, -CF₃, $\text{C}_1\text{-C}_6$ alkoxy, -O-phenyl, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(C) $\text{C}_2\text{-C}_6$ alkenyl with one or two double bonds, optionally substituted with one, two or three substituents selected from the group consisting of $\text{C}_1\text{-C}_3$ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C \equiv N, -CF₃, $\text{C}_1\text{-C}_6$ alkoxy, -O-phenyl, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(D) $\text{C}_2\text{-C}_6$ alkynyl with one or two triple bonds, optionally substituted with one, two or three substituents selected from the group consisting of $\text{C}_1\text{-C}_3$ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C \equiv N, -CF₃, $\text{C}_1\text{-C}_6$ alkoxy, -O-phenyl, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

20 (E) $-(\text{CH}_2)_{1-2}-\text{S}(\text{O})_{0-2}-(\text{C}_1\text{-C}_6 \text{ alkyl})$,

(F) $-(\text{CH}_2)_{0-4}-\text{C}_3\text{-C}_7$ cycloalkyl, optionally substituted with one, two or three substituents selected from the group consisting of $\text{C}_1\text{-C}_3$ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C \equiv N, -CF₃, $\text{C}_1\text{-C}_6$ alkoxy, -O-phenyl, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

25 (G) $-(\text{C}_1\text{-C}_4 \text{ alkyl})-\text{R}_{\text{A'-aryl}}$ where $\text{R}_{\text{A'-aryl}}$ is as defined for R_{1-aryl},

(H) $-(\text{C}_1\text{-C}_4 \text{ alkyl})-\text{R}_{\text{A-heteroaryl}}$ where $\text{R}_{\text{A-heteroaryl}}$ is as defined above,

(I) $-(\text{C}_1\text{-C}_4 \text{ alkyl})-\text{R}_{\text{A-heterocycle}}$ where $\text{R}_{\text{A-heterocycle}}$ is as defined above,

(J) $-\text{R}_{\text{A-heteroaryl}}$ where $\text{R}_{\text{A-heteroaryl}}$ is as defined above,

(K) $-\text{R}_{\text{A-heterocycle}}$ where $\text{R}_{\text{A-heterocycle}}$ is as defined above,

(M) $-(\text{CH}_2)_{1-4}-\text{R}_{\text{A-4}}-(\text{CH}_2)_{0-4}-\text{R}_{\text{A'-aryl}}$ where $\text{R}_{\text{A-4}}$ is -O-, -S- or

—NR_{A-5}— where R_{A-5} is C₁-C₆ alkyl, and where R_{A'-aryl} is defined above,

(N) —(CH₂)₁₋₄—R_{A-4}—(CH₂)₀₋₄—R_{A-heteroaryl} where R_{A-4} and R_{A-heteroaryl} are as defined above, and

(O) —R_{A'-aryl} where R_{A'-aryl} is as defined above,

5 and where R_{A-3} is the same or different and is:

(A) —H,

(B) —C₁-C₆ alkyl optionally substituted with one, two or three substituents selected from the group consisting of C₁-C₃ alkyl, —F, —Cl, —Br, —I, —OH, —SH, —C≡N, —CF₃, C₁-C₆ alkoxy, —O-phenyl, —NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

10 defined above,

(C) C₂-C₆ alkenyl with one or two double bonds, optionally substituted with one, two or three substituents selected from the group consisting of C₁-C₃ alkyl, —F, —Cl, —Br, —I, —OH, —SH, —C≡N, —CF₃, C₁-C₆ alkoxy, —O-phenyl, and —NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

15 (D) C₂-C₆ alkynyl with one or two triple bonds, optionally substituted with one, two or three substituents selected from the group consisting of C₁-C₃ alkyl, —F, —Cl, —Br, —I, —OH, —SH, —C≡N, —CF₃, C₁-C₆ alkoxy, —O-phenyl, and —NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(E) —(CH₂)₀₋₄—C₃-C₇ cycloalkyl, optionally substituted with one, two or three substituents selected from the group consisting of C₁-C₃ alkyl, —F, —Cl, —Br, —I, —OH, —SH, —C≡N, —CF₃, C₁-C₆ alkoxy, —O-phenyl, and —NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

20 (F) —R_{A'-aryl} where R_{A'-aryl} is as defined above,

(G) —R_{A-heteroaryl} where R_{A-heteroaryl} is as defined above,

25 (H) —R_{A-heterocycle} where R_{A-heterocycle} is as defined above,

(I) —(C₁-C₄ alkyl)—R_{A'-aryl} where R_{A'-aryl} is as defined above,

(J) —(C₁-C₄ alkyl)—R_{A-heteroaryl} where R_{A-heteroaryl} is as defined

above,

(K) —(C₁-C₄ alkyl)—R_{A-heterocycle} where R_{A-heterocycle} is as defined

30 above, or

(XVI) —CH(R_{A-aryl})₂ where R_{A-aryl} are the same or different and are as defined above,

(XVII) $-\text{CH}(\text{R}_{\text{A-heteroaryl}})_2$ where $\text{R}_{\text{A-heteroaryl}}$ are the same or different and are as defined above,

(XVIII) $-\text{CH}(\text{R}_{\text{A-aryl}})(\text{R}_{\text{A-heteroaryl}})$ where $\text{R}_{\text{A-aryl}}$ and $\text{R}_{\text{A-heteroaryl}}$ are as defined above,

5 (XIX) -cyclopentyl, -cyclohexyl, or -cycloheptyl ring fused to $\text{R}_{\text{A-aryl}}$, $\text{R}_{\text{A-heteroaryl}}$, $\text{R}_{\text{A-heterocycle}}$ where $\text{R}_{\text{A-aryl}}$ or $\text{R}_{\text{A-heteroaryl}}$ or $\text{R}_{\text{A-heterocycle}}$ are as defined above where one carbon of cyclopentyl, cyclohexyl, or -cycloheptyl is optionally replaced with NH, $\text{NR}_{\text{N-5}}$, O, or $\text{S}(=\text{O})_{0-2}$, and where cyclopentyl, cyclohexyl, or -cycloheptyl can be optionally substituted with one or two $-\text{C}_1-\text{C}_3$ alkyl, $-\text{F}$, $-\text{OH}$, $-\text{SH}$, $-\text{C}\equiv\text{N}$, $-\text{CF}_3$, C_1-C_6 alkoxy, $=\text{O}$, or $-\text{NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

10 (XX) C_2-C_{10} alkenyl containing one or two double bonds optionally substituted with one, two or three substituents selected from the group consisting of C_1-C_3 alkyl, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, $-\text{OH}$, $-\text{SH}$, $-\text{C}\equiv\text{N}$, $-\text{CF}_3$, C_1-C_6 alkoxy, $-\text{O-phenyl}$, and $-\text{NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

15 (XXI) C_2-C_{10} alkynyl containing one or two triple bonds optionally substituted with one, two or three substituents selected from the group consisting of C_1-C_3 alkyl, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, $-\text{OH}$, $-\text{SH}$, $-\text{C}\equiv\text{N}$, $-\text{CF}_3$, C_1-C_6 alkoxy, $-\text{O-phenyl}$, $-\text{NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

20 (XXI) $-(\text{CH}_2)_{0-1}-\text{CHR}_{\text{A-6}}-(\text{CH}_2)_{0-1}-\text{R}_{\text{A-aryl}}$ where $\text{R}_{\text{A-aryl}}$ is as defined above and $\text{R}_{\text{A-6}}$ is $-(\text{CH}_2)_{0-6}-\text{OH}$,

(XXII) $-(\text{CH}_2)_{0-1}-\text{CHR}_{\text{A-6}}-(\text{CH}_2)_{0-1}-\text{R}_{\text{A-heteroaryl}}$ where $\text{R}_{\text{A-heteroaryl}}$ and $\text{R}_{\text{A-6}}$ is as defined above,

(XXIII) $-\text{CH}(-\text{R}_{\text{A-aryl}} \text{ or } \text{R}_{\text{A-heteroaryl}})-\text{CO}-\text{O}(\text{C}_1-\text{C}_4 \text{ alkyl})$ where $\text{R}_{\text{A-aryl}}$ and $\text{R}_{\text{A-heteroaryl}}$ are as defined above,

25 (XXIV) $-\text{CH}(-\text{CH}_2-\text{OH})-\text{CH}(-\text{OH})-\text{micro-NO}_2$,

(XXV) $(\text{C}_1-\text{C}_6 \text{ alkyl})-\text{O}-(\text{C}_1-\text{C}_6 \text{ alkyl})-\text{OH}$,

(XXVII) $-\text{CH}_2-\text{NH}-\text{CH}_2-\text{CH}(-\text{O}-\text{CH}_2-\text{CH}_3)_2$,

(XXVIII) $-\text{H}$,

30 (XXIX) $-(\text{CH}_2)_{0-6}-\text{C}(=\text{NR}_{1-a})(\text{NR}_{1-a}\text{R}_{1-b})$ where R_{1-a} and R_{1-b} are as defined above; or

(XXX)

$-\text{C}=\text{OC}(\text{HR}_6)\text{NHR}_7$, where R_6 and R_7 are as defined below,

$-\text{C}=\text{OR}_7$, where R_7 is as defined below,

-C=OOR₇, where R₇ is as defined below, or

- SOOR₇ where R₇ is as defined below,

wherein R₆ is:

- hydrogen,
- 5 C₁ - C₃ alkyl,
- phenyl,
- thioalkoxyalkyl,
- alkyl substituted aryl,
- cycloalkyl,
- 10 cycloalkylalkyl,
- hydroxyalkyl,
- alkoxyalkyl,
- aryloxyalkyl,
- haloalkyl,
- 15 carboxyalkyl,
- alkoxycarbonylalkyl,
- aminoalkyl,
- (N-protected)aminoalkyl,
- alkylaminoalkyl,
- 20 ((N-protected)(alkyl)amino)alkyl,
- dialkylaminoalkyl,
- guanidinoalkyl,
- lower alkenyl,
- heterocyclic,
- 25 (heterocyclic)alkyl),
- arylthioalkyl,
- arylsulfonylalkyl,
- (heterocyclic)thioalkyl,
- (heterocyclic)sulfonylalkyl,
- 30 (heterocyclic)oxyalkyl,
- arylalkoxyalkyl,
- arylthioalkoxyalkyl,
- arylalkylsulfonylalkyl,
- (heterocyclic))alkoxyalkyl,

- (heterocyclic)thioalkoxyalkyl,
 (heterocyclic)alkylsulfonylalkyl,
 cycloalkyloxyalkyl,
 cycloalkylthioalkyl,
 5 cycloalkylsulfonylalkyl,
 cycloalkylalkoxyalkyl,
 cycloalkylthioalkoxyalkyl,
 cycloalkylalkylsulfonylalkyl,
 aminocarbonyl,
 10 alkylaminocarbonyl,
 dialkylaminocarbonyl,
 aroylalkyl,
 (heterocyclic)carbonylalkyl,
 polyhydroxyalkyl,
 15 aminocarbonylalkyl,
 alkylaminocarbonylalkyl,
 dialkylaminocarbonylalkyl,
 aryloxyalkyl, or
 alkylsulfonylalkyl,
 20 wherein heterocyclic is pyridyl, thiazolyl, isothiazolyl,
 oxazolyl, isoxazolyl, furanyl, thienyl, tetrahydrofuranyl, tetrahydrothienyl and
 tetrahydro[2H]pyranyl and wherein the heterocycle is unsubstituted or substituted
 with one to three substituents independently selected from hydroxy, halo, amino,
 alkylamino, dialkylamino, alkoxy, polyalkoxy, haloalkyl, cycloalkyl,
 25 cycloalkylalkyl, aryl, arylalkyl, COOH, -SO₃H, lower alkenyl or lower alkyl;
 wherein R₇ is:
 C₁ - C₃ alkyl,
 phenyl,
 thioalkoxyalkyl,
 30 (aryl)alkyl,
 cycloalkyl,
 cycloalkylalkyl,
 hydroxyalkyl,
 alkoxyalkyl,

5 aryloxyalkyl,
 haloalkyl,
 carboxyalkyl,
 alkoxycarbonylalkyl,
 aminoalkyl,
 (N-protected)aminocalkyl,
 alkylaminoalkyl,
 ((N-protected)(alkyl)amino)alkyl,
 dialkylaminoalkyl,
10 guanidinoalkyl,
 lower alkenyl,
 heterocyclic,
 (heterocyclic)alkyl),
 arylthioalkyl,
15 arylsulfonyalkyl,
 (heterocyclic)thioalkyl,
 (heterocyclic)sulfonylalkyl,
 (heterocyclic)oxyalkyl,
 arylalkoxyalkyl,
20 arylthioalkoxyalkyl,
 arylalkylsulfonylalkyl,
 (heterocyclic)alkoxyalkyl,
 (heterocyclic)thioalkoxyalkyl,
 (heterocyclic)alkylsulfonylalkyl,
25 cycloalkyloxyalkyl,
 cycloalkylthioalkyl,
 cycloalkylsulfonylalkyl,
 cycloalkylalkoxyalkyl,
 cycloalkylthioalkoxyalkyl,
30 cycloalkylalkylsulfonylalkyl,
 aminocarbonyl,
 alkylaminocarbonyl,
 dialkylaminocarbonyl,
 aroalkyl,

(heterocyclic)carbonylalkyl,
 polyhydroxyalkyl,
 aminocarbonylalkyl,
 alkylaminocarbonylalkyl,
 5 dialkylaminocarbonylalkyl,
 aryloxyalkyl, or
 alkylsulfonylalkyl,
 wherein heterocyclic is pyridyl, thiazolyl, isothiazolyl,
 oxazolyl, isoxazolyl, furanyl, thienyl, tetrahydrofuranyl, tetrahydrothienyl and
 10 tetrahydro[2H]pyranyl and wherein the heterocycle is unsubstituted or substituted
 with one to three substituents independently selected from hydroxy, halo, amino,
 alkylamino, dialkylamino, alkoxy, polyalkoxy, haloalkyl, cycloalkyl,
 cycloalkylalkyl, aryl, arylalkyl, COOH, -SO₃H, lower alkenyl or lower alkyl;

15 where X is -N or -O, with the proviso that when X is O, R_B is absent; and
 when X is N,

R_B, is:

(I)-C₁-C₁₀ alkyl optionally substituted with one, two or three
 substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH,
 20 -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as
 defined above, -OC=O NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above, -S(=O)₀₋₂
 R_{1-a} where R_{1-a} is as defined above, - NR_{1-a}C=O NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as
 defined above, -C=O NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above, and -
 S(=O)₂ NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

25 (II) -(CH₂)₀₋₃-(C₃-C₈) cycloalkyl where cycloalkyl can be optionally
 substituted with one, two or three substituents selected from the group consisting of
 C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, -CO-
 OH, -CO-O-(C₁-C₄ alkyl), and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(III) -(CR_{B-x}R_{B-y})₀₋₄-R_{B-aryl} where R_{B-x} and R_{B-y} are

30 (A) -H,
 (B) C₁-C₄ alkyl optionally substituted with one or two -OH,
 (C) C₁-C₄ alkoxy optionally substituted with one, two, or
 three of -F,

(D) $-(CH_2)_{0-4}-C_3-C_7$ cycloalkyl,

(E) C_2-C_6 alkenyl containing one or two double bonds,

(F) C_2-C_6 alkynyl containing one or two triple bonds, or

(G) phenyl,

5 and where R_{B-x} and R_{B-y} are taken together with the carbon to which they are attached to form a carbocycle of three, four, five, six or seven carbon atoms, optionally where one carbon atom is replaced by a heteroatom selected from the group consisting of $-O-$, $-S-$, $-SO_2-$, and $-NR_{N-2}$ where R_{N-2} is as defined above, and R_{B-aryl} is the same as R_{N-aryl} and is defined above

10 (IV) $-(CR_{B-x}R_{B-y})_{0-4}-R_{B-heteroaryl}$ where $R_{B-heteroaryl}$ is the same as $R_{N-heteroaryl}$, R_{B-x} , and R_{B-y} are as defined above,

(V) $-(CR_{B-x}R_{B-y})_{0-4}-R_{B-aryl}-R_{B-aryl}$ where R_{B-aryl} , R_{B-x} , and R_{B-y} are as defined above,

(VI) $-(CR_{B-x}R_{B-y})_{0-4}-R_{B-aryl}-R_{B-heteroaryl}$ where R_{B-aryl} , $R_{B-heteroaryl}$, R_{B-x} and R_{B-y} are as defined above,

(VII) $-(CR_{B-x}R_{B-y})_{0-4}-R_{B-heteroaryl}-R_{B-aryl}$ where $R_{B-heteroaryl}$, R_{B-aryl} , R_{B-x} and R_{B-y} are as defined above,

(VIII) $-(CR_{B-x}R_{B-y})_{0-4}-R_{B-heteroaryl}-R_{B-heteroaryl}$ where $R_{B-heteroaryl}$, R_{B-x} and R_{B-y} are as defined above,

20 (IX) $-(CR_{B-x}R_{B-y})_{0-4}-R_{B-aryl}-R_{B-heterocycle}$ where $R_{B-heterocycle}$ is defined as $R_{1-heterocycle}$, and where R_{B-aryl} , R_{B-x} and R_{B-y} are as defined above,

(X) $-(CR_{B-x}R_{B-y})_{0-4}-R_{B-heteroaryl}-R_{B-heterocycle}$ where $R_{B-heteroaryl}$, $R_{B-heterocycle}$, R_{B-x} and R_{B-y} are as defined above,

(XI) $-(CR_{B-x}R_{B-y})_{0-4}-R_{B-heterocycle}-R_{B-aryl}$ where $R_{B-heterocycle}$, R_{B-aryl} , R_{B-x} and R_{B-y} are as defined above,

(XII) $-(CR_{B-x}R_{B-y})_{0-4}-R_{B-heterocycle}-R_{B-heteroaryl}$ where $R_{B-heterocycle}$, $R_{B-heteroaryl}$, R_{B-x} and R_{B-y} are as defined above,

(XIII) $-(CR_{B-x}R_{B-y})_{0-4}-R_{B-heterocycle}-R_{B-heterocycle}$ where $R_{B-heterocycle}$, R_{B-x} and R_{B-y} are as defined above,

30 (XIV) $-(CR_{B-x}R_{B-y})_{0-4}-R_{B-heterocycle}$ where $R_{B-heterocycle}$, R_{B-x} and R_{B-y} are as defined above,

(XV) $-[C(R_{B-1})(R_{B-2})]_{1-3}-CO-N-(R_{B-3})_2$ where R_{B-1} and R_{B-2} are the same or different and are selected from the group consisting of:

(A) $-H$,

(B) $-C_1-C_6$ alkyl, optionally substituted with one, two or three substituents selected from the group consisting of C_1-C_3 alkyl, $-F$, $-Cl$, $-Br$, $-I$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, C_1-C_6 alkoxy, $-O$ -phenyl, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

5 (C) C_2-C_6 alkenyl with one or two double bonds, optionally substituted with one, two or three substituents selected from the group consisting of C_1-C_3 alkyl, $-F$, $-Cl$, $-Br$, $-I$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, C_1-C_6 alkoxy, $-O$ -phenyl, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

(D) C_2-C_6 alkynyl with one or two triple bonds, optionally substituted with one, two or three substituents selected from the group consisting of C_1-C_3 alkyl, $-F$, $-Cl$, $-Br$, $-I$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, C_1-C_6 alkoxy, $-O$ -phenyl, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

(E) $-(CH_2)_{1-2}-S(O)_{0-2}-(C_1-C_6 \text{ alkyl})$,

(F) $-(CH_2)_{0-4}-C_3-C_7$ cycloalkyl, optionally substituted with one, two or three substituents selected from the group consisting of C_1-C_3 alkyl, $-F$, $-Cl$, $-Br$, $-I$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, C_1-C_6 alkoxy, $-O$ -phenyl, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

(G) $-(C_1-C_4 \text{ alkyl})-R_{B'-aryl}$ where $R_{B'-aryl}$ is as defined above for R_{1-aryl} ,

(H) $-(C_1-C_4 \text{ alkyl})-R_{B-heteroaryl}$ where $R_{B-heteroaryl}$ is as defined above,

(I) $-(C_1-C_4 \text{ alkyl})-R_{B-heterocycle}$ where $R_{B-heterocycle}$ is as defined above,

(J) $-R_{B-heteroaryl}$ where $R_{B-heteroaryl}$ is as defined above,

(K) $-R_{B-heterocycle}$ where $R_{B-heterocycle}$ is as defined above,

(M) $-(CH_2)_{1-4}-R_{B-4}-(CH_2)_{0-4}-R_{B'-aryl}$ where R_{B-4} is $-O-$, $-S-$ or $-NR_{B-5}-$ where R_{B-5} is C_1-C_6 alkyl, and where $R_{B'-aryl}$ is defined above,

(N) $-(CH_2)_{1-4}-R_{B-4}-(CH_2)_{0-4}-R_{B-heteroaryl}$ where R_{B-4} and $R_{B-heteroaryl}$ are as defined above, and

(O) $-R_{B'-aryl}$ where $R_{B'-aryl}$ is as defined above,

and where R_{B-3} is the same or different and is:

(A) $-H$,

(B) $-C_1-C_6$ alkyl optionally substituted with one, two or three substituents selected from the group consisting of C_1-C_3 alkyl, $-F$, $-Cl$, $-Br$, $-I$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, C_1-C_6 alkoxy, $-O$ -phenyl, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

5 (C) C_2-C_6 alkenyl with one or two double bonds, optionally substituted with one, two or three substituents selected from the group consisting of C_1-C_3 alkyl, $-F$, $-Cl$, $-Br$, $-I$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, C_1-C_6 alkoxy, $-O$ -phenyl, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

(D) C_2-C_6 alkynyl with one or two triple bonds, optionally substituted with one, two or three substituents selected from the group consisting of C_1-C_3 alkyl, $-F$, $-Cl$, $-Br$, $-I$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, C_1-C_6 alkoxy, $-O$ -phenyl, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

(E) $-(CH_2)_{0-4}-C_3-C_7$ cycloalkyl, optionally substituted with one, two or three substituents selected from the group consisting of C_1-C_3 alkyl, $-F$, $-Cl$, $-Br$, $-I$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, C_1-C_6 alkoxy, $-O$ -phenyl, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

(F) $-R_{B'-aryl}$ where $R_{B'-aryl}$ is as defined above,

(G) $-R_{B-heteroaryl}$ where $R_{B-heteroaryl}$ is as defined above,

(H) $-R_{B-heterocycle}$ where $R_{B-heterocycle}$ is as defined above,

20 (I) $-(C_1-C_4 \text{ alkyl})-R_{B'-aryl}$ where $R_{B'-aryl}$ is as defined above,

(J) $-(C_1-C_4 \text{ alkyl})-R_{B-heteroaryl}$ where $R_{B-heteroaryl}$ is as defined

above,

(K) $-(C_1-C_4 \text{ alkyl})-R_{B-heterocycle}$ where $R_{B-heterocycle}$ is as defined above, or

25 (XVI) $-\text{CH}(R_{B-aryl})_2$ where R_{B-aryl} are the same or different and are as defined above,

(XVII) $-\text{CH}(R_{B-heteroaryl})_2$ where $R_{B-heteroaryl}$ are the same or different and are as defined above,

(XVIII) $-\text{CH}(R_{B-aryl})(R_{B-heteroaryl})$ where R_{B-aryl} and $R_{B-heteroaryl}$ are as defined above,

30 (XIX) -cyclopentyl, -cyclohexyl, or -cycloheptyl ring fused to R_{B-aryl} or $R_{B-heteroaryl}$ or $R_{B-heterocycle}$ where R_{B-aryl} or $R_{B-heteroaryl}$ or $R_{B-heterocycle}$ are as defined above where one carbon of cyclopentyl, cyclohexyl, or -cycloheptyl is optionally

replaced with NH, NR_{N-5}, O, or S(=O)₀₋₂, and where cyclopentyl, cyclohexyl, or -cycloheptyl can be optionally substituted with one or two -C₁-C₃ alkyl, -F, -OH, -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, =O, or -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

5 (XX) C₂-C₁₀ alkenyl containing one or two double bonds optionally substituted with one, two or three substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(XXI) C₂-C₁₀ alkynyl containing one or two triple bonds optionally
10 substituted with one, two or three substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(XXI) -(CH₂)₀₋₁-CHR_{C-6}-(CH₂)₀₋₁-R_{B-aryl} where R_{B-aryl} is as defined above and R_{C-6} is -(CH₂)₀₋₆-OH,

15 (XXII) -(CH₂)₀₋₁-CHR_{B-6}-(CH₂)₀₋₁-R_{B-heteroaryl} where R_{B-heteroaryl} and R_{C-6} is as defined above,

(XXIII) -CH(-R_{B-aryl} or R_{B-heteroaryl})-CO-O(C₁-C₄ alkyl) where R_{B-aryl} and R_{B-heteroaryl} are as defined above,

(XXIV) -CH(-CH₂-OH)-CH(-OH)-micro-NO₂,

20 (XXV) (C₁-C₆ alkyl)-O-(C₁-C₆ alkyl)-OH,


(XXVII) -CH₂-NH-CH₂-CH(-O-CH₂-CH₃)₂,

(XXVIII) -H, or

(XXIX) -(CH₂)₀₋₆-C(=NR_{1-a})(NR_{1-a}R_{1-b}) where R_{1-a} and R_{1-b} are as defined above,

25 and pharmaceutically acceptable salts thereof.

Disclosed is the use of a compound of formula (XV)



where R₁, R₂, R₃, R_N, R_A, R_B, and X are as defined above for the compound of formula (XV), and pharmaceutically acceptable salts thereof for the manufacture of
30 a medicament for use in treating a patient who has, or in preventing a patient from

getting, a disease or condition selected from the group consisting of Alzheimer's disease, for helping prevent or delay the onset of Alzheimer's disease, for treating patients with mild cognitive impairment (MCI) and preventing or delaying the onset of Alzheimer's disease in those who would progress from MCI to AD, for treating
5 Down's syndrome, for treating humans who have Hereditary Cerebral Hemorrhage with Amyloidosis of the Dutch-Type, for treating cerebral amyloid angiopathy and preventing its potential consequences, i.e. single and recurrent lobar hemorrhages, for treating other degenerative dementias, including dementias of mixed vascular and degenerative origin, dementia associated with Parkinson's disease, dementia
10 associated with progressive supranuclear palsy, dementia associated with cortical basal degeneration, diffuse Lewy body type of Alzheimer's disease and who is in need of such treatment.

The present invention provides compounds, compositions, kits, and methods for inhibiting beta-secretase-mediated cleavage of amyloid precursor protein (APP).
15 More particularly, the compounds, compositions, and methods of the invention are effective to inhibit the production of A beta peptide and to treat or prevent any human or veterinary disease or condition associated with a pathological form of A beta peptide.

The compounds, compositions, and methods of the invention are useful for
20 treating humans who have Alzheimer's Disease (AD), for helping prevent or delay the onset of AD, for treating patients with mild cognitive impairment (MCI), and preventing or delaying the onset of AD in those patients who would otherwise be expected to progress from MCI to AD, for treating Down's syndrome, for treating Hereditary Cerebral Hemorrhage with Amyloidosis of the Dutch Type, for treating
25 cerebral beta-amyloid angiopathy and preventing its potential consequences such as single and recurrent lobar hemorrhages, for treating other degenerative dementias, including dementias of mixed vascular and degenerative origin, for treating dementia associated with Parkinson's disease, dementia associated with progressive supranuclear palsy, dementia associated with cortical basal degeneration, and diffuse
30 Lewy body type AD.

The compounds employed in the methods of the invention possess beta-secretase inhibitory activity. The inhibitory activities of the compounds employed in the methods of the invention are readily demonstrated, for example, using one or more of the assays described herein or known in the art.

DETAILED DESCRIPTION OF THE INVENTION

The invention includes compounds of formula (XV) that are useful in
5 treating and preventing Alzheimer's disease. The anti-Alzheimer's compounds of
formula (XV) are made by methods well known to those skilled in the art from
starting compounds known to those skilled in the art. The process chemistry is well
known to those skilled in the art. Examples of preparing various compounds of
formula (XV) are included in charts A-C. One skilled in the art will appreciate that
10 these are all well known reactions in organic chemistry. A chemist skilled in the art,
knowing the chemical structure of the biologically active compounds of formula
(XV) of the invention would be able to prepare them by known methods from
known starting materials without any additional information. The explanation below
therefore is not necessary but is deemed helpful to those skilled in the art who desire
15 to make compounds of the invention.

CHART A illustrates a general method of synthesizing compounds of the
invention. The anti-Alzheimer's compounds of formula (XV) are prepared by
starting with the corresponding epoxide (I). The epoxides (I) are well known to
those skilled in the art or can be readily prepared from known compounds by
20 methods well known to those skilled in the art. The compounds of formula (XV) of
the present invention have at least two enantiomeric centers which give four
enantiomers. The first of these enantiomeric centers derives from the epoxide
starting material (I). If a desired enantiomer is preferred, it is preferred to
commercially obtain or produce the desired enantiomer (S or R) rather than produce
25 an enantiomerically impure mixture and then have to separate out the desired
enantiomer. For the epoxide (I), R_1 is:

(I) C_1-C_6 alkyl, optionally substituted with one, two or three
substituents selected from the group consisting of C_1-C_3 alkyl, C_1-C_7 alkyl
(optionally substituted with C_1-C_3 alkyl and C_1-C_3 alkoxy), -F, -Cl, -Br, -I, -OH, -
30 SH, -C≡N, -CF₃, C_1-C_3 alkoxy, -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are -H or C_1-C_6 alkyl,
and -OC=O NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(II) $-CH_2-S(O)_{0-2}-(C_1-C_6 \text{ alkyl})$,

(III) $-CH_2-CH_2-S(O)_{0-2}-(C_1-C_6 \text{ alkyl})$,

- (IV) C₂-C₆ alkenyl with one or two double bonds, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, -C≡N, -CF₃, C₁-C₃ alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are -H or C₁-C₆ alkyl,
- 5 (V) C₂-C₆ alkynyl with one or two triple bonds, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, -C≡N, -CF₃, C₁-C₃ alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are -H or C₁-C₆ alkyl,
- 10 (VI) -(CH₂)_{n1}-(R_{1-aryl}) where n₁ is zero or one and where R_{1-aryl} is phenyl, 1-naphthyl, 2-naphthyl and indanyl, indenyl, dihydronaphthalyl, or tetralinyl optionally substituted with one, two, three or four of the following substituents on the aryl ring:
- (A) C₁-C₆ alkyl optionally substituted with one, two or three substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, and C₁-C₃ alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,
- 15 (B) C₂-C₆ alkenyl with one or two double bonds, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, -C≡N, -CF₃, C₁-C₃ alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are -H or C₁-C₆ alkyl,
- 20 (C) C₂-C₆ alkynyl with one or two triple bonds, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, -C≡N, -CF₃, C₁-C₃ alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are -H or C₁-C₆ alkyl,
- 25 (D) -F, Cl, -Br or -I,
- (F) -C₁-C₆ alkoxy optionally substituted with one, two or three of -F,
- (G) -NR_{N-2}R_{N-3} where R_{N-2} and R_{N-3} are as defined below,
- (H) -OH,
- 30 (I) -C≡N,
- (J) C₃-C₇ cycloalkyl, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, -C≡N, -CF₃, C₁-C₃ alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are -H or C₁-C₆ alkyl,

(K) $-\text{CO}-(\text{C}_1\text{-C}_4 \text{ alkyl})$,

(L) $-\text{SO}_2\text{-NR}_{1\text{-a}}\text{R}_{1\text{-b}}$ where $\text{R}_{1\text{-a}}$ and $\text{R}_{1\text{-b}}$ are as defined above,

(M) $-\text{CO-NR}_{1\text{-a}}\text{R}_{1\text{-b}}$ where $\text{R}_{1\text{-a}}$ and $\text{R}_{1\text{-b}}$ are as defined above,

or

5 (N) $-\text{SO}_2-(\text{C}_1\text{-C}_4 \text{ alkyl})$,

(VII) $-(\text{CH}_2)_{n_1}\text{-(R}_{1\text{-heteroaryl}})$ where n_1 is as defined above and where

$\text{R}_{1\text{-heteroaryl}}$ is selected from the group consisting of:

pyridinyl,

pyrimidinyl,

10 quinolinyl,

benzothienyl,

indolyl,

indolinyl,

pyridazinyl,

15 pyrazinyl,

isoquinolyl,

quinazolinyl,

quinoxalinyl,

phthalazinyl,

20 imidazolyl,

isoxazolyl,

pyrazolyl,

oxazolyl,

thiazolyl,

25 indolizinyl,

indazolyl,

benzothiazolyl,

benzimidazolyl,

benzofuranyl,

30 furanyl,

thienyl,

pyrrolyl,

oxadiazolyl,

thiadiazolyl,

	triazolyl,
	tetrazolyl,
	oxazolopyridinyl,
	imidazopyridinyl,
5	isothiazolyl,
	naphthyridinyl,
	cinnolinyl,
	carbazolyl,
	beta-carbolinyl,
10	isochromanlyl,
	chromanlyl,
	tetrahydroisoquinolinyl,
	isoindolinyl,
	isobenzotetrahydrofuranyl,
15	isobenzotetrahydrothienyl,
	isobenzothienyl,
	benzoxazolyl,
	pyridopyridinyl,
	benzotetrahydrofuranyl,
20	benzotetrahydrothienyl,
	purinyl,
	benzodioxolyl,
	triazinyl,
	phenoxazinyl,
25	phenothiazinyl,
	pteridinyl,
	benzothiazolyl,
	imidazopyridinyl,
	imidazothiazolyl,
30	dihydrobenzisoxazinyl,
	benzisoxazinyl,
	benzoxazinyl,
	dihydrobenzisoctiazinyl,
	benzopyranyl,

	benzothiopyranyl,
	coumarinyl,
	isocoumarinyl,
	chromonyl,
5	chromanonyl,
	pyridinyl-N-oxide,
	tetrahydroquinolinyl
	dihydroquinolinyl
	dihydroquinolinonyl
10	dihydroisoquinolinonyl
	dihydrocoumarinyl
	dihydroisocoumarinyl
	isoindolinonyl
	benzodioxanyl
15	benzoxazolinonyl
	pyrrolyl N-oxide,
	pyrimidinyl N-oxide,
	pyridazinyl N-oxide,
	pyrazinyl N-oxide,
20	quinolinyl N-oxide,
	indolyl N-oxide,
	indolinyl N-oxide,
	isoquinolyl N-oxide,
	quinazolinyl N-oxide,
25	quinoxalinyl N-oxide,
	phthalazinyl N-oxide,
	imidazolyl N-oxide,
	isoxazolyl N-oxide,
	oxazolyl N-oxide,
30	thiazolyl N-oxide,
	indolizinyl N-oxide,
	indazolyl N-oxide,
	benzothiazolyl N-oxide,
	benzimidazolyl N-oxide,

- pyrrolyl N-oxide,
 oxadiazolyl N-oxide,
 thiadiazolyl N-oxide,
 triazolyl N-oxide,
 5 tetrazolyl N-oxide,
 benzothiopyranyl S-oxide, and
 benzothiopyranyl S,S-dioxide,

where the $R_{1\text{-heteroaryl}}$ group is bonded to $-(CH_2)_{n1}-$ by any ring atom of the parent $R_{1\text{-heteroaryl}}$ group substituted by hydrogen such that the new bond to the
 10 $R_{1\text{-heteroaryl}}$ group replaces the hydrogen atom and its bond, where heteroaryl is optionally substituted with one, two, three or four of:

(1) C_1 - C_6 alkyl optionally substituted with one, two or three substituents selected from the group consisting of C_1 - C_3 alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C \equiv N, -CF₃, C_1 - C_3 alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as
 15 defined above,

(2) C_2 - C_6 alkenyl with one or two double bonds, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, -C \equiv N, -CF₃, C_1 - C_3 alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are -H or C_1 - C_6 alkyl,

(3) C_2 - C_6 alkynyl with one or two triple bonds, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, -C \equiv N, -CF₃, C_1 - C_3 alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are -H or C_1 - C_6 alkyl,
 20

(4) -F, Cl, -Br or -I,

(6) - C_1 - C_6 alkoxy optionally substituted with one, two, or three of -F,
 25

(7) -NR_{N-2}R_{N-3} where R_{N-2} and R_{N-3} are as defined below,

(8) -OH,

(9) -C \equiv N,
 30

(10) C_3 - C_7 cycloalkyl, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, -C \equiv N, -CF₃, C_1 - C_3 alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are -H or C_1 - C_6 alkyl,

(11) $-\text{CO}-(\text{C}_1\text{-C}_4 \text{ alkyl})$,

(12) $-\text{SO}_2\text{-NR}_{1\text{-a}}\text{R}_{1\text{-b}}$ where $\text{R}_{1\text{-a}}$ and $\text{R}_{1\text{-b}}$ are as defined

above,

(13) $-\text{CO-NR}_{1\text{-a}}\text{R}_{1\text{-b}}$ where $\text{R}_{1\text{-a}}$ and $\text{R}_{1\text{-b}}$ are as defined

5 above, or

(14) $-\text{SO}_2-(\text{C}_1\text{-C}_4 \text{ alkyl})$, with the proviso that when n_1 is zero $\text{R}_{1\text{-heteroaryl}}$ is not bonded to the carbon chain by nitrogen; or

(VIII) $-(\text{CH}_2)_{n_1}-(\text{R}_{1\text{-heterocycle}})$ where n_1 is as defined above and $\text{R}_{1\text{-heterocycle}}$ is selected from the group consisting of:

- | | |
|----|----------------------------------|
| 10 | morpholinyl, |
| | thiomorpholinyl, |
| | thiomorpholinyl S-oxide, |
| | thiomorpholinyl S,S-dioxide, |
| | piperazinyl, |
| 15 | homopiperazinyl, |
| | pyrrolidinyl, |
| | pyrrolinyl, |
| | tetrahydropyranyl, |
| | piperidinyl, |
| 20 | tetrahydrofuranyl, |
| | tetrahydrothienyl, |
| | homopiperidinyl, |
| | homomorpholinyl, |
| | homothiomorpholinyl, |
| 25 | homothiomorpholinyl S,S-dioxide, |
| | oxazolidinonyl, |
| | dihydropyrazolyl, |
| | dihydropyrrolyl, |
| | dihydropyrazinyl, |
| 30 | dihydropyridinyl, |
| | dihydropyrimidinyl, |
| | dihydrofuryl, |
| | dihydropyranyl, |
| | tetrahydrothienyl S-oxide, |

tetrahydrothienyl S,S-dioxide, and
homothiomorpholinyl S-oxide,

where the $R_{1\text{-heterocycle}}$ group is bonded by any atom of the parent $R_{1\text{-heterocycle}}$ group substituted by hydrogen such that the new bond to the $R_{1\text{-heterocycle}}$ group replaces the hydrogen atom and its bond, where heterocycle is optionally substituted with one, two, three or four:

- (1) C_1 - C_6 alkyl optionally substituted with one, two or three substituents selected from the group consisting of C_1 - C_3 alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C \equiv N, -CF₃, C_1 - C_3 alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,
- (2) C_2 - C_6 alkenyl with one or two double bonds, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, -C \equiv N, -CF₃, C_1 - C_3 alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are -H or C_1 - C_6 alkyl,
- (3) C_2 - C_6 alkynyl with one or two triple bonds, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, -C \equiv N, -CF₃, C_1 - C_3 alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are -H or C_1 - C_6 alkyl,
- (4) -F, Cl, -Br or -I,
- (5) C_1 - C_6 alkoxy,
- (6) - C_1 - C_6 alkoxy optionally substituted with one, two, or three -F,
- (7) -NR_{N-2}R_{N-3} where R_{N-2} and R_{N-3} are as defined below,
- (8) -OH,
- (9) -C \equiv N,
- (10) C_3 - C_7 cycloalkyl, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, -C \equiv N, -CF₃, C_1 - C_3 alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are -H or C_1 - C_6 alkyl,
- (11) -CO-(C_1 - C_4 alkyl),
- (12) -SO₂-NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(13) $-\text{CO}-\text{NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are as defined

above;

(14) $-\text{SO}_2-(\text{C}_1-\text{C}_4 \text{ alkyl})$, or

(15) $=\text{O}$, with the proviso that when n_1 is zero R_{1-}

5 heterocycle is not bonded to the carbon chain by nitrogen.

When R_1 is $\text{R}_{1-\text{heteroaryl}}$ or $\text{R}_{1-\text{heterocycle}}$ the bond from the $\text{R}_{1-\text{heteroaryl}}$ or $\text{R}_{1-\text{heterocycle}}$ group to the $-(\text{CH}_2)_{n1}-$ group can be from any ring atom which has an available valence provided that such bond does not result in formation of a charged species or unstable valence. This means that the $\text{R}_{1-\text{heteroaryl}}$ or $\text{R}_{1-\text{heterocycle}}$ group is
 10 bonded to $-(\text{CH}_2)_{n1}-$ by any ring atom of the parent $\text{R}_{1-\text{heteroaryl}}$ or $\text{R}_{1-\text{heterocycle}}$ group which was substituted by hydrogen such that the new bond to the $\text{R}_{1-\text{heteroaryl}}$ or $\text{R}_{1-\text{heterocycle}}$ group replaces the hydrogen atom and its bond.

The epoxide (I) also contains the R_2 and R_3 groups. In the epoxide (I), R_2 and R_3 are each independently:

15 (I)-H,

(II) C_1-C_6 alkyl, optionally substituted with one, two or three substituents selected from the group consisting of C_1-C_3 alkyl, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, $-\text{OH}$, $-\text{SH}$, $-\text{C}\equiv\text{N}$, $-\text{CF}_3$, C_1-C_3 alkoxy, and $-\text{NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

20 (III) $-(\text{CH}_2)_{0-4}-\text{R}_{2-1}$ where R_{2-1} is $\text{R}_{1-\text{aryl}}$ or $\text{R}_{1-\text{heteroaryl}}$ where $\text{R}_{1-\text{aryl}}$ and $\text{R}_{1-\text{heteroaryl}}$ are as defined above

(IV) C_2-C_6 alkenyl with one or two double bonds,

(V) C_2-C_6 alkynyl with one or two triple bonds; or

(VI) $-(\text{CH}_2)_{0-4}-\text{C}_3-\text{C}_7$ cycloalkyl, optionally substituted with one, two
 25 or three substituents selected from the group consisting of $-\text{F}$, $-\text{Cl}$, $-\text{OH}$, $-\text{SH}$, $-\text{C}\equiv\text{N}$, $-\text{CF}_3$, C_1-C_3 alkoxy, and $-\text{NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are $-\text{H}$ or C_1-C_6 alkyl,

and where R_2 and R_3 are taken together with the carbon to which they are attached to form a carbocycle of three, four, five, six, or seven carbon atoms, optionally where one carbon atom is replaced by a heteroatom selected from the
 30 group consisting of $-\text{O}-$, $-\text{S}-$, $-\text{SO}_2-$, and $-\text{NR}_{\text{N}-2}-$, where $\text{R}_{\text{N}-2}$ is as defined below.

It is preferred that R_2 and R_3 both be $-\text{H}$. If R_2 and R_3 are not the same, an additional enantiomeric center is added to the molecule.

Before the synthesis is begun, the free amino group of the epoxide (I) must be protected with an amino protecting group. There are a number of methods well known to those skilled in the art for accomplishing this step. Amino protecting groups are well known to those skilled in the art. See for example, "Protecting Groups in Organic Synthesis", John Wiley and sons, New York, N.Y., 1981, Chapter 7; "Protecting Groups in Organic Chemistry", Plenum Press, New York, N.Y., 1973, Chapter 2. The function of the amino protecting group is to protect the free amino functionality (-NH₂) during subsequent reactions on the epoxide (I) which would not proceed well, either because the amino group would react and be functionalized in a way that is inconsistent with its need to be free for subsequent reactions, or the free amino group would interfere in the reaction. When the amino protecting group is no longer needed, it is removed by methods well known to those skilled in the art. By definition the amino protecting group must be readily removable as is known to those skilled in the art by methods well known to those skilled in the art.

Suitable amino PROTECTING GROUP is selected from the group consisting of *t*-butoxycarbonyl, benzyloxycarbonyl, formyl, trityl, acetyl, trichloroacetyl, dichloroacetyl, chloroacetyl, trifluoroacetyl, difluoroacetyl, fluoroacetyl, 4-phenylbenzyloxycarbonyl, 2-methylbenzyloxycarbonyl, 4-ethoxybenzyloxycarbonyl, 4-fluorobenzyloxycarbonyl, 4-chlorobenzyloxycarbonyl, 3-chlorobenzyloxycarbonyl, 2-chlorobenzyloxycarbonyl, 2,4-dichlorobenzyloxycarbonyl, 4-bromobenzyloxycarbonyl, 3-bromobenzyloxycarbonyl, 4-nitrobenzyloxycarbonyl, 4-cyanobenzyloxycarbonyl, 2-(4-phenyl)isopropoxycarbonyl, 1,1-diphenyleth-1-yloxycarbonyl, 1,1-diphenylprop-1-yloxycarbonyl, 2-phenylprop-2-yloxycarbonyl, 2-(*p*-toluyl)prop-2-yloxycarbonyl, cyclopentanyloxycarbonyl, 1-methylcyclopentanyloxycarbonyl, cyclohexanyloxycarbonyl, 1-methylcyclohexanyloxycarbonyl, 2-methylcyclohexanyloxycarbonyl, 2-(4-toluylsulfonyl)ethoxycarbonyl, 2-(methylsulfonyl)ethoxycarbonyl, 2-(triphenylphosphino)ethoxycarbonyl, fluorenylmethoxycarbonyl, 2-(trimethylsilyl)ethoxycarbonyl, allyloxycarbonyl, 1-(trimethylsilylmethyl)prop-1-enyloxycarbonyl, 5-benzisoxalylmethoxycarbonyl, 4-acetoxybenzyloxycarbonyl, 2,2,2-trichloroethoxycarbonyl, 2-ethynyl-2-propoxycarbonyl, cyclopropylmethoxycarbonyl, 4-(decyloxyl)benzyloxycarbonyl, isobornyloxycarbonyl and 1-piperidyloxycarbonyl, 9-fluorenylmethyl carbonate, -

CH-CH=CH₂ and phenyl-C(=N)-H. It is preferred that the protecting group be *t*-butoxycarbonyl (BOC) and benzyloxycarbonyl (CBZ), it is more preferred that the protecting group be *t*-butoxycarbonyl. One skilled in the art will understand the preferred methods of introducing a *t*-butoxycarbonyl or benzyloxycarbonyl

5 protecting group and may additionally consult T.W. Green and P.G.M. Wuts in "Protective Groups in Organic Chemistry," John Wiley and Sons, 1991 for guidance.

Once the epoxide (I) is protected, the synthesis begins with reaction of a protected epoxide (I) with a hydrazine. The hydrazine provides R_A, and R_B that are present in the final compound (XV). For the hydrazine, R_A is:

10 (I)-C₁-C₁₀ alkyl optionally substituted with one, two or three substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above, -OC(=O)NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above, -S(=O)₀₋₂R_{1-a} where R_{1-a} is as defined above, -NR_{1-a}C(=O)NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above, -C(=O)NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above, and -S(=O)₂NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(II) -(CH₂)₀₋₃-(C₃-C₈) cycloalkyl where cycloalkyl can be optionally substituted with one, two or three substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, -CO-OH, -CO-O-(C₁-C₄ alkyl), and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(III) -(CR_{A-x}R_{A-y})₀₋₄-R_{A-aryl} where R_{A-x} and R_{A-y} are

- (A) -H,
- (B) C₁-C₄ alkyl optionally substituted with one or two -OH,
- 25 (C) C₁-C₄ alkoxy optionally substituted with one, two, or three of -F,
- (D) -(CH₂)₀₋₄-C₃-C₇ cycloalkyl,
- (E) C₂-C₆ alkenyl containing one or two double bonds,
- (F) C₂-C₆ alkynyl containing one or two triple bonds, or
- 30 (G) phenyl,

and where R_{A-x} and R_{A-y} are taken together with the carbon to which they are attached to form a carbocycle of three, four, five, six or seven carbon atoms,

optionally where one carbon atom is replaced by a heteroatom selected from the group consisting of -O-, -S-, -SO₂-, and -NR_{N-2}- and R_{A-aryl} is the same as R_{N-aryl},

(IV) -(CR_{A-x}R_{A-y})₀₋₄-R_{A-heteroaryl} where R_{A-heteroaryl} is the same as R_{N-heteroaryl} and R_{A-x} and R_{A-y} are as defined above,

5 (V) -(CR_{A-x}R_{A-y})₀₋₄-R_{A-aryl}-R_{A-aryl} where R_{A-aryl}, R_{A-x} and R_{A-y} are as defined above,

(VI) -(CR_{A-x}R_{A-y})₀₋₄-R_{A-aryl}-R_{A-heteroaryl} where R_{A-aryl}, R_{A-heteroaryl}, R_{A-x} and R_{A-y} are as defined above,

(VII) -(CR_{A-x}R_{A-y})₀₋₄-R_{A-heteroaryl}-R_{A-aryl} where R_{A-heteroaryl}, R_{A-aryl}, R_{A-x} and R_{A-y} are as defined above,

10 (VIII) -(CR_{A-x}R_{A-y})₀₋₄-R_{A-heteroaryl}-R_{A-heteroaryl} where R_{A-heteroaryl}, R_{A-x} and R_{A-y} are as defined above,

(IX) -(CR_{A-x}R_{A-y})₀₋₄-R_{A-aryl}-R_{A-heterocycle} where R_{A-heterocycle} is defined as R_{1-heterocycle}, and where R_{A-aryl}, R_{A-x} and R_{A-y} are as defined above,

15 (X) -(CR_{A-x}R_{A-y})₀₋₄-R_{A-heteroaryl}-R_{A-heterocycle} where R_{A-heteroaryl}, R_{A-heterocycle}, R_{A-x} and R_{A-y} are as defined above,

(XI) -(CR_{A-x}R_{A-y})₀₋₄-R_{A-heterocycle}-R_{A-aryl} where R_{A-heterocycle}, R_{A-aryl}, R_{A-x} and R_{A-y} are as defined above,

(XII) -(CR_{A-x}R_{A-y})₀₋₄-R_{A-heterocycle}-R_{A-heteroaryl} where R_{A-heterocycle}, R_{A-heteroaryl}, R_{A-x} and R_{A-y} are as defined above,

20 (XIII) -(CR_{A-x}R_{A-y})₀₋₄-R_{A-heterocycle}-R_{A-heterocycle} where R_{A-heterocycle}, R_{A-x} and R_{A-y} are as defined above,

(XIV) -(CR_{A-x}R_{A-y})₀₋₄-R_{A-heterocycle} where R_{A-heterocycle}, R_{A-x} and R_{A-y} are as defined above,

25 (XV) -[C(R_{A-1})(R_{A-2})]₁₋₃-CO-N-(R_{A-3})₂ where R_{A-1} and R_{A-2} are the same or different and are selected from the group consisting of:

(A) -H,

(B) -C₁-C₆ alkyl, optionally substituted with one, two or three substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(C) C₂-C₆ alkenyl with one or two double bonds, optionally substituted with one, two or three substituents selected from the group consisting of

C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(D) C₂-C₆ alkynyl with one or two triple bonds, optionally substituted with one, two or three substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(E) -(CH₂)₁₋₂-S(O)₀₋₂-(C₁-C₆ alkyl),

(F) -(CH₂)₀₋₄-C₃-C₇ cycloalkyl, optionally substituted with one, two or three substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(G) -(C₁-C₄ alkyl)-R_{A'-aryl} where R_{A'-aryl} is as defined for R_{1-aryl},

(H) -(C₁-C₄ alkyl)-R_{A-heteroaryl} where R_{A-heteroaryl} is as defined above,

(I) -(C₁-C₄ alkyl)-R_{A-heterocycle} where R_{A-heterocycle} is as defined above,

(J) -R_{A-heteroaryl} where R_{A-heteroaryl} is as defined above,

(K) -R_{A-heterocycle} where R_{A-heterocycle} is as defined above,

(M) -(CH₂)₁₋₄-R_{A-4}-(CH₂)₀₋₄-R_{A'-aryl} where R_{A-4} is -O-, -S- or -NR_{A-5}- where R_{A-5} is C₁-C₆ alkyl, and where R_{A'-aryl} is defined above,

(N) -(CH₂)₁₋₄-R_{A-4}-(CH₂)₀₋₄-R_{A-heteroaryl} where R_{A-4} and R_{A-heteroaryl} are as defined above, and

(O) -R_{A'-aryl} where R_{A'-aryl} is as defined above,

and where R_{A-3} is the same or different and is:

(A) -H,

(B) -C₁-C₆ alkyl optionally substituted with one, two or three substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(C) C₂-C₆ alkenyl with one or two double bonds, optionally substituted with one, two or three substituents selected from the group consisting of

C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(D) C₂-C₆ alkynyl with one or two triple bonds, optionally substituted with one, two or three substituents selected from the group consisting of
 5 C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(E) -(CH₂)₀₋₄-C₃-C₇ cycloalkyl, optionally substituted with one, two or three substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, and -NR_{1-a}R_{1-b} where
 10 R_{1-a} and R_{1-b} are as defined above,

(F) -R_{A'-aryl} where R_{A'-aryl} is as defined above,

(G) -R_{A-heteroaryl} where R_{A-heteroaryl} is as defined above,

(H) -R_{A-heterocycle} where R_{A-heterocycle} is as defined above,

(I) -(C₁-C₄ alkyl)-R_{A'-aryl} where R_{A'-aryl} is as defined
 15 above,

(J) -(C₁-C₄ alkyl)-R_{A-heteroaryl} where R_{A-heteroaryl} is as defined above,

(K) -(C₁-C₄ alkyl)-R_{A-heterocycle} where R_{A-heterocycle} is as defined above, or

(XVI) -CH(R_{A-aryl})₂ where R_{A-aryl} are the same or different and are as defined above,

(XVII) -CH(R_{A-heteroaryl})₂ where R_{A-heteroaryl} are the same or different and are as defined above,

(XVIII) -CH(R_{A-aryl})(R_{A-heteroaryl}) where R_{A-aryl} and R_{A-heteroaryl} are as defined above,
 25 defined above,

(XIX) -cyclopentyl, -cyclohexyl, or -cycloheptyl ring fused to R_{A-aryl}, R_{A-heteroaryl}, R_{A-heterocycle} where R_{A-aryl} or R_{A-heteroaryl} or R_{A-heterocycle} are as defined above where one carbon of cyclopentyl, cyclohexyl, or -cycloheptyl is optionally replaced with NH, NR_{N-5}, O, or S(=O)₀₋₂, and where cyclopentyl, cyclohexyl, or -cycloheptyl
 30 can be optionally substituted with one or two -C₁-C₃ alkyl, -F, -OH, -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, =O, or -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(XX) C₂-C₁₀ alkenyl containing one or two double bonds optionally substituted with one, two or three substituents selected from the group consisting of

C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(XXI) C₂-C₁₀ alkynyl containing one or two triple bonds optionally substituted with one, two or three substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(XXI) -(CH₂)₀₋₁-CHR_{A-6}-(CH₂)₀₋₁-R_{A-aryl} where R_{A-aryl} is as defined above and R_{A-6} is -(CH₂)₀₋₆-OH,

(XXII) -(CH₂)₀₋₁-CHR_{A-6}-(CH₂)₀₋₁-R_{A-heteroaryl} where R_{A-heteroaryl} and R_{A-6} is as defined above,

(XXIII) -CH(-R_{A-aryl} or R_{A-heteroaryl})-CO-O(C₁-C₄ alkyl) where R_{A-aryl} and R_{A-heteroaryl} are as defined above,

(XXIV) -CH(-CH₂-OH)-CH(-OH)-micro-NO₂,

(XXV) (C₁-C₆ alkyl)-O-(C₁-C₆ alkyl)-OH,

(XXVII) -CH₂-NH-CH₂-CH(-O-CH₂-CH₃)₂,

(XXVIII) -H,

(XXIX) -(CH₂)₀₋₆-C(=NR_{1-a})(NR_{1-a}R_{1-b}) where R_{1-a} and R_{1-b} are as defined above; or

(XXX)

-C=OC(HR₆)NHR₇, where R₆ and R₇ are as defined below,

-C=OR₇, where R₇ is as defined below,

-C=OOR₇, where R₇ is as defined below, or

-SOOR₇ where R₇ is as defined below,

wherein R₆ is:

hydrogen,

C₁ - C₃ alkyl,

phenyl,

thioalkoxyalkyl,

alkyl substituted aryl,

cycloalkyl,

cycloalkylalkyl,

hydroxyalkyl,

alkoxyalkyl,

aryloxyalkyl,
haloalkyl,
carboxyalkyl,
alkoxycarbonylalkyl,
5 aminoalkyl,
(N-protected)aminoalkyl,
alkylaminoalkyl,
((N-protected)(alkyl)amino)alkyl,
dialkylaminoalkyl,
10 guanidinoalkyl,
lower alkenyl,
heterocyclic,
(heterocyclic)alkyl),
arylthioalkyl,
15 arylsulfonylalkyl,
(heterocyclic)thioalkyl,
(heterocyclic)sulfonylalkyl,
(heterocyclic)oxyalkyl,
arylalkoxyalkyl,
20 arylthioalkoxyalkyl,
arylalkylsulfonylalkyl,
(heterocyclic))alkoxyalkyl,
(heterocyclic)thioalkoxyalkyl,
(heterocyclic)alkylsulfonylalkyl,
25 cycloalkyloxyalkyl,
cycloalkylthioalkyl,
cycloalkylsulfonylalkyl,
cycloalkylalkoxyalkyl,
cycloalkylthioalkoxyalkyl,
30 cycloalkylalkylsulfonylalkyl,
aminocarbonyl,
alkylaminocarbonyl,
dialkylaminocarbonyl,
aroalkyl,

(heterocyclic)carbonylalkyl,
polyhydroxyalkyl,
aminocarbonylalkyl,
alkylaminocarbonylalkyl,
5 dialkylaminocarbonylalkyl,
aryloxyalkyl, or
alkylsulfonylalkyl,

wherein heterocyclic is pyridyl, thiazolyl, isothiazolyl,
oxazolyl, isoxazolyl, furanyl, thienyl, tetrahydrofuranyl, tetrahydrothienyl and
10 tetrahydro[2H]pyranyl and wherein the heterocycle is unsubstituted or substituted
with one to three substituents independently selected from hydroxy, halo, amino,
alkylamino, dialkylamino, alkoxy, polyalkoxy, haloalkyl, cycloalkyl,
cycloalkylalkyl, aryl, arylalkyl, COOH, -SO₃H, lower alkenyl or lower alkyl;

wherein R₇ is:

15 C₁ - C₃ alkyl,
phenyl,
thioalkoxyalkyl,
(aryl)alkyl,
cycloalkyl,
20 cycloalkylalkyl,
hydroxyalkyl,
alkoxyalkyl,
aryloxyalkyl,
haloalkyl,
25 carboxyalkyl,
alkoxycarbonylalkyl,
aminoalkyl,
(N-protected)aminocalkyl,
alkylaminoalkyl,
30 ((N-protected)(alkyl)amino)alkyl,
dialkylaminoalkyl,
guanidinoalkyl,
lower alkenyl,
heterocyclic,

- (heterocyclic)alkyl),
arylthioalkyl,
arylsulfonylalkyl,
(heterocyclic)thioalkyl,
5 (heterocyclic)sulfonylalkyl,
(heterocyclic)oxyalkyl,
arylalkoxyalkyl,
arylthioalkoxyalkyl,
arylalkylsulfonylalkyl,
10 (heterocyclic))alkoxyalkyl,
(heterocyclic)thioalkoxyalkyl,
(heterocyclic)alkylsulfonylalkyl,
cycloalkyloxyalkyl,
cycloalkylthioalkyl,
15 cycloalkylsulfonylalkyl,
cycloalkylalkoxyalkyl,
cycloalkylthioalkoxyalkyl,
cycloalkylalkylsulfonylalkyl,
aminocarbonyl,
20 alkylaminocarbonyl,
dialkylaminocarbonyl,
aroylalkyl,
(heterocyclic)carbonylalkyl,
polyhydroxyalkyl,
25 aminocarbonylalkyl,
alkylaminocarbonylalkyl,
dialkylaminocarbonylalkyl,
aryloxyalkyl, or
alkylsulfonylalkyl,
30 wherein heterocyclic is pyridyl, thiazolyl, isothiazolyl,
oxazolyl, isoxazolyl, furanyl, thienyl, tetrahydrofuranyl, tetrahydrothienyl and
tetrahydro[2H]pyranyl and wherein the heterocycle is unsubstituted or substituted
with one to three substituents independently selected from hydroxy, halo, amino,

alkylamino, dialkylamino, alkoxy, polyalkoxy, haloalkyl, cycloalkyl, cycloalkylalkyl, aryl, arylalkyl, COOH, -SO₃H, lower alkenyl or lower alkyl.

The hydrazine also provides R_B in the final compound (XV). For the hydrazine, R_B is:

- 5 (I)-C₁-C₁₀ alkyl optionally substituted with one, two or three substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above, -OC=O NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above, -S(=O)₀₋₂ R_{1-a} where R_{1-a} is as defined above, -NR_{1-a}C=O NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above, -C=O NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above, and -S(=O)₂ NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

- (II) -(CH₂)₀₋₃-(C₃-C₈) cycloalkyl where cycloalkyl can be optionally substituted with one, two or three substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, -CO-OH, -CO-O-(C₁-C₄ alkyl), and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(III) -(CR_{B-x}R_{B-y})₀₋₄-R_{B-aryl} where R_{B-x} and R_{B-y} are

- (A) -H,
 (B) C₁-C₄ alkyl optionally substituted with one or two -OH,
 (C) C₁-C₄ alkoxy optionally substituted with one, two, or
 20 three of -F,
 (D) -(CH₂)₀₋₄-C₃-C₇ cycloalkyl,
 (E) C₂-C₆ alkenyl containing one or two double bonds,
 (F) C₂-C₆ alkynyl containing one or two triple bonds, or
 (G) phenyl,

- 25 and where R_{B-x} and R_{B-y} are taken together with the carbon to which they are attached to form a carbocycle of three, four, five, six or seven carbon atoms, optionally where one carbon atom is replaced by a heteroatom selected from the group consisting of -O-, -S-, -SO₂-, and -NR_{N-2} where R_{N-2} is as defined above, and R_{B-aryl} is the same as R_{N-aryl} and is defined above

- 30 (IV) -(CR_{B-x}R_{B-y})₀₋₄-R_{B-heteroaryl} where R_{B-heteroaryl} is the same as R_{N-heteroaryl}, R_{B-x}, and R_{B-y} are as defined above,

(V) -(CR_{B-x}R_{B-y})₀₋₄-R_{B-aryl}-R_{B-aryl} where R_{B-aryl}, R_{B-x}, and R_{B-y} are as defined above,

(VI) $-(\text{CR}_{\text{B-x}}\text{R}_{\text{B-y}})_{0-4}-\text{R}_{\text{B-aryl}}-\text{R}_{\text{B-heteroaryl}}$ where $\text{R}_{\text{B-aryl}}$, $\text{R}_{\text{B-heteroaryl}}$, $\text{R}_{\text{B-x}}$ and $\text{R}_{\text{B-y}}$ are as defined above,

(VII) $-(\text{CR}_{\text{B-x}}\text{R}_{\text{B-y}})_{0-4}-\text{R}_{\text{B-heteroaryl}}-\text{R}_{\text{B-aryl}}$ where $\text{R}_{\text{B-heteroaryl}}$, $\text{R}_{\text{B-aryl}}$, $\text{R}_{\text{B-x}}$ and $\text{R}_{\text{B-y}}$ are as defined above,

5 (VIII) $-(\text{CR}_{\text{B-x}}\text{R}_{\text{B-y}})_{0-4}-\text{R}_{\text{B-heteroaryl}}-\text{R}_{\text{B-heteroaryl}}$ where $\text{R}_{\text{B-heteroaryl}}$, $\text{R}_{\text{B-x}}$ and $\text{R}_{\text{B-y}}$ are as defined above,

(IX) $-(\text{CR}_{\text{B-x}}\text{R}_{\text{B-y}})_{0-4}-\text{R}_{\text{B-aryl}}-\text{R}_{\text{B-heterocycle}}$ where $\text{R}_{\text{B-heterocycle}}$ is defined as $\text{R}_{1\text{-heterocycle}}$, and where $\text{R}_{\text{B-aryl}}$, $\text{R}_{\text{B-x}}$ and $\text{R}_{\text{B-y}}$ are as defined above,

(X) $-(\text{CR}_{\text{B-x}}\text{R}_{\text{B-y}})_{0-4}-\text{R}_{\text{B-heteroaryl}}-\text{R}_{\text{B-heterocycle}}$ where $\text{R}_{\text{B-heteroaryl}}$, $\text{R}_{\text{B-heterocycle}}$, $\text{R}_{\text{B-x}}$ and $\text{R}_{\text{B-y}}$ are as defined above,

(XI) $-(\text{CR}_{\text{B-x}}\text{R}_{\text{B-y}})_{0-4}-\text{R}_{\text{B-heterocycle}}-\text{R}_{\text{B-aryl}}$ where $\text{R}_{\text{B-heterocycle}}$, $\text{R}_{\text{B-aryl}}$, $\text{R}_{\text{B-x}}$ and $\text{R}_{\text{B-y}}$ are as defined above,

(XII) $-(\text{CR}_{\text{B-x}}\text{R}_{\text{B-y}})_{0-4}-\text{R}_{\text{B-heterocycle}}-\text{R}_{\text{B-heteroaryl}}$ where $\text{R}_{\text{B-heterocycle}}$, $\text{R}_{\text{B-heteroaryl}}$, $\text{R}_{\text{B-x}}$ and $\text{R}_{\text{B-y}}$ are as defined above,

15 (XIII) $-(\text{CR}_{\text{B-x}}\text{R}_{\text{B-y}})_{0-4}-\text{R}_{\text{B-heterocycle}}-\text{R}_{\text{B-heterocycle}}$ where $\text{R}_{\text{B-heterocycle}}$, $\text{R}_{\text{B-x}}$ and $\text{R}_{\text{B-y}}$ are as defined above,

(XIV) $-(\text{CR}_{\text{B-x}}\text{R}_{\text{B-y}})_{0-4}-\text{R}_{\text{B-heterocycle}}$ where $\text{R}_{\text{B-heterocycle}}$, $\text{R}_{\text{B-x}}$ and $\text{R}_{\text{B-y}}$ are as defined above,

(XV) $-[\text{C}(\text{R}_{\text{B-1}})(\text{R}_{\text{B-2}})]_{1-3}-\text{CO}-\text{N}-(\text{R}_{\text{B-3}})_2$ where $\text{R}_{\text{B-1}}$ and $\text{R}_{\text{B-2}}$ are the same or different and are selected from the group consisting of:

(A) -H,

(B) $\text{C}_1\text{-C}_6$ alkyl, optionally substituted with one, two or three substituents selected from the group consisting of $\text{C}_1\text{-C}_3$ alkyl, -F, -Cl, -Br, -I, -OH, -SH, $\text{-C}\equiv\text{N}$, -CF_3 , $\text{C}_1\text{-C}_6$ alkoxy, -O-phenyl, and $\text{-NR}_{1\text{-a}}\text{R}_{1\text{-b}}$ where $\text{R}_{1\text{-a}}$ and $\text{R}_{1\text{-b}}$ are as defined above,

(C) $\text{C}_2\text{-C}_6$ alkenyl with one or two double bonds, optionally substituted with one, two or three substituents selected from the group consisting of $\text{C}_1\text{-C}_3$ alkyl, -F, -Cl, -Br, -I, -OH, -SH, $\text{-C}\equiv\text{N}$, -CF_3 , $\text{C}_1\text{-C}_6$ alkoxy, -O-phenyl, and $\text{-NR}_{1\text{-a}}\text{R}_{1\text{-b}}$ where $\text{R}_{1\text{-a}}$ and $\text{R}_{1\text{-b}}$ are as defined above,

30 (D) $\text{C}_2\text{-C}_6$ alkynyl with one or two triple bonds, optionally substituted with one, two or three substituents selected from the group consisting of $\text{C}_1\text{-C}_3$ alkyl, -F, -Cl, -Br, -I, -OH, -SH, $\text{-C}\equiv\text{N}$, -CF_3 , $\text{C}_1\text{-C}_6$ alkoxy, -O-phenyl, and $\text{-NR}_{1\text{-a}}\text{R}_{1\text{-b}}$ where $\text{R}_{1\text{-a}}$ and $\text{R}_{1\text{-b}}$ are as defined above,

(E) $-(\text{CH}_2)_{1-2}-\text{S}(\text{O})_{0-2}-(\text{C}_1-\text{C}_6 \text{ alkyl})$,

(F) $-(\text{CH}_2)_{0-4}-\text{C}_3-\text{C}_7$ cycloalkyl, optionally substituted with one, two or three substituents selected from the group consisting of C_1-C_3 alkyl, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, $-\text{OH}$, $-\text{SH}$, $-\text{C}\equiv\text{N}$, $-\text{CF}_3$, C_1-C_6 alkoxy, $-\text{O}$ -phenyl, and $-\text{NR}_{1-a}\text{R}_{1-b}$ where

5 R_{1-a} and R_{1-b} are as defined above,

(G) $-(\text{C}_1-\text{C}_4 \text{ alkyl})-\text{R}_{\text{B}'\text{-aryl}}$ where $\text{R}_{\text{B}'\text{-aryl}}$ is as defined above for $\text{R}_{1\text{-aryl}}$,

(H) $-(\text{C}_1-\text{C}_4 \text{ alkyl})-\text{R}_{\text{B-heteroaryl}}$ where $\text{R}_{\text{B-heteroaryl}}$ is as defined above,

10 (I) $-(\text{C}_1-\text{C}_4 \text{ alkyl})-\text{R}_{\text{B-heterocycle}}$ where $\text{R}_{\text{B-heterocycle}}$ is as defined above,

(J) $-\text{R}_{\text{B-heteroaryl}}$ where $\text{R}_{\text{B-heteroaryl}}$ is as defined above,

(K) $-\text{R}_{\text{B-heterocycle}}$ where $\text{R}_{\text{B-heterocycle}}$ is as defined above,

(M) $-(\text{CH}_2)_{1-4}-\text{R}_{\text{B-4}}-(\text{CH}_2)_{0-4}-\text{R}_{\text{B}'\text{-aryl}}$ where $\text{R}_{\text{B-4}}$ is $-\text{O}-$, $-\text{S}-$ or $-\text{NR}_{\text{B-5}}-$ where $\text{R}_{\text{B-5}}$ is C_1-C_6 alkyl, and where $\text{R}_{\text{B}'\text{-aryl}}$ is defined above,

(N) $-(\text{CH}_2)_{1-4}-\text{R}_{\text{B-4}}-(\text{CH}_2)_{0-4}-\text{R}_{\text{B-heteroaryl}}$ where $\text{R}_{\text{B-4}}$ and $\text{R}_{\text{B-heteroaryl}}$ are as defined above, and

(O) $-\text{R}_{\text{B}'\text{-aryl}}$ where $\text{R}_{\text{B}'\text{-aryl}}$ is as defined above,

and where $\text{R}_{\text{B-3}}$ is the same or different and is:

20 (A) $-\text{H}$,

(B) C_1-C_6 alkyl optionally substituted with one, two or three substituents selected from the group consisting of C_1-C_3 alkyl, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, $-\text{OH}$, $-\text{SH}$, $-\text{C}\equiv\text{N}$, $-\text{CF}_3$, C_1-C_6 alkoxy, $-\text{O}$ -phenyl, and $-\text{NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

25 (C) C_2-C_6 alkenyl with one or two double bonds, optionally substituted with one, two or three substituents selected from the group consisting of C_1-C_3 alkyl, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, $-\text{OH}$, $-\text{SH}$, $-\text{C}\equiv\text{N}$, $-\text{CF}_3$, C_1-C_6 alkoxy, $-\text{O}$ -phenyl, and $-\text{NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

(D) C_2-C_6 alkynyl with one or two triple bonds, optionally
30 substituted with one, two or three substituents selected from the group consisting of C_1-C_3 alkyl, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, $-\text{OH}$, $-\text{SH}$, $-\text{C}\equiv\text{N}$, $-\text{CF}_3$, C_1-C_6 alkoxy, $-\text{O}$ -phenyl, and $-\text{NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

(E) $-(CH_2)_{0-4}-C_3-C_7$ cycloalkyl, optionally substituted with one, two or three substituents selected from the group consisting of C_1-C_3 alkyl, $-F$, $-Cl$, $-Br$, $-I$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, C_1-C_6 alkoxy, $-O$ -phenyl, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

5 (F) $-R_{B'-aryl}$ where $R_{B'-aryl}$ is as defined above,

(G) $-R_{B-heteroaryl}$ where $R_{B-heteroaryl}$ is as defined above,

(H) $-R_{B-heterocycle}$ where $R_{B-heterocycle}$ is as defined above,

(I) $-(C_1-C_4 \text{ alkyl})-R_{B'-aryl}$ where $R_{B'-aryl}$ is as defined above,

10 (J) $-(C_1-C_4 \text{ alkyl})-R_{B-heteroaryl}$ where $R_{B-heteroaryl}$ is as defined above,

(K) $-(C_1-C_4 \text{ alkyl})-R_{B-heterocycle}$ where $R_{B-heterocycle}$ is as defined above, or

(XVI) $-\text{CH}(R_{B-aryl})_2$ where R_{B-aryl} are the same or different and are as defined above,

(XVII) $-\text{CH}(R_{B-heteroaryl})_2$ where $R_{B-heteroaryl}$ are the same or different and are as defined above,

(XVIII) $-\text{CH}(R_{B-aryl})(R_{B-heteroaryl})$ where R_{B-aryl} and $R_{B-heteroaryl}$ are as defined above,

20 (XIX) -cyclopentyl, -cyclohexyl, or -cycloheptyl ring fused to R_{B-aryl} or $R_{B-heteroaryl}$ or $R_{B-heterocycle}$ where R_{B-aryl} or $R_{B-heteroaryl}$ or $R_{B-heterocycle}$ are as defined above where one carbon of cyclopentyl, cyclohexyl, or -cycloheptyl is optionally replaced with NH , NR_{N-5} , O , or $S(=O)_{0-2}$, and where cyclopentyl, cyclohexyl, or -cycloheptyl can be optionally substituted with one or two $-C_1-C_3$ alkyl, $-F$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, C_1-C_6 alkoxy, $=O$, or $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

(XX) C_2-C_{10} alkenyl containing one or two double bonds optionally substituted with one, two or three substituents selected from the group consisting of C_1-C_3 alkyl, $-F$, $-Cl$, $-Br$, $-I$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, C_1-C_6 alkoxy, $-O$ -phenyl, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

30 (XXI) C_2-C_{10} alkynyl containing one or two triple bonds optionally substituted with one, two or three substituents selected from the group consisting of

C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(XXI) -(CH₂)₀₋₁-CHR_{C-6}-(CH₂)₀₋₁-RB_{B-aryl} where R_{B-aryl} is as defined above and R_{C-6} is -(CH₂)₀₋₆-OH,

5 (XXII) -(CH₂)₀₋₁-CHR_{B-6}-(CH₂)₀₋₁-R_{B-heteroaryl} where R_{B-heteroaryl} and R_{C-6} is as defined above,

(XXIII) -CH(-R_{B-aryl} or R_{B-heteroaryl})-CO-O(C₁-C₄ alkyl) where R_{B-aryl} and R_{B-heteroaryl} are as defined above,

(XXIV) -CH(-CH₂-OH)-CH(-OH)-micro-NO₂,

10 (XXV) (C₁-C₆ alkyl)-O-(C₁-C₆ alkyl)-OH,

(XXVII) -CH₂-NH-CH₂-CH(-O-CH₂-CH₃)₂,

(XXVIII) -H, or

(XXIX) -(CH₂)₀₋₆-C(=NR_{1-a})(NR_{1-a}R_{1-b}) where R_{1-a} and R_{1-b} are as defined above.

15 It is preferred that R_A and R_B are, independently, C₁-C₈ alkyl, (CH₂)₀₋₃-(C₃-C₇) cycloalkyl, (CR_{A-x}R_{A-y})₀₋₄-R_{A-aryl}, (CR_{A-x}R_{A-y})₀₋₄-R_{A-heteroaryl}, (CR_{A-x}R_{A-y})₀₋₄-R_{A-heterocycle}, cyclopentyl or cyclohexyl ring fused to R_{A-aryl} or R_{A-heteroaryl} or R_{A-heterocycle}. It is more preferred that -(CH₂)₀₋₃-(C₃-C₇) cycloalkyl, (CR_{A-x}R_{A-y})₀₋₄-R_{A-aryl}, (CR_{A-x}R_{A-y})₀₋₄-R_{A-heteroaryl}, (CR_{A-x}R_{A-y})₀₋₄-R_{A-heterocycle}, or cyclopentyl or -cyclohexyl ring
20 fused to a R_{A-aryl} or R_{A-heteroaryl} or R_{A-heterocycle}. It is most preferred that R_B is (CR_{C-x}R_{C-y})₀₋₄-R_{C-aryl}, (CR_{C-x}R_{C-y})₀₋₄-R_{C-heteroaryl}, cyclopentyl or -cyclohexyl ring fused to a R_{C-aryl} or R_{C-heteroaryl} or R_{C-heterocycle}.

The epoxide (I) is combined with the hydrazine in hot isopropanol resulting in the selective formation of the hydrazine (II) arising from alkylation of the
25 unsubstituted nitrogen (M. Nakakata, *Tetrahedron Letters* **1993**, 6095-6098). Monoacylation of the hydrazine -NH-NH- with benzyloxycarbonyl chloride or other acylating agent gives (III) and reduces the reactivity of this group to further acylation irrespective of which hydrazine nitrogen the first acyl group becomes attached to (B. Gisin, *Helv. Chim. Acta* **1970**, vol 53, 1030-1043. S. Shinagawa,
30 *Chem. Pharm. Bull.* **1981**, vol 29, 3630-3638). Removal of the *tert*-butoxycarbonyl protecting group of (III) will provide the free amine (IV), which is coupled to the compound that provides R_N. R_N is:

(I) R_{N-1}-X_N- where X_N is selected from the group consisting of:

(A) $-\text{CO}-$,

(B) $-\text{SO}_2-$,

(C) $-(\text{CR}'\text{R}'')_{1-6}$ where R' and R'' are the same or different and are $-\text{H}$ and $\text{C}_1\text{-C}_4$ alkyl,

5 (D) $-\text{CO}-(\text{CR}'\text{R}'')_{1-6}-\text{X}_{\text{N}-1}$ where $\text{X}_{\text{N}-1}$ is selected from the group consisting of $-\text{O}-$, $-\text{S}-$ and $-\text{NR}'-$ and where R' and R'' are as defined above, and

(E) a single bond;

where $\text{R}_{\text{N}-1}$ is selected from the group consisting of:

10 (A) $\text{R}_{\text{N-aryl}}$ where $\text{R}_{\text{N-aryl}}$ is phenyl, 1-naphthyl, 2-naphthyl, tetralinyl, indanyl, 6,7,8,9-tetrahydro-5H-benzo[a]cycloheptenyl, or dihydronaphthyl optionally substituted with one, two or three of the following substituents which can be the same or different and are:

(1) $\text{C}_1\text{-C}_6$ alkyl, optionally substituted with one, two or
15 three substituents selected from the group consisting of $\text{C}_1\text{-C}_3$ alkyl, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, $-\text{OH}$, $-\text{SH}$, $-\text{C}\equiv\text{N}$, $-\text{CF}_3$, $\text{C}_1\text{-C}_3$ alkoxy, and $-\text{NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

(2) $-\text{OH}$,

(3) $-\text{NO}_2$,

20 (4) $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$,

(5) $-\text{CO}-\text{OH}$,

(6) $-\text{C}\equiv\text{N}$,

(7) $-(\text{CH}_2)_{0-4}-\text{CO}-\text{NR}_{\text{N}-2}\text{R}_{\text{N}-3}$ where $\text{R}_{\text{N}-2}$ and $\text{R}_{\text{N}-3}$ are the same or different and are selected from the group consisting of:

25 (a) $-\text{H}$,

(b) $-\text{C}_1\text{-C}_6$ alkyl optionally substituted with one substituent selected from the group consisting of:

(i) $-\text{OH}$, and

(ii) $-\text{NH}_2$,

30 (c) $-\text{C}_1\text{-C}_6$ alkyl optionally substituted with one, two, or three $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, or $-\text{I}$,

(d) $-\text{C}_3\text{-C}_7$ cycloalkyl,

(e) $-(\text{C}_1\text{-C}_2 \text{ alkyl})-(\text{C}_3\text{-C}_7 \text{ cycloalkyl})$,

- (f) $-(C_1-C_6 \text{ alkyl})-O-(C_1-C_3 \text{ alkyl})$,
 (g) $-C_2-C_6 \text{ alkenyl}$ with one or two double bonds,
 (h) $-C_2-C_6 \text{ alkynyl}$ with one or two triple bonds,
 (i) $-C_1-C_6 \text{ alkyl chain}$ with one double bond and one triple bond,
 (j) $-R_{1\text{-aryl}}$ where $R_{1\text{-aryl}}$ is as defined above, and
 (k) $-R_{1\text{-heteroaryl}}$ where $R_{1\text{-heteroaryl}}$ is as defined above,
 (8) $-(CH_2)_{0-4}-CO-(C_1-C_{12} \text{ alkyl})$,
 (9) $-(CH_2)_{0-4}-CO-(C_2-C_{12} \text{ alkenyl})$ with one, two or three double bonds),
 (10) $-(CH_2)_{0-4}-CO-(C_2-C_{12} \text{ alkynyl})$ with one, two or three triple bonds),
 (11) $-(CH_2)_{0-4}-CO-(C_3-C_7 \text{ cycloalkyl})$,
 (12) $-(CH_2)_{0-4}-CO-R_{1\text{-aryl}}$ where $R_{1\text{-aryl}}$ is as defined above,
 (13) $-(CH_2)_{0-4}-CO-R_{1\text{-heteroaryl}}$ where $R_{1\text{-heteroaryl}}$ is as defined above,
 (14) $-(CH_2)_{0-4}-CO-R_{1\text{-heterocycle}}$ where $R_{1\text{-heterocycle}}$ is as defined above,
 (15) $-(CH_2)_{0-4}-CO-R_{N-4}$ where R_{N-4} is selected from the group consisting of morpholinyl, thiomorpholinyl, piperazinyl, piperidinyl, homomorpholinyl, homothiomorpholinyl, homothiomorpholinyl S-oxide, homothiomorpholinyl S,S-dioxide, pyrrolinyl and pyrrolidinyl where each group is optionally substituted with one, two, three, or four of $C_1-C_6 \text{ alkyl}$,
 (16) $-(CH_2)_{0-4}-CO-O-R_{N-5}$ where R_{N-5} is selected from the group consisting of:
 (a) $C_1-C_6 \text{ alkyl}$,
 (b) $-(CH_2)_{0-2}-(R_{1\text{-aryl}})$ where $R_{1\text{-aryl}}$ is as defined above,
 (c) $C_2-C_6 \text{ alkenyl}$ containing one or two double bonds,

- (d) C₂-C₆ alkynyl containing one or two triple bonds,
- (e) C₃-C₇ cycloalkyl, and
- (f) -(CH₂)₀₋₂-(R_{1-heteroaryl}) where R_{1-heteroaryl} is as defined above,
- (17) -(CH₂)₀₋₄-SO₂-NR_{N-2}R_{N-3} where R_{N-2} and R_{N-3} are as defined above,
- (18) -(CH₂)₀₋₄-SO-(C₁-C₈ alkyl),
- (19) -(CH₂)₀₋₄-SO₂-(C₁-C₁₂ alkyl),
- (20) -(CH₂)₀₋₄-SO₂-(C₃-C₇ cycloalkyl),
- (21) -(CH₂)₀₋₄-N(H or R_{N-5})-CO-O-R_{N-5} where R_{N-5} can be the same or different and is as defined above,
- (22) -(CH₂)₀₋₄-N(H or R_{N-5})-CO-N(R_{N-5})₂, where R_{N-5} can be the same or different and is as defined above,
- (23) -(CH₂)₀₋₄-N-CS-N(R_{N-5})₂, where R_{N-5} can be the same or different and is as defined above,
- (24) -(CH₂)₀₋₄-N(H or R_{N-5})-CO-R_{N-2} where R_{N-5} and R_{N-2} can be the same or different and are as defined above,
- (25) -(CH₂)₀₋₄-NR_{N-2}R_{N-3} where R_{N-2} and R_{N-3} can be the same or different and are as defined above,
- (26) -(CH₂)₀₋₄-R_{N-4} where R_{N-4} is as defined above,
- (27) -(CH₂)₀₋₄-O-CO-(C₁-C₆ alkyl),
- (28) -(CH₂)₀₋₄-O-P(O)-(OR_{N-aryl-1})₂ where R_{N-aryl-1} is -H or C₁-C₄ alkyl,
- (29) -(CH₂)₀₋₄-O-CO-N(R_{N-5})₂ where R_{N-5} is as defined above,
- (30) -(CH₂)₀₋₄-O-CS-N(R_{N-5})₂ where R_{N-5} is as defined above,
- (31) -(CH₂)₀₋₄-O-(R_{N-5})₂ where R_{N-5} is as defined above,
- (32) -(CH₂)₀₋₄-O-(R_{N-5})₂-COOH where R_{N-5} is as defined above,
- (33) -(CH₂)₀₋₄-S-(R_{N-5})₂ where R_{N-5} is as defined above,

(34) $-(\text{CH}_2)_{0-4}-\text{O}-(\text{C}_1-\text{C}_6 \text{ alkyl optionally substituted with one, two, three, four, or five of }-\text{F})$,

(35) $\text{C}_3-\text{C}_7 \text{ cycloalkyl}$,

(36) $\text{C}_2-\text{C}_6 \text{ alkenyl with one or two double bonds}$

5 optionally substituted with $\text{C}_1-\text{C}_3 \text{ alkyl}$, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, $-\text{OH}$, $-\text{SH}$, $-\text{C}\equiv\text{N}$, $-\text{CF}_3$, $\text{C}_1-\text{C}_3 \text{ alkoxy}$, or $-\text{NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

(37) $\text{C}_2-\text{C}_6 \text{ alkynyl with one or two triple bonds}$

optionally substituted with $\text{C}_1-\text{C}_3 \text{ alkyl}$, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, $-\text{OH}$, $-\text{SH}$, $-\text{C}\equiv\text{N}$, $-\text{CF}_3$, $\text{C}_1-\text{C}_3 \text{ alkoxy}$, or $-\text{NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

10 (38) $-(\text{CH}_2)_{0-4}-\text{N}(-\text{H or } \text{R}_{\text{N}-5})-\text{SO}_2-\text{R}_{\text{N}-2}$ where $\text{R}_{\text{N}-5}$ and $\text{R}_{\text{N}-2}$ can be the same or different and are as described above, or

(39) $-(\text{CH}_2)_{0-4}-\text{C}_3-\text{C}_7 \text{ cycloalkyl}$,

(B) $-\text{R}_{\text{N-heteroaryl}}$ where $\text{R}_{\text{N-heteroaryl}}$ is selected from the group

consisting of:

15 pyridinyl,
pyrimidinyl,
quinolinyl,
benzothienyl,
indolyl,
20 indolinyl,
pyridazinyl,
pyrazinyl,
isoindolyl,
isoquinolyl,
25 quinazolinyl,
quinoxalinyl,
phthalazinyl,
imidazolyl,
isoxazolyl,
30 pyrazolyl,
oxazolyl,
thiazolyl,
indolizinyl,

	indazolyl,
	benzothiazolyl,
	benzimidazolyl,
	benzofuranyl,
5	furanyl,
	thienyl,
	pyrrolyl,
	oxadiazolyl,
	thiadiazolyl,
10	triazolyl,
	tetrazolyl,
	oxazolopyridinyl,
	imidazopyridinyl,
	isothiazolyl,
15	naphthyridinyl,
	cinnoliny,
	carbazolyl,
	beta-carbolinyl,
	isochromanyl,
20	chromanyl,
	tetrahydroisoquinoliny,
	isoindoliny,
	isobenzotetrahydrofuranyl,
	isobenzotetrahydrothienyl,
25	isobenzothieryl,
	benzoxazolyl,
	pyridopyridinyl,
	benzotetrahydrofuranyl,
	benzotetrahydrothienyl,
30	purinyl,
	benzodioxolyl,
	triazinyl,
	phenoxazinyl,
	phenothiazinyl,

	pteridinyI,
	benzothiazolyl,
	imidazopyridinyI,
	imidazothiazolyl,
5	dihydrobenzisoXazinyI,
	benzisoXazinyI,
	benzoxazinyI,
	dihydrobenzisothiazinyI,
	benzopyranyI,
10	benzothiopyranyI,
	coumarinyI,
	isocoumarinyI,
	chromonyI,
	chromanonyI,
15	pyridinyI-N-oxide,
	tetrahydroquinolinyI,
	dihydroquinolinyI,
	dihydroquinolinonyI,
	dihydroisoquinolinonyI,
20	dihydrocoumarinyI,
	dihydroisocoumarinyI,
	isoindolinonyI,
	benzodioxanyI,
	benzoxazolinonyI,
25	pyrrolyI N-oxide,
	pyrimidinyI N-oxide,
	pyridazinyI N-oxide,
	pyrazinyI N-oxide,
	quinolinyI N-oxide,
30	indolyI N-oxide,
	indolinyI N-oxide,
	isoquinolyI N-oxide,
	quinazolinyI N-oxide,
	quinoxalinyI N-oxide,

- phthalazinyl N-oxide,
 imidazolyl N-oxide,
 isoxazolyl N-oxide,
 oxazolyl N-oxide,
 5 thiazolyl N-oxide,
 indoliziny N-oxide,
 indazolyl N-oxide,
 benzothiazolyl N-oxide,
 benzimidazolyl N-oxide,
 10 pyrrolyl N-oxide,
 oxadiazolyl N-oxide,
 thiadiazolyl N-oxide,
 triazolyl N-oxide,
 tetrazolyl N-oxide,
 15 benzothiopyranyl S-oxide, and
 benzothiopyranyl S,S-dioxide

where the $R_{N\text{-heteroaryl}}$ group is bonded by any atom of the parent $R_{N\text{-heteroaryl}}$ group substituted by hydrogen such that the new bond to the $R_{N\text{-heteroaryl}}$ group replaces the hydrogen atom and its bond, where heteroaryl is
 20 optionally substituted with one, two, three, or four of:

- (1) $C_1\text{-}C_6$ alkyl, optionally substituted with one, two or three substituents selected from the group consisting of $C_1\text{-}C_3$ alkyl, -F, -Cl, -Br, -I, -OH, -SH, $-C\equiv N$, $-CF_3$, $C_1\text{-}C_3$ alkoxy, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,
 25 (2) -OH,
 (3) $-NO_2$,
 (4) -F, -Cl, -Br, or -I,
 (5) -CO-OH,
 (6) $-C\equiv N$,
 30 (7) $-(CH_2)_{0-4}\text{-CO-NR}_{N-2}R_{N-3}$ where R_{N-2} and R_{N-3} are the same or different and are selected from the group consisting of:

- (a) -H,

(b) -C₁-C₆ alkyl optionally substituted with one substituent selected from the group consisting of:

(i) -OH, and

(ii) -NH₂,

5 (c) -C₁-C₆ alkyl optionally substituted with one, two, or three -F, -Cl, -Br, -I,

(d) -C₃-C₇ cycloalkyl,

(e) -(C₁-C₂ alkyl)-(C₃-C₇ cycloalkyl),

(f) -(C₁-C₆ alkyl)-O-(C₁-C₃ alkyl),

10 (g) -C₂-C₆ alkenyl with one or two double bonds,

(h) -C₂-C₆ alkynyl with one or two triple

bonds,

(i) -C₁-C₆ alkyl chain with one double bond

15 and one triple bond,

(j) -R_{1-aryl} where R_{1-aryl} is as defined above,

(k) -R_{1-heteroaryl} where R_{1-heteroaryl} is as defined

above,

(8) -(CH₂)₀₋₄-CO-(C₁-C₁₂ alkyl),

20 (9) -(CH₂)₀₋₄-CO-(C₂-C₁₂ alkenyl with one, two or three double bonds),

(10) -(CH₂)₀₋₄-CO-(C₂-C₁₂ alkynyl with one, two or

three triple bonds),

(11) -(CH₂)₀₋₄-CO-(C₃-C₇ cycloalkyl),

25 (12) -(CH₂)₀₋₄-CO-R_{1-aryl} where R_{1-aryl} is as defined above,

(13) -(CH₂)₀₋₄-CO-R_{1-heteroaryl} where R_{1-heteroaryl} is as

defined above,

(14) -(CH₂)₀₋₄-CO-R_{1-heterocycle} where R_{1-heterocycle} is as

30 defined above,

(15) -(CH₂)₀₋₄-CO-R_{N-4} where R_{N-4} is selected from

the group consisting of morpholinyl, thiomorpholinyl, piperazinyl, piperidinyl, homomorpholinyl, homothiomorpholinyl, homothiomorpholinyl S-oxide,

homothiomorpholinyl S,S-dioxide, pyrrolinyl and pyrrolidinyl where each group is optionally substituted with one, two, three, or four of C₁-C₆ alkyl,

(16) $-(CH_2)_{0-4}-CO-O-R_{N-5}$ where R_{N-5} is selected from

the group consisting of:

- 5 (a) C₁-C₆ alkyl,
 (b) $-(CH_2)_{0-2}-(R_{1-aryl})$ where R_{1-aryl} is as defined
 above,
 (c) C₂-C₆ alkenyl containing one or two double
 bonds,
 10 (d) C₂-C₆ alkynyl containing one or two triple
 bonds,
 (e) C₃-C₇ cycloalkyl, and
 (f) $-(CH_2)_{0-2}-(R_{1-heteroaryl})$ where R_{1-heteroaryl} is as
 defined above,

15 (17) $-(CH_2)_{0-4}-SO_2-NR_{N-2}R_{N-3}$ where R_{N-2} and R_{N-3} are
 as defined above,

(18) $-(CH_2)_{0-4}-SO-(C_1-C_8 \text{ alkyl})$,

(19) $-(CH_2)_{0-4}-SO_2-(C_1-C_{12} \text{ alkyl})$,

(20) $-(CH_2)_{0-4}-SO_2-(C_3-C_7 \text{ cycloalkyl})$,

20 (21) $-(CH_2)_{0-4}-N(H \text{ or } R_{N-5})-CO-O-R_{N-5}$ where R_{N-5}
 can be the same or different and is as defined above,

(22) $-(CH_2)_{0-4}-N(H \text{ or } R_{N-5})-CO-N(R_{N-5})_2$, where R_{N-5}
 can be the same or different and is as defined above,

(23) $-(CH_2)_{0-4}-N-CS-N(R_{N-5})_2$, where R_{N-5} can be the
 25 same or different and is as defined above,

(24) $-(CH_2)_{0-4}-N(-H \text{ or } R_{N-5})-CO-R_{N-2}$ where R_{N-5} and
 R_{N-2} can be the same or different and are as defined above,

(25) $-(CH_2)_{0-4}-NR_{N-2}R_{N-3}$ where R_{N-2} and R_{N-3} can be
 the same or different and are as defined above,

30 (26) $-(CH_2)_{0-4}-R_{N-4}$ where R_{N-4} is as defined above,

(27) $-(CH_2)_{0-4}-O-CO-(C_1-C_6 \text{ alkyl})$,

(28) $-(CH_2)_{0-4}-O-P(O)-(OR_{N-aryl-1})_2$ where R_{N-aryl-1} is -

H or C₁-C₄ alkyl,

- (29) $-(\text{CH}_2)_{0-4}-\text{O}-\text{CO}-\text{N}(\text{R}_{\text{N}-5})_2$ where $\text{R}_{\text{N}-5}$ is as defined above,
- (30) $-(\text{CH}_2)_{0-4}-\text{O}-\text{CS}-\text{N}(\text{R}_{\text{N}-5})_2$ where $\text{R}_{\text{N}-5}$ is as defined above,
- 5 (31) $-(\text{CH}_2)_{0-4}-\text{O}-(\text{R}_{\text{N}-5})_2$ where $\text{R}_{\text{N}-5}$ is as defined above,
- (32) $-(\text{CH}_2)_{0-4}-\text{O}-(\text{R}_{\text{N}-5})_2-\text{COOH}$ where $\text{R}_{\text{N}-5}$ is as defined above,
- (33) $-(\text{CH}_2)_{0-4}-\text{S}-(\text{R}_{\text{N}-5})_2$ where $\text{R}_{\text{N}-5}$ is as defined above,
- 10 (34) $-(\text{CH}_2)_{0-4}-\text{O}-(\text{C}_1-\text{C}_6 \text{ alkyl optionally substituted with one, two, three, four, or five of } -\text{F}),$
- (35) $\text{C}_3-\text{C}_7 \text{ cycloalkyl},$
- (36) $\text{C}_2-\text{C}_6 \text{ alkenyl with one or two double bonds}$
- 15 optionally substituted with $\text{C}_1-\text{C}_3 \text{ alkyl}, -\text{F}, -\text{Cl}, -\text{Br}, -\text{I}, -\text{OH}, -\text{SH}, -\text{C}\equiv\text{N}, -\text{CF}_3, \text{C}_1-\text{C}_3 \text{ alkoxy}, \text{ or } -\text{NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,
- (37) $\text{C}_2-\text{C}_6 \text{ alkynyl with one or two triple bonds}$
- optionally substituted with $\text{C}_1-\text{C}_3 \text{ alkyl}, -\text{F}, -\text{Cl}, -\text{Br}, -\text{I}, -\text{OH}, -\text{SH}, -\text{C}\equiv\text{N}, -\text{CF}_3, \text{C}_1-\text{C}_3 \text{ alkoxy}, \text{ or } -\text{NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,
- 20 (38) $-(\text{CH}_2)_{0-4}-\text{N}(-\text{H} \text{ or } \text{R}_{\text{N}-5})-\text{SO}_2-\text{R}_{\text{N}-2}$ where $\text{R}_{\text{N}-5}$ and $\text{R}_{\text{N}-2}$ can be the same or different and are as defined above, or
- (39) $-(\text{CH}_2)_{0-4}-\text{C}_3-\text{C}_7 \text{ cycloalkyl},$
- (C) $\text{R}_{\text{N-aryl}}-\text{W}-\text{R}_{\text{N-aryl}},$ where $\text{R}_{\text{N-aryl}},$ and $\text{R}_{\text{N-aryl}}$ are as defined above,
- 25 (D) $\text{R}_{\text{N-aryl}}-\text{W}-\text{R}_{\text{N-heteroaryl}},$ where $\text{R}_{\text{N-aryl}},$ and $\text{R}_{\text{N-heteroaryl}},$ are as defined above,
- (E) $\text{R}_{\text{N-aryl}}-\text{W}-\text{R}_{\text{N-1-heterocycle}},$ wherein $\text{R}_{\text{N-1-heterocycle}}$ is the same as $\text{R}_{1-\text{heterocycle}},$ and $\text{R}_{1-\text{heterocycle}}$ is as defined above
- (F) $\text{R}_{\text{N-heteroaryl}}-\text{W}-\text{R}_{\text{N-aryl}},$ where $\text{R}_{\text{N-aryl}},$ and $\text{R}_{\text{N-heteroaryl}},$ are as defined above,
- 30 (G) $\text{R}_{\text{N-heteroaryl}}-\text{W}-\text{R}_{\text{N-heteroaryl}},$ where $\text{R}_{\text{N-heteroaryl}}$ is as defined above,

(H) $R_{N\text{-heteroaryl}}\text{-W-}R_{N\text{-1-heterocycle}}$, where $R_{N\text{-heteroaryl}}$, and $R_{N\text{-1-heterocycle}}$, are as defined above,

(I) $R_{N\text{-heterocycle}}\text{-W-}R_{N\text{-aryl}}$, wherein $R_{N\text{-heterocycle}}$ is the same as $R_{1\text{-heterocycle}}$, and $R_{1\text{-heterocycle}}$ is as defined above, and $R_{N\text{-aryl}}$ is as defined above,

5 (J) $R_{N\text{-heterocycle}}\text{-W-}R_{N\text{-heteroaryl}}$, where $R_{N\text{-heteroaryl}}$, and $R_{N\text{-heterocycle}}$, are as defined above, and

(K) $R_{N\text{-heterocycle}}\text{-W-}R_{N\text{-1-heterocycle}}$, where $R_{N\text{-heterocycle}}$, and $R_{N\text{-1-heterocycle}}$, are as defined above,

where W is

10 (5) $\text{-(CH}_2\text{)}_{0-4}\text{-}$,

(6) -O- ,

(7) $\text{-S(O)}_{0-2}\text{-}$,

(8) $\text{-N(R}_{N\text{-5}}\text{)-}$ where $R_{N\text{-5}}$ is as defined above, or

(5) -CO- ;

15 (II) $\text{-CO-(C}_1\text{-C}_{10}\text{ alkyl)}$ where alkyl is optionally substituted with one three substituents selected from the group consisting of:

(A) -OH ,

(B) $\text{-C}_1\text{-C}_6\text{ alkoxy}$,

(C) $\text{-C}_1\text{-C}_6\text{ thioalkoxy}$,

20 (D) $\text{-CO-O-}R_{N\text{-8}}$ where $R_{N\text{-8}}$ is -H , $\text{C}_1\text{-C}_6\text{ alkyl}$ or -phenyl ,

(E) $\text{-CO-NR}_{N\text{-2}}R_{N\text{-3}}$ where $R_{N\text{-2}}$ and $R_{N\text{-3}}$ are the same or different and are as defined above,

(F) $\text{-CO-}R_{N\text{-4}}$ where $R_{N\text{-4}}$ is as defined above,

(G) $\text{-SO}_2\text{-(C}_1\text{-C}_8\text{ alkyl)}$,

25 (H) $\text{-SO}_2\text{-NR}_{N\text{-2}}R_{N\text{-3}}$ where $R_{N\text{-2}}$ and $R_{N\text{-3}}$ are the same or different and are as defined above,

(I) $\text{-NH-CO-(C}_1\text{-C}_6\text{ alkyl)}$,

(J) $\text{-NH-CO-O-}R_{N\text{-8}}$ where $R_{N\text{-8}}$ is as defined above,

(K) $\text{-NR}_{N\text{-2}}R_{N\text{-3}}$ where $R_{N\text{-2}}$ and $R_{N\text{-3}}$ are the same or different

30 and are as defined above,

(L) $\text{-R}_{N\text{-4}}$ where $R_{N\text{-4}}$ is as defined above,

(M) $\text{-O-CO-(C}_1\text{-C}_6\text{ alkyl)}$,

(N) $\text{-O-CO-NR}_{N\text{-8}}R_{N\text{-8}}$ where $R_{N\text{-8}}$ are the same or different and are as defined above,

- (O) -O-(C₁-C₅ alkyl)-COOH,
 (P) -O-(C₁-C₆ alkyl optionally substituted with one, two, or
 three of -F, -Cl, -Br, -I),
 (Q) -NH-SO₂-(C₁-C₆ alkyl), and
 5 (R) -F, or -Cl,
 (III) -CO-(C₁-C₆ alkyl)-O-(C₁-C₆ alkyl) where alkyl is optionally
 substituted with one, two, or three of substituents selected from the group consisting
 of:
- (A) -OH,
 10 (B) -C₁-C₆ alkoxy,
 (C) -C₁-C₆ thioalkoxy,
 (D) -CO-O-R_{N-8} where R_{N-8} is -H, C₁-C₆ alkyl or -phenyl,
 (E) -CO-NR_{N-2}R_{N-3} where R_{N-2} and R_{N-3} are the same or
 different and are as defined above,
 15 (F) -CO-R_{N-4} where R_{N-4} is as defined above,
 (G) -SO₂-(C₁-C₈ alkyl),
 (H) -SO₂-NR_{N-2}R_{N-3} where R_{N-2} and R_{N-3} are the same or
 different and are as defined above,
 (I) -NH-CO-(C₁-C₆ alkyl),
 20 (J) -NH-CO-O-R_{N-8} where R_{N-8} is as defined above,
 (K) -NR_{N-2}R_{N-3} where R_{N-2} and R_{N-3} are the same or different
 and are as defined above,
 (L) -R_{N-4} where R_{N-4} is as defined above,
 (M) -O-CO-(C₁-C₆ alkyl),
 25 (N) -O-CO-NR_{N-8}R_{N-8} where R_{N-8} are the same or different
 and are as defined above,
 (O) -O-(C₁-C₅ alkyl)-COOH,
 (P) -O-(C₁-C₆ alkyl optionally substituted with one, two, or
 three of -F, -Cl, -Br, or -I),
 30 (Q) -NH-SO₂-(C₁-C₆ alkyl), and
 (R) -F, or -Cl,

(IV) $-\text{CO}-(\text{C}_1\text{-C}_6 \text{ alkyl})-\text{S}-(\text{C}_1\text{-C}_6 \text{ alkyl})$ where alkyl is optionally substituted with one, two, or three of substituents selected from the group consisting of:

- 5 (A) $-\text{OH}$,
 (B) $-\text{C}_1\text{-C}_6 \text{ alkoxy}$,
 (C) $-\text{C}_1\text{-C}_6 \text{ thioalkoxy}$,
 (D) $-\text{CO}-\text{O}-\text{R}_{\text{N-8}}$ where $\text{R}_{\text{N-8}}$ is as defined above,
 (E) $-\text{CO}-\text{NR}_{\text{N-2}}\text{R}_{\text{N-3}}$ where $\text{R}_{\text{N-2}}$ and $\text{R}_{\text{N-3}}$ are the same or different and are as defined above,
 10 (F) $-\text{CO}-\text{R}_{\text{N-4}}$ where $\text{R}_{\text{N-4}}$ is as defined above,
 (G) $-\text{SO}_2-(\text{C}_1\text{-C}_8 \text{ alkyl})$,
 (H) $-\text{SO}_2-\text{NR}_{\text{N-2}}\text{R}_{\text{N-3}}$ where $\text{R}_{\text{N-2}}$ and $\text{R}_{\text{N-3}}$ are the same or different and are as defined above,
 (I) $-\text{NH}-\text{CO}-(\text{C}_1\text{-C}_6 \text{ alkyl})$,
 15 (J) $-\text{NH}-\text{CO}-\text{O}-\text{R}_{\text{N-8}}$ where $\text{R}_{\text{N-8}}$ is as defined above,
 (K) $-\text{NR}_{\text{N-2}}\text{R}_{\text{N-3}}$ where $\text{R}_{\text{N-2}}$ and $\text{R}_{\text{N-3}}$ are the same or different and are as defined above,
 (L) $-\text{R}_{\text{N-4}}$ where $\text{R}_{\text{N-4}}$ is as defined above,
 (M) $-\text{O}-\text{CO}-(\text{C}_1\text{-C}_6 \text{ alkyl})$,
 20 (N) $-\text{O}-\text{CO}-\text{NR}_{\text{N-8}}\text{R}_{\text{N-8}}$ where $\text{R}_{\text{N-8}}$ are the same or different and are as defined above,
 (O) $-\text{O}-(\text{C}_1\text{-C}_5 \text{ alkyl})-\text{COOH}$,
 (P) $-\text{O}-(\text{C}_1\text{-C}_6 \text{ alkyl})$ optionally substituted with one, two, or three of $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, or $-\text{I}$,
 25 (Q) $-\text{NH}-\text{SO}_2-(\text{C}_1\text{-C}_6 \text{ alkyl})$, and
 (R) $-\text{F}$, or $-\text{Cl}$,
 (V) $-\text{CO}-\text{CH}(-(\text{CH}_2)_{0-2}-\text{O}-\text{R}_{\text{N-10}})-(\text{CH}_2)_{0-2}-\text{R}_{\text{N-aryl}}/\text{R}_{\text{N-heteroaryl}}$ where $\text{R}_{\text{N-aryl}}$ and $\text{R}_{\text{N-heteroaryl}}$ are as defined above, where $\text{R}_{\text{N-10}}$ is selected from the group consisting of:
 30 (A) $-\text{H}$,
 (B) $\text{C}_1\text{-C}_6 \text{ alkyl}$,
 (C) $\text{C}_3\text{-C}_7 \text{ cycloalkyl}$,
 (D) $\text{C}_2\text{-C}_6 \text{ alkenyl}$ with one double bond,

- (E) C₂-C₆ alkynyl with one triple bond,
 (F) R_{1-aryl} where R_{1-aryl} is as defined above, and
 (G) R_{N-heteroaryl} where R_{N-heteroaryl} is as defined above, or
 (VI) -CO-(C₃-C₈ cycloalkyl) where alkyl is optionally substituted
 5 with one or two substituents selected from the group consisting of:
 (A) -(CH₂)₀₋₄-OH,
 (B) -(CH₂)₀₋₄-C₁-C₆ alkoxy,
 (C) -(CH₂)₀₋₄-C₁-C₆ thioalkoxy,
 (D) -(CH₂)₀₋₄-CO-O-R_{N-8} where R_{N-8} is -H, C₁-C₆ alkyl or
 10 phenyl,
 (E) -(CH₂)₀₋₄-CO-NR_{N-2}R_{N-3} where R_{N-2} and R_{N-3} are the
 same or different and are as defined above,
 (F) -(CH₂)₀₋₄-CO-R_{N-4} where R_{N-4} is as defined above,
 (G) -(CH₂)₀₋₄-SO₂-(C₁-C₈ alkyl),
 15 (H) -(CH₂)₀₋₄-SO₂-NR_{N-2}R_{N-3} where R_{N-2} and R_{N-3} are the
 same or different and are as defined above,
 (I) -(CH₂)₀₋₄-NH-CO-(C₁-C₆ alkyl),
 (J) -NH-CO-O-R_{N-8} where R_{N-8} is as defined above,
 (K) -(CH₂)₀₋₄-NR_{N-2}R_{N-3} where R_{N-2} and R_{N-3} are the same or
 20 different and are as defined above,
 (L) -(CH₂)₀₋₄-R_{N-4} where R_{N-4} is as defined above,
 (M) -O-CO-(C₁-C₆ alkyl),
 (N) -O-CO-NR_{N-8}R_{N-8} where R_{N-8} are the same or different
 and are as defined above,
 25 (O) -O-(C₁-C₅ alkyl)-COOH,
 (P) -O-(C₁-C₆ alkyl optionally substituted with one, two, or
 three of -F, -Cl, -Br, or -I),
 (Q) -NH-SO₂-(C₁-C₆ alkyl), and
 (R) -F, or -Cl.
 30 The compound that is the source of R_N can be coupled with any well known
 coupling agents, an example of which is carbodiimide. Cleavage of the
 acylhydrazine linkage gives the compounds (XV).

CHART A' gives a more specific example of one method of synthesizing compounds of the invention (XV). The anti-Alzheimer's compounds of formula (XV) are prepared by starting with the corresponding epoxide (I). The epoxides (I) are well known to those skilled in the art or can be readily prepared from known compounds by methods well known to those skilled in the art. The compounds of formula (XV) of the present invention have at least two enantiomeric centers which give four enantiomers. The first of these enantiomeric centers derives from the epoxide starting material (I). If a desired enantiomer is preferred, it is preferred to commercially obtain or produce the desired enantiomer (S or R) rather than produce an enantiomerically impure mixture and then have to separate out the desired enantiomer.

The exemplary synthesis begins by reacting the epoxide (I) with an aromatic hydrazine in hot isopropanol results in the selective formation of the hydrazines (II) arising from alkylation of the unsubstituted nitrogen (M. Nakakata, *Tetrahedron Letters* **1993**, 6095-6098). Monoacylation of the hydrazine -NH-NH- with benzyloxycarbonyl chloride or other acylating agent gives (III) and reduces the reactivity of this group to further acylation irrespective of which hydrazine nitrogen the first acyl group becomes attached to (B. Gisin, *Helv. Chim. Acta* **1970**, vol 53, 1030-1043. S. Shinagawa, *Chem. Pharm. Bull.* **1981**, vol 29, 3630-3638). Removal of the *tert*-butoxycarbonyl protecting group of (III) will provide the free amine (IV), which is coupled to the isophthalic acid (XIV) using carbodiimide or other well known coupling agents. Cleavage of the acylhydrazine linkage gives a compound of the invention (XV).

CHART B offers another example of a method that can be utilized to make compounds of the invention. Selective acylation of methylhydrazine on the substituted nitrogen (D. Butler, *J. Medicinal Chemistry* **1971**, vol. 14, 1052-1054) will provide acylhydrazine VI. Treating this hydrazide with epoxide I in hot isopropanol will provide adduct VII (S. Wang, *J. Medicinal Chemistry* **1997**, vol 40, 937-941. G. Bold, *J. Medicinal Chemistry* **1998**, vol 41, 3387-3401). Cleavage of the *tert*-butoxycarbonyl protecting group and coupling to isophthalic acid (XIV) will provide a compound of the invention (XV).

CHART C offers a general method of making compounds (XV) of the invention, wherein X is O. A general method of synthesizing compounds (XV) of the invention wherein X is O, begins with a protected epoxide (I). The epoxide (I)

again serves to provide R_1 , R_2 , and R_3 of the final product (XV), the discussion of these compounds offered above applies equally here. The epoxide is opened with a hydroxylamine having the formula R_A-O-NH_2 . The hydroxylamine serves both to open the epoxide ring and provide R_A to the final product (XV). Once the hydroxylamine has been reacted with the epoxide (I), the adduct (XI) is formed. Adduct (XI) has R_1 , R_2 , R_3 , and R_A of the compounds (XV) of the invention. The possible identities of R_1 , R_2 , R_3 , and R_A , as well as the protecting group discussed above, apply to adduct (XI) as well. The next step in the synthesis of compounds (XV) of the invention, wherein X is O is cleavage of the protecting group. The protecting groups and methods of cleaving them discussed above apply similarly to these compounds. After the protecting group has been cleaved from adduct XI, the next step involves acylation with the source of R_N .

CHART C' offers another more specific illustrative example of one method of making compounds (XV) of the invention, wherein X is O. Epoxide (I) opening with O-benzylhydroxylamine gives the adduct XI (S. Rosenberg, *J. Medicinal Chemistry* **1990**, vol 33, 1582-1590). Cleavage of the *tert*-butoxycarbonyl protecting group and acylation with isophthallic acid (as prepared, for example, by the method below) provides the target compound XIII.

The preparation of isophthallic acid for use in the above synthesis can be accomplished for example, by the below synthesis, referring to CHART D below. Methyl isophthalate (1 equiv, 11.1 mmol) was dissolved in 50:50 THF:DMF (20 mL) before the addition of 1,1'-carbonyldiimidazole (CDI) (1.2 equiv, 13.3 mmol) at ambient temperature. Upon addition of CDI, a color change from colorless to yellow, as well as evolution of gas (CO_2), was observed. After gas evolution subsided (approximately one minute or less), the amine (1.2 equiv, 13.3 mmol) dissolved in DMF and diisopropylethyl amine (1.2 equiv, 13.3 mmol) was added. After 12 hours of stirring at ambient temperature, the reaction was partitioned between saturated aqueous NH_4Cl and ethyl acetate, and the aqueous layer was extracted twice more with ethyl acetate. The organic extracts were then washed with saturated aqueous solutions of $NaHCO_3$ and $NaCl$, and dried over anhydrous $MgSO_4$ or $NaSO_4$. Filtration of the drying agent and removal of solvents *in vacuo* gave the crude white solid or clear oil. Purification of these compounds *if needed* was achieved via chromatography on silica gel with 30–40% ethyl acetate in hexanes (80-90% yield).

The methyl isophthalate mono-alkyl or di-alkyl amide (1 equiv, 11.1 mmol) was then treated with LiOH·H₂O (3 equiv, 33.3 mmol) in a minimum amount of 1:2:1 THF:MeOH:H₂O and allowed to stir overnight at ambient temperature. After 12 hours, the solvents were removed *in vacuo* and subsequently partitioned between H₂O and ethyl acetate. If emulsions prohibit separation of the two layers, a small amount of brine was added to aid in separation. The aqueous layer was extracted once more with ethyl acetate (to remove any unreacted starting material). The aqueous layer was then acidified with concentrated HCl until pH ≤ 3. The cloudy-white acidic aqueous solution thus obtained was then extracted three times with ethyl acetate. These combined organic extracts were dried over anhydrous MgSO₄ or Na₂SO₄. Filtration of the drying agent and removal of solvents *in vacuo* gave the crude white solid. The mono- or di-alkyl amide isophthalate was used crude in the next reaction (90-100% yield).

Compounds of the invention may contain geometric or optical isomers as well as tautomers. Thus, the invention includes all tautomers and pure geometric isomers, such as the *E* and *Z* geometric isomers, as well as mixtures thereof. Furthermore, the invention includes pure enantiomers and diastereomers as well as mixtures thereof, including racemic mixtures. The individual geometric isomers, enantiomers, or diastereomers may be prepared or isolated by methods known in the art.

Compounds of the invention with the stereochemistry designated in formula XV may be included in mixtures, including racemic mixtures, with other enantiomers, diastereomers, geometric isomers or tautomers. Compounds of the invention with the stereochemistry designated in formula XV are typically present in these mixtures in excess of 50 percent. Preferably, compounds of the invention with the stereochemistry designated in formula XV are present in these mixtures in excess of 80 percent. Most preferably, compounds of the invention with the stereochemistry designated in formula XV are present in these mixtures in excess of 90 percent.

The (S,R)-substituted amines (XV) are amines and as such form salts when reacted with acids. Pharmaceutically acceptable salts are preferred over the corresponding (S,R)-substituted amines (XV) since they produce compounds which are more water soluble, stable and/or more crystalline.

Pharmaceutically acceptable salts are any salt which retains the activity of the parent compound and does not impart any deleterious or undesirable effect on the subject to whom it is administered and in the context in which it is administered. Pharmaceutically acceptable salts include salts of both inorganic and organic acids.

- 5 The preferred pharmaceutically acceptable salts include salts of the following acids acetic, aspartic, benzenesulfonic, benzoic, bicarbonic, bisulfuric, bitartaric, butyric, calcium edetate, camsyllic, carbonic, chlorobenzoic, citric, edetic, edisylic, estolic, esyl, esylic, formic, fumaric, gluceptic, gluconic, glutamic, glycolylarsanilic, hexamic, hexylresorcinoic, hydrabamic, hydrobromic, hydrochloric, hydroiodic, 10 hydroxynaphthoic, isethionic, lactic, lactobionic, maleic, malic, malonic, mandelic, methanesulfonic, methylnitric, methylsulfuric, mucic, muconic, napsylic, nitric, oxalic, p-nitromethanesulfonic, pamoic, pantothenic, phosphoric, monohydrogen phosphoric, dihydrogen phosphoric, phthalic, polygalactouronic, propionic, salicylic, stearic, succinic, succinic, sulfamic, sulfanilic, sulfonic, sulfuric, tannic, 15 tartaric, teoclic and toluenesulfonic. For other acceptable salts, see *Int. J. Pharm.*, 33, 201-217 (1986) and *J.Pharm.Sci.*, 66(1), 1, (1977).

The present invention provides compounds, compositions, kits, and methods for inhibiting beta-secretase enzyme activity and A beta peptide production.

- Inhibition of beta-secretase enzyme activity halts or reduces the production of A 20 beta from APP and reduces or eliminates the formation of beta-amyloid deposits in the brain.

Methods of the Invention

- The compounds employed in the methods of the invention, and 25 pharmaceutically acceptable salts thereof, are useful for treating humans or animals suffering from a condition characterized by a pathological form of beta-amyloid peptide, such as beta-amyloid plaques, and for helping to prevent or delay the onset of such a condition. For example, the compounds are useful for treating Alzheimer's disease, for helping prevent or delay the onset of Alzheimer's disease, for treating 30 patients with MCI (mild cognitive impairment) and preventing or delaying the onset of Alzheimer's disease in those who would progress from MCI to AD, for treating Down's syndrome, for treating humans who have Hereditary Cerebral Hemorrhage with Amyloidosis of the Dutch-Type, for treating cerebral amyloid angiopathy and preventing its potential consequences, i.e. single and recurrent lobal hemorrhages,

for treating other degenerative dementias, including dementias of mixed vascular and degenerative origin, dementia associated with Parkinson's disease, dementia associated with progressive supranuclear palsy, dementia associated with cortical basal degeneration, and diffuse Lewy body type Alzheimer's disease. The
5 compounds and compositions of the invention are particularly useful for treating or preventing Alzheimer's disease. When treating or preventing these diseases, the compounds employed in the methods of the invention can either be used individually or in combination, as is best for the patient.

As used herein, the term "treating" means that the compounds employed in
10 the methods of the invention can be used in humans with at least a tentative diagnosis of disease. The compounds employed in the methods of the invention will delay or slow the progression of the disease thereby giving the individual a more useful life span.

The term "preventing" means that the compounds employed in the method of
15 the invention are useful when administered to a patient who has not been diagnosed as possibly having the disease at the time of administration, but who would normally be expected to develop the disease or be at increased risk for the disease. The compounds employed in the methods of the invention will slow the development of disease symptoms, delay the onset of the disease, or prevent the individual from
20 developing the disease at all. Preventing also includes administration of the compounds employed in the methods of the invention to those individuals thought to be predisposed to the disease due to age, familial history, genetic or chromosomal abnormalities, and/or due to the presence of one or more biological markers for the disease, such as a known genetic mutation of APP or APP cleavage products in brain
25 tissues or fluids.

In treating or preventing the above diseases, the compounds employed in the methods of the invention are administered in a therapeutically effective amount. The therapeutically effective amount will vary depending on the particular compound used and the route of administration, as is known to those skilled in the
30 art.

In treating a patient displaying any of the diagnosed above conditions a physician may administer a compound employed in the method of the invention immediately and continue administration indefinitely, as needed. In treating patients who are not diagnosed as having Alzheimer's disease, but who are believed to be at

substantial risk for Alzheimer's disease, the physician should preferably start treatment when the patient first experiences early pre-Alzheimer's symptoms such as, memory or cognitive problems associated with aging. In addition, there are some patients who may be determined to be at risk for developing Alzheimer's through the detection of a genetic marker such as APOE4 or other biological indicators that are predictive for Alzheimer's disease. In these situations, even though the patient does not have symptoms of the disease, administration of the compounds employed in the methods of the invention may be started before symptoms appear, and treatment may be continued indefinitely to prevent or delay the onset of the disease.

Dosage Forms and Amounts

The compounds employed in the methods of the invention can be administered orally, parenterally, (IV, IM, depo-IM, SQ, and depo SQ), sublingually, intranasally (inhalation), intrathecally, topically, or rectally. Dosage forms known to those of skill in the art are suitable for delivery of the compounds employed in the methods of the invention.

Compositions are provided that contain therapeutically effective amounts of the compounds employed in the methods of the invention. The compounds are preferably formulated into suitable pharmaceutical preparations such as tablets, capsules, or elixirs for oral administration or in sterile solutions or suspensions for parenteral administration. Typically the compounds described above are formulated into pharmaceutical compositions using techniques and procedures well known in the art.

About 1 to 500 mg of a compound or mixture of compounds employed in the methods of the invention or a physiologically acceptable salt or ester is compounded with a physiologically acceptable vehicle, carrier, excipient, binder, preservative, stabilizer, flavor, etc., in a unit dosage form as called for by accepted pharmaceutical practice. The amount of active substance in those compositions or preparations is such that a suitable dosage in the range indicated is obtained. The compositions are preferably formulated in a unit dosage form, each dosage containing from about 2 to about 100 mg, more preferably about 10 to about 30 mg of the active ingredient. The term "unit dosage form" refers to physically discrete units suitable as unitary dosages for human subjects and other mammals, each unit containing a

predetermined quantity of active material calculated to produce the desired therapeutic effect, in association with a suitable pharmaceutical excipient.

To prepare compositions, one or more compounds employed in the methods of the invention are mixed with a suitable pharmaceutically acceptable carrier.

- 5 Upon mixing or addition of the compound(s), the resulting mixture may be a solution, suspension, emulsion, or the like. Liposomal suspensions may also be suitable as pharmaceutically acceptable carriers. These may be prepared according to methods known to those skilled in the art. The form of the resulting mixture depends upon a number of factors, including the intended mode of administration and the solubility of the compound in the selected carrier or vehicle. The effective concentration is sufficient for lessening or ameliorating at least one symptom of the disease, disorder, or condition treated and may be empirically determined.

- Pharmaceutical carriers or vehicles suitable for administration of the compounds provided herein include any such carriers known to those skilled in the art to be suitable for the particular mode of administration. In addition, the active materials can also be mixed with other active materials that do not impair the desired action, or with materials that supplement the desired action, or have another action. The compounds may be formulated as the sole pharmaceutically active ingredient in the composition or may be combined with other active ingredients.

- 20 Where the compounds exhibit insufficient solubility, methods for solubilizing may be used. Such methods are known and include, but are not limited to, using cosolvents such as dimethylsulfoxide (DMSO), using surfactants such as Tween®, and dissolution in aqueous sodium bicarbonate. Derivatives of the compounds, such as salts or prodrugs may also be used in formulating effective pharmaceutical compositions.

The concentration of the compound is effective for delivery of an amount upon administration that lessens or ameliorates at least one symptom of the disorder for which the compound is administered. Typically, the compositions are formulated for single dosage administration.

- 30 The compounds employed in the methods of the invention may be prepared with carriers that protect them against rapid elimination from the body, such as time-release formulations or coatings. Such carriers include controlled release formulations, such as, but not limited to, microencapsulated delivery systems. The active compound is included in the pharmaceutically acceptable carrier in an amount

sufficient to exert a therapeutically useful effect in the absence of undesirable side effects on the patient treated. The therapeutically effective concentration may be determined empirically by testing the compounds in known *in vitro* and *in vivo* model systems for the treated disorder.

5 The compounds and compositions of the invention can be enclosed in multiple or single dose containers. The enclosed compounds and compositions can be provided in kits, for example, including component parts that can be assembled for use. For example, a compound inhibitor in lyophilized form and a suitable diluent may be provided as separated components for combination prior to use. A
10 kit may include a compound inhibitor and a second therapeutic agent for co-administration. The inhibitor and second therapeutic agent may be provided as separate component parts. A kit may include a plurality of containers, each container holding one or more unit dose of the compound employed in the method of the invention. The containers are preferably adapted for the desired mode of
15 administration, including, but not limited to tablets, gel capsules, sustained-release capsules, and the like for oral administration; depot products, pre-filled syringes, ampoules, vials, and the like for parenteral administration; and patches, medipads, creams, and the like for topical administration.

 The concentration of active compound in the drug composition will depend
20 on absorption, inactivation, and excretion rates of the active compound, the dosage schedule, and amount administered as well as other factors known to those of skill in the art.

 The active ingredient may be administered at once, or may be divided into a number of smaller doses to be administered at intervals of time. It is understood that
25 the precise dosage and duration of treatment is a function of the disease being treated and may be determined empirically using known testing protocols or by extrapolation from *in vivo* or *in vitro* test data. It is to be noted that concentrations and dosage values may also vary with the severity of the condition to be alleviated. It is to be further understood that for any particular subject, specific dosage regimens
30 should be adjusted over time according to the individual need and the professional judgment of the person administering or supervising the administration of the compositions, and that the concentration ranges set forth herein are exemplary only and are not intended to limit the scope or practice of the claimed compositions.

If oral administration is desired, the compound should be provided in a composition that protects it from the acidic environment of the stomach. For example, the composition can be formulated in an enteric coating that maintains its integrity in the stomach and releases the active compound in the intestine. The composition may also be formulated in combination with an antacid or other such ingredient.

Oral compositions will generally include an inert diluent or an edible carrier and may be compressed into tablets or enclosed in gelatin capsules. For the purpose of oral therapeutic administration, the active compound or compounds can be incorporated with excipients and used in the form of tablets, capsules, or troches. Pharmaceutically compatible binding agents and adjuvant materials can be included as part of the composition.

The tablets, pills, capsules, troches, and the like can contain any of the following ingredients or compounds of a similar nature: a binder such as, but not limited to, gum tragacanth, acacia, corn starch, or gelatin; an excipient such as microcrystalline cellulose, starch, or lactose; a disintegrating agent such as, but not limited to, alginic acid and corn starch; a lubricant such as, but not limited to, magnesium stearate; a gildant, such as, but not limited to, colloidal silicon dioxide; a sweetening agent such as sucrose or saccharin; and a flavoring agent such as peppermint, methyl salicylate, or fruit flavoring.

When the dosage unit form is a capsule, it can contain, in addition to material of the above type, a liquid carrier such as a fatty oil. In addition, dosage unit forms can contain various other materials, which modify the physical form of the dosage unit, for example, coatings of sugar and other enteric agents. The compounds can also be administered as a component of an elixir, suspension, syrup, wafer, chewing gum or the like. A syrup may contain, in addition to the active compounds, sucrose as a sweetening agent and certain preservatives, dyes and colorings, and flavors.

The active materials can also be mixed with other active materials that do not impair the desired action, or with materials that supplement the desired action.

Solutions or suspensions used for parenteral, intradermal, subcutaneous, or topical application can include any of the following components: a sterile diluent such as water for injection, saline solution, fixed oil, a naturally occurring vegetable oil such as sesame oil, coconut oil, peanut oil, cottonseed oil, and the like, or a synthetic fatty vehicle such as ethyl oleate, and the like, polyethylene glycol,

glycerine, propylene glycol, or other synthetic solvent; antimicrobial agents such as benzyl alcohol and methyl parabens; antioxidants such as ascorbic acid and sodium bisulfite; chelating agents such as ethylenediaminetetraacetic acid (EDTA); buffers such as acetates, citrates, and phosphates; and agents for the adjustment of tonicity
5 such as sodium chloride and dextrose. Parenteral preparations can be enclosed in ampoules, disposable syringes, or multiple dose vials made of glass, plastic, or other suitable material. Buffers, preservatives, antioxidants, and the like can be incorporated as required.

Where administered intravenously, suitable carriers include physiological
10 saline, phosphate buffered saline (PBS), and solutions containing thickening and solubilizing agents such as glucose, polyethylene glycol, polypropyleneglycol, and mixtures thereof. Liposomal suspensions including tissue-targeted liposomes may also be suitable as pharmaceutically acceptable carriers. These may be prepared according to methods known for example, as described in U.S. Patent No.
15 4,522,811.

The active compounds may be prepared with carriers that protect the compound against rapid elimination from the body, such as time-release formulations or coatings. Such carriers include controlled release formulations, such as, but not limited to, implants and microencapsulated delivery systems, and
20 biodegradable, biocompatible polymers such as collagen, ethylene vinyl acetate, polyanhydrides, polyglycolic acid, polyorthoesters, polylactic acid, and the like. Methods for preparation of such formulations are known to those skilled in the art.

The compounds employed in the methods of the invention can be administered orally, parenterally (IV, IM, depo-IM, SQ, and depo-SQ), sublingually,
25 intranasally (inhalation), intrathecally, topically, or rectally. Dosage forms known to those skilled in the art are suitable for delivery of the compounds employed in the methods of the invention.

Compounds employed in the methods of the invention may be administered enterally or parenterally. When administered orally, compounds employed in the
30 methods of the invention can be administered in usual dosage forms for oral administration as is well known to those skilled in the art. These dosage forms include the usual solid unit dosage forms of tablets and capsules as well as liquid dosage forms such as solutions, suspensions, and elixirs. When the solid dosage forms are used, it is preferred that they be of the sustained release type so that the

compounds employed in the methods of the invention need to be administered only once or twice daily.

The oral dosage forms are administered to the patient 1, 2, 3, or 4 times daily. It is preferred that the compounds employed in the methods of the invention be administered either three or fewer times, more preferably once or twice daily. Hence, it is preferred that the compounds employed in the methods of the invention be administered in oral dosage form. It is preferred that whatever oral dosage form is used, that it be designed so as to protect the compounds employed in the methods of the invention from the acidic environment of the stomach. Enteric coated tablets are well known to those skilled in the art. In addition, capsules filled with small spheres each coated to protect from the acidic stomach, are also well known to those skilled in the art.

When administered orally, an administered amount therapeutically effective to inhibit beta-secretase activity, to inhibit A beta production, to inhibit A beta deposition, or to treat or prevent AD is from about 0.1 mg/day to about 1,000 mg/day. It is preferred that the oral dosage is from about 1 mg/day to about 100 mg/day. It is more preferred that the oral dosage is from about 5 mg/day to about 50 mg/day. It is understood that while a patient may be started at one dose, that dose may be varied over time as the patient's condition changes.

Compounds employed in the methods of the invention may also be advantageously delivered in a nano crystal dispersion formulation. Preparation of such formulations is described, for example, in U.S. Patent 5,145,684. Nano crystalline dispersions of HIV protease inhibitors and their method of use are described in U.S. Patent No. 6,045,829. The nano crystalline formulations typically afford greater bioavailability of drug compounds.

The compounds employed in the methods of the invention can be administered parenterally, for example, by IV, IM, depo-IM, SC, or depo-SC. When administered parenterally, a therapeutically effective amount of about 0.5 to about 100 mg/day, preferably from about 5 to about 50 mg daily should be delivered. When a depot formulation is used for injection once a month or once every two weeks, the dose should be about 0.5 mg/day to about 50 mg/day, or a monthly dose of from about 15 mg to about 1,500 mg. In part because of the forgetfulness of the patients with Alzheimer's disease, it is preferred that the parenteral dosage form be a depo formulation.

The compounds employed in the methods of the invention can be administered sublingually. When given sublingually, the compounds employed in the methods of the invention should be given one to four times daily in the amounts described above for IM administration.

5 The compounds employed in the methods of the invention can be administered intranasally. When given by this route, the appropriate dosage forms are a nasal spray or dry powder, as is known to those skilled in the art. The dosage of the compounds employed in the methods of the invention for intranasal administration is the amount described above for IM administration.

10 The compounds employed in the methods of the invention can be administered intrathecally. When given by this route the appropriate dosage form can be a parenteral dosage form as is known to those skilled in the art. The dosage of the compounds employed in the methods of the invention for intrathecal administration is the amount described above for IM administration.

15 The compounds employed in the methods of the invention can be administered topically. When given by this route, the appropriate dosage form is a cream, ointment, or patch. Because of the amount of the compounds employed in the methods of the invention to be administered, the patch is preferred. When administered topically, the dosage is from about 0.5 mg/day to about 200 mg/day.

20 Because the amount that can be delivered by a patch is limited, two or more patches may be used. The number and size of the patch is not important, what is important is that a therapeutically effective amount of the compounds employed in the methods of the invention be delivered as is known to those skilled in the art. The compounds employed in the methods of the invention can be administered rectally
25 by suppository as is known to those skilled in the art. When administered by suppository, the therapeutically effective amount is from about 0.5 mg to about 500 mg.

 The compounds employed in the methods of the invention can be administered by implants as is known to those skilled in the art. When
30 administering a compound employed in the method of the invention by implant, the therapeutically effective amount is the amount described above for depot administration.

 The invention here is the new compounds employed in the methods of the invention and new methods of using the compounds employed in the methods of the

invention. Given a particular compound employed in the method of the invention and a desired dosage form, one skilled in the art would know how to prepare and administer the appropriate dosage form.

The compounds employed in the methods of the invention are used in the same manner, by the same routes of administration, using the same pharmaceutical dosage forms, and at the same dosing schedule as described above, for preventing disease or treating patients with MCI (mild cognitive impairment) and preventing or delaying the onset of Alzheimer's disease in those who would progress from MCI to AD, for treating or preventing Down's syndrome, for treating humans who have Hereditary Cerebral Hemorrhage with Amyloidosis of the Dutch-Type, for treating cerebral amyloid angiopathy and preventing its potential consequences, i.e. single and recurrent lobar hemorrhages, for treating other degenerative dementias, including dementias of mixed vascular and degenerative origin, dementia associated with Parkinson's disease, dementia associated with progressive supranuclear palsy, dementia associated with cortical basal degeneration, and diffuse Lewy body type of Alzheimer's disease.

The compounds employed in the methods of the invention can be used in combination, with each other or with other therapeutic agents or approaches used to treat or prevent the conditions listed above. Such agents or approaches include: acetylcholine esterase inhibitors such as tacrine (tetrahydroaminoacridine, marketed as COGNEX®), donepezil hydrochloride, (marketed as Aricept® and rivastigmine (marketed as Exelon®); gamma-secretase inhibitors; anti-inflammatory agents such as cyclooxygenase II inhibitors; anti-oxidants such as Vitamin E and ginkgolides; immunological approaches, such as, for example, immunization with A beta peptide or administration of anti-A beta peptide antibodies; statins; and direct or indirect neurotropic agents such as Cerebrolysin®, AIT-082 (Emilieu, 2000, *Arch. Neurol.* 57:454), and other neurotropic agents of the future.

It should be apparent to one skilled in the art that the exact dosage and frequency of administration will depend on the particular compounds employed in the methods of the invention administered, the particular condition being treated, the severity of the condition being treated, the age, weight, general physical condition of the particular patient, and other medication the individual may be taking as is well known to administering physicians who are skilled in this art.

Inhibition of APP Cleavage

The compounds employed in the methods of the invention inhibit cleavage of APP between Met595 and Asp596 numbered for the APP695 isoform, or a mutant thereof, or at a corresponding site of a different isoform, such as APP751 or APP770, or a mutant thereof (sometimes referred to as the "beta secretase site"). While not wishing to be bound by a particular theory, inhibition of beta-secretase activity is thought to inhibit production of beta amyloid peptide (A beta). Inhibitory activity is demonstrated in one of a variety of inhibition assays, whereby cleavage of an APP substrate in the presence of a beta-secretase enzyme is analyzed in the presence of the inhibitory compound, under conditions normally sufficient to result in cleavage at the beta-secretase cleavage site. Reduction of APP cleavage at the beta-secretase cleavage site compared with an untreated or inactive control is correlated with inhibitory activity. Assay systems that can be used to demonstrate efficacy of the compound inhibitors of the invention are known. Representative assay systems are described, for example, in U.S. Patents No. 5,942,400, 5,744,346, as well as in the Examples below.

The enzymatic activity of beta-secretase and the production of A beta can be analyzed *in vitro* or *in vivo*, using natural, mutated, and/or synthetic APP substrates, natural, mutated, and/or synthetic enzyme, and the test compound. The analysis may involve primary or secondary cells expressing native, mutant, and/or synthetic APP and enzyme, animal models expressing native APP and enzyme, or may utilize transgenic animal models expressing the substrate and enzyme. Detection of enzymatic activity can be by analysis of one or more of the cleavage products, for example, by immunoassay, fluorometric or chromogenic assay, HPLC, or other means of detection. Inhibitory compounds are determined as those having the ability to decrease the amount of beta-secretase cleavage product produced in comparison to a control, where beta-secretase mediated cleavage in the reaction system is observed and measured in the absence of inhibitory compounds.

Beta-Secretase

Various forms of beta-secretase enzyme are known, and are available and useful for assay of enzyme activity and inhibition of enzyme activity. These include native, recombinant, and synthetic forms of the enzyme. Human beta-secretase is known as Beta Site APP Cleaving Enzyme (BACE), Asp2, and memapsin 2, and has

been characterized, for example, in U.S. Patent No. 5,744,346 and published PCT patent applications WO98/22597, WO00/03819, WO01/23533, and WO00/17369, as well as in literature publications (Hussain et al., 1999, *Mol. Cell. Neurosci.* 14:419-427; Vassar et al., 1999, *Science* 286:735-741; Yan et al., 1999, *Nature* 402:533-537; Sinha et al., 1999, *Nature* 40:537-540; and Lin et al., 2000, *PNAS USA* 97:1456-1460). Synthetic forms of the enzyme have also been described (WO98/22597 and WO00/17369). Beta-secretase can be extracted and purified from human brain tissue and can be produced in cells, for example mammalian cells expressing recombinant enzyme.

Useful inhibitory compounds are effective to inhibit 50% of beta-secretase enzymatic activity at a concentration of less than 50 micromolar, preferably at a concentration of 10 micromolar or less, more preferably 1 micromolar or less, and most preferably 10 nanomolar or less.

APP Substrate

Assays that demonstrate inhibition of beta-secretase-mediated cleavage of APP can utilize any of the known forms of APP, including the 695 amino acid "normal" isotype described by Kang et al., 1987, *Nature* 325:733-6, the 770 amino acid isotype described by Kitaguchi et. al., 1981, *Nature* 331:530-532, and variants such as the Swedish Mutation (KM670-1NL) (APP-SW), the London Mutation (V7176F), and others. See, for example, U.S. Patent No. 5,766,846 and also Hardy, 1992, *Nature Genet.* 1:233-234, for a review of known variant mutations. Additional useful substrates include the dibasic amino acid modification, APP-KK disclosed, for example, in WO 00/17369, fragments of APP, and synthetic peptides containing the beta-secretase cleavage site, wild type (WT) or mutated form, e.g., SW, as described, for example, in U.S. Patent No 5,942,400 and WO00/03819.

The APP substrate contains the beta-secretase cleavage site of APP (KM-DA or NL-DA) for example, a complete APP peptide or variant, an APP fragment, a recombinant or synthetic APP, or a fusion peptide. Preferably, the fusion peptide includes the beta-secretase cleavage site fused to a peptide having a moiety useful for enzymatic assay, for example, having isolation and/or detection properties. A useful moiety may be an antigenic epitope for antibody binding, a label or other detection moiety, a binding substrate, and the like.

Antibodies

Products characteristic of APP cleavage can be measured by immunoassay using various antibodies, as described, for example, in Pirttila et al., 1999, *Neuro. Lett.* 249:21-4, and in U.S. Patent No. 5,612,486. Useful antibodies to detect A beta include, for example, the monoclonal antibody 6E10 (Senetek, St. Louis, MO) that specifically recognizes an epitope on amino acids 1-16 of the A beta peptide; antibodies 162 and 164 (New York State Institute for Basic Research, Staten Island, NY) that are specific for human A beta 1-40 and 1-42, respectively; and antibodies that recognize the junction region of beta-amyloid peptide, the site between residues 16 and 17, as described in U.S. Patent No. 5,593,846. Antibodies raised against a synthetic peptide of residues 591 to 596 of APP and SW192 antibody raised against 590-596 of the Swedish mutation are also useful in immunoassay of APP and its cleavage products, as described in U.S. Patent Nos. 5,604,102 and 5,721,130.

Assay Systems

Assays for determining APP cleavage at the beta-secretase cleavage site are well known in the art. Exemplary assays, are described, for example, in U.S. Patent Nos. 5,744,346 and 5,942,400, and described in the Examples below.

Cell Free Assays

Exemplary assays that can be used to demonstrate the inhibitory activity of the compounds employed in the methods of the invention are described, for example, in WO00/17369, WO 00/03819, and U.S. Patents No. 5,942,400 and 5,744,346. Such assays can be performed in cell-free incubations or in cellular incubations using cells expressing a beta-secretase and an APP substrate having a beta-secretase cleavage site.

An APP substrate containing the beta-secretase cleavage site of APP, for example, a complete APP or variant, an APP fragment, or a recombinant or synthetic APP substrate containing the amino acid sequence: KM-DA or NL-DA, is incubated in the presence of beta-secretase enzyme, a fragment thereof, or a synthetic or recombinant polypeptide variant having beta-secretase activity and effective to cleave the beta-secretase cleavage site of APP, under incubation conditions suitable for the cleavage activity of the enzyme. Suitable substrates optionally include derivatives that may be fusion proteins or peptides that contain

the substrate peptide and a modification useful to facilitate the purification or detection of the peptide or its beta-secretase cleavage products. Useful modifications include the insertion of a known antigenic epitope for antibody binding; the linking of a label or detectable moiety, the linking of a binding substrate, and the like.

Suitable incubation conditions for a cell-free *in vitro* assay include, for example: approximately 200 nanomolar to 10 micromolar substrate, approximately 10 to 200 picomolar enzyme, and approximately 0.1 nanomolar to 10 micromolar inhibitor compound, in aqueous solution, at an approximate pH of 4 -7, at approximately 37 degrees C, for a time period of approximately 10 minutes to 3 hours. These incubation conditions are exemplary only, and can be varied as required for the particular assay components and/or desired measurement system. Optimization of the incubation conditions for the particular assay components should account for the specific beta-secretase enzyme used and its pH optimum, any additional enzymes and/or markers that might be used in the assay, and the like. Such optimization is routine and will not require undue experimentation.

One useful assay utilizes a fusion peptide having maltose binding protein (MBP) fused to the C-terminal 125 amino acids of APP-SW. The MBP portion is captured on an assay substrate by anti-MBP capture antibody. Incubation of the captured fusion protein in the presence of beta-secretase results in cleavage of the substrate at the beta-secretase cleavage site. Analysis of the cleavage activity can be, for example, by immunoassay of cleavage products. One such immunoassay detects a unique epitope exposed at the carboxy terminus of the cleaved fusion protein, for example, using the antibody SW192. This assay is described, for example, in U.S. Patent No 5,942,400.

Cellular Assay

Numerous cell-based assays can be used to analyze beta-secretase activity and/or processing of APP to release A beta. Contact of an APP substrate with a beta-secretase enzyme within the cell and in the presence or absence of a compound inhibitor of the invention can be used to demonstrate beta-secretase inhibitory activity of the compound. Preferably, assay in the presence of a useful inhibitory compound provides at least about 30%, most preferably at least about 50% inhibition of the enzymatic activity, as compared with a non-inhibited control.

In one embodiment, cells that naturally express beta-secretase are used.

Alternatively, cells are modified to express a recombinant beta-secretase or synthetic variant enzyme as discussed above. The APP substrate may be added to the culture medium and is preferably expressed in the cells. Cells that naturally express APP, variant or mutant forms of APP, or cells transformed to express an isoform of APP, mutant or variant APP, recombinant or synthetic APP, APP fragment, or synthetic APP peptide or fusion protein containing the beta-secretase APP cleavage site can be used, provided that the expressed APP is permitted to contact the enzyme and enzymatic cleavage activity can be analyzed.

Human cell lines that normally process A beta from APP provide a useful means to assay inhibitory activities of the compounds employed in the methods of the invention. Production and release of A beta and/or other cleavage products into the culture medium can be measured, for example by immunoassay, such as Western blot or enzyme-linked immunoassay (EIA) such as by ELISA.

Cells expressing an APP substrate and an active beta-secretase can be incubated in the presence of a compound inhibitor to demonstrate inhibition of enzymatic activity as compared with a control. Activity of beta-secretase can be measured by analysis of one or more cleavage products of the APP substrate. For example, inhibition of beta-secretase activity against the substrate APP would be expected to decrease release of specific beta-secretase induced APP cleavage products such as A beta.

Although both neural and non-neural cells process and release A beta, levels of endogenous beta-secretase activity are low and often difficult to detect by EIA. The use of cell types known to have enhanced beta-secretase activity, enhanced processing of APP to A beta, and/or enhanced production of A beta are therefore preferred. For example, transfection of cells with the Swedish Mutant form of APP (APP-SW); with APP-KK; or with APP-SW-KK provides cells having enhanced beta-secretase activity and producing amounts of A beta that can be readily measured.

In such assays, for example, the cells expressing APP and beta-secretase are incubated in a culture medium under conditions suitable for beta-secretase enzymatic activity at its cleavage site on the APP substrate. On exposure of the cells to the compound inhibitor, the amount of A beta released into the medium and/or the amount of CTF99 fragments of APP in the cell lysates is reduced as compared with

the control. The cleavage products of APP can be analyzed, for example, by immune reactions with specific antibodies, as discussed above.

Preferred cells for analysis of beta-secretase activity include primary human neuronal cells, primary transgenic animal neuronal cells where the transgene is APP, and other cells such as those of a stable 293 cell line expressing APP, for example, APP-SW.

***In vivo* Assays: Animal Models**

Various animal models can be used to analyze beta-secretase activity and /or processing of APP to release A beta, as described above. For example, transgenic animals expressing APP substrate and beta-secretase enzyme can be used to demonstrate inhibitory activity of the compounds employed in the methods of the invention. Certain transgenic animal models have been described, for example, in U.S. Patent Nos.: 5,877,399; 5,612,486; 5,387,742; 5,720,936; 5,850,003; 5,877,015,, and 5,811,633, and in Ganes et al., 1995, *Nature* 373:523. Preferred are animals that exhibit characteristics associated with the pathophysiology of AD. Administration of the compound inhibitors of the invention to the transgenic mice described herein provides an alternative method for demonstrating the inhibitory activity of the compounds. Administration of the compounds in a pharmaceutically effective carrier and via an administrative route that reaches the target tissue in an appropriate therapeutic amount is also preferred.

Inhibition of beta-secretase mediated cleavage of APP at the beta-secretase cleavage site and of A beta release can be analyzed in these animals by measure of cleavage fragments in the animal's body fluids such as cerebral fluid or tissues. Analysis of brain tissues for A beta deposits or plaques is preferred.

On contacting an APP substrate with a beta-secretase enzyme in the presence of an inhibitory compound employed in the method of the invention and under conditions sufficient to permit enzymatic mediated cleavage of APP and/or release of A beta from the substrate, the compounds employed in the methods of the invention are effective to reduce beta-secretase-mediated cleavage of APP at the beta-secretase cleavage site and/or effective to reduce released amounts of A beta. Where such contacting is the administration of the inhibitory compounds employed in the methods of the invention to an animal model, for example, as described above, the compounds are effective to reduce A beta deposition in brain tissues of the

animal, and to reduce the number and/or size of beta amyloid plaques. Where such administration is to a human subject, the compounds are effective to inhibit or slow the progression of disease characterized by enhanced amounts of A beta, to slow the progression of AD in the, and/or to prevent onset or development of AD in a patient at risk for the disease.

Unless defined otherwise, all scientific and technical terms used herein have the same meaning as commonly understood by one of skill in the art to which this invention belongs. All patents and publications referred to herein are hereby incorporated by reference for all purposes.

10

DEFINITIONS AND CONVENTIONS

The definitions and explanations below are for the terms as used throughout this entire document including both the specification and the claims.

15

I. CONVENTIONS FOR FORMULAS AND DEFINITIONS OF VARIABLES

The chemical formulas representing various compounds or molecular fragments in the specification and claims may contain variable substituents in addition to expressly defined structural features. These variable substituents are identified by a letter or a letter followed by a numerical subscript, for example, "Z₁" or "R_i" where "i" is an integer. These variable substituents are either monovalent or bivalent, that is, they represent a group attached to the formula by one or two chemical bonds. For example, a group Z₁ would represent a bivalent variable if attached to the formula CH₃-C(=Z₁)H. Groups R_i and R_j would represent monovalent variable substituents if attached to the formula CH₃-CH₂-C(R_i)(R_j)H₂. When chemical formulas are drawn in a linear fashion, such as those above, variable substituents contained in parentheses are bonded to the atom immediately to the left of the variable substituent enclosed in parenthesis. When two or more consecutive variable substituents are enclosed in parentheses, each of the consecutive variable substituents is bonded to the immediately preceding atom to the left which is not enclosed in parentheses. Thus, in the formula above, both R_i and R_j are bonded to the preceding carbon atom. Also, for any molecule with an established system of carbon atom numbering, such as steroids, these carbon atoms are designated as C_i,

where "i" is the integer corresponding to the carbon atom number. For example, C₆ represents the 6 position or carbon atom number in the steroid nucleus as traditionally designated by those skilled in the art of steroid chemistry. Likewise the term "R₆" represents a variable substituent (either monovalent or bivalent) at the C₆ position.

Chemical formulas or portions thereof drawn in a linear fashion represent atoms in a linear chain. The symbol "-" in general represents a bond between two atoms in the chain. Thus CH₃-O-CH₂-CH(R_i)-CH₃ represents a 2-substituted-1-methoxypropane compound. In a similar fashion, the symbol "=" represents a double bond, e.g., CH₂=C(R_i)-O-CH₃, and the symbol "≡" represents a triple bond, e.g., HC≡C-CH(R_i)-CH₂-CH₃. Carbonyl groups are represented in either one of two ways: -CO- or -C(=O)-, with the former being preferred for simplicity.

Chemical formulas of cyclic (ring) compounds or molecular fragments can be represented in a linear fashion. Thus, the compound 4-chloro-2-methylpyridine can be represented in linear fashion by N^{*}=C(CH₃)-CH=CCl-CH=C^{*}H with the convention that the atoms marked with an asterisk (*) are bonded to each other resulting in the formation of a ring. Likewise, the cyclic molecular fragment, 4-(ethyl)-1-piperazinyll can be represented by -N^{*}-(CH₂)₂-N(C₂H₅)-CH₂-C^{*}H₂.

A rigid cyclic (ring) structure for any compounds herein defines an orientation with respect to the plane of the ring for substituents attached to each carbon atom of the rigid cyclic compound. For saturated compounds which have two substituents attached to a carbon atom which is part of a cyclic system, -C(X₁)(X₂)- the two substituents may be in either an axial or equatorial position relative to the ring and may change between axial/equatorial. However, the position of the two substituents relative to the ring and each other remains fixed. While either substituent at times may lie in the plane of the ring (equatorial) rather than above or below the plane (axial), one substituent is always above the other. In chemical structural formulas depicting such compounds, a substituent (X₁) which is "below" another substituent (X₂) will be identified as being in the alpha configuration and is identified by a broken, dashed or dotted line attachment to the carbon atom, i.e., by the symbol "- - -" or "...". The corresponding substituent attached "above" (X₂) the other (X₁) is identified as being in the beta configuration and is indicated by an unbroken line attachment to the carbon atom.

When a variable substituent is bivalent, the valences may be taken together or separately or both in the definition of the variable. For example, a variable R_i attached to a carbon atom as $-C(=R_i)-$ might be bivalent and be defined as oxo or keto (thus forming a carbonyl group $(-CO-)$ or as two separately attached

5 monovalent variable substituents $\alpha-R_{i-j}$ and $\beta-R_{i-k}$. When a bivalent variable, R_i , is defined to consist of two monovalent variable substituents, the convention used to define the bivalent variable is of the form " $\alpha-R_{i-j}:\beta-R_{i-k}$ " or some variant thereof. In such a case both $\alpha-R_{i-j}$ and $\beta-R_{i-k}$ are attached to the carbon atom to give $-C(\alpha-R_{i-j})(\beta-R_{i-k})-$. For example, when the bivalent

10 variable R_6 , $-C(=R_6)-$ is defined to consist of two monovalent variable substituents, the two monovalent variable substituents are $\alpha-R_{6-1}:\beta-R_{6-2}$, ..., $\alpha-R_{6-9}:\beta-R_{6-10}$, etc, giving $-C(\alpha-R_{6-1})(\beta-R_{6-2})-$, ..., $-C(\alpha-R_{6-9})(\beta-R_{6-10})-$, etc. Likewise, for the bivalent variable R_{11} , $-C(=R_{11})-$, two monovalent variable substituents are $\alpha-R_{11-1}:\beta-R_{11-2}$. For a ring substituent for which separate alpha and

15 beta orientations do not exist (e.g. due to the presence of a carbon carbon double bond in the ring), and for a substituent bonded to a carbon atom which is not part of a ring the above convention is still used, but the alpha and beta designations are omitted.

Just as a bivalent variable may be defined as two separate monovalent

20 variable substituents, two separate monovalent variable substituents may be defined to be taken together to form a bivalent variable. For example, in the formula $-C_1(R_i)H-C_2(R_j)H-$ (C_1 and C_2 define arbitrarily a first and second carbon atom, respectively) R_i and R_j may be defined to be taken together to form (1) a second bond between C_1 and C_2 or (2) a bivalent group such as oxa $(-O-)$ and the formula

25 thereby describes an epoxide. When R_i and R_j are taken together to form a more complex entity, such as the group $-X-Y-$, then the orientation of the entity is such that C_1 in the above formula is bonded to X and C_2 is bonded to Y . Thus, by convention the designation "... R_i and R_j are taken together to form $-CH_2-CH_2-O-CO-$..." means a lactone in which the carbonyl is bonded to C_2 . However, when

30 designated "... R_j and R_i are taken together to form $-CO-O-CH_2-CH_2-$ the convention means a lactone in which the carbonyl is bonded to C_1 .

The carbon atom content of variable substituents is indicated in one of two ways. The first method uses a prefix to the entire name of the variable such as " C_1-C_4 ", where both "1" and "4" are integers representing the minimum and maximum

number of carbon atoms in the variable. The prefix is separated from the variable by a space. For example, "C₁-C₄ alkyl" represents alkyl of 1 through 4 carbon atoms, (including isomeric forms thereof unless an express indication to the contrary is given). Whenever this single prefix is given, the prefix indicates the entire carbon atom content of the variable being defined. Thus C₂-C₄ alkoxy-carbonyl describes a group CH₃-(CH₂)_n-O-CO- where n is zero, one or two. By the second method the carbon atom content of only each portion of the definition is indicated separately by enclosing the "C_i-C_j" designation in parentheses and placing it immediately (no intervening space) before the portion of the definition being defined. By this optional convention (C₁-C₃)alkoxy-carbonyl has the same meaning as C₂-C₄ alkoxy-carbonyl because the "C₁-C₃" refers only to the carbon atom content of the alkoxy group. Similarly while both C₂-C₆ alkoxyalkyl and (C₁-C₃)alkoxy(C₁-C₃)alkyl define alkoxyalkyl groups containing from 2 to 6 carbon atoms, the two definitions differ since the former definition allows either the alkoxy or alkyl portion alone to contain 4 or 5 carbon atoms while the latter definition limits either of these groups to 3 carbon atoms.

When the claims contain a fairly complex (cyclic) substituent, at the end of the phrase naming/designating that particular substituent will be a notation in (parentheses) which will correspond to the same name/designation in one of the CHARTS which will also set forth the chemical structural formula of that particular substituent.

II. DEFINITIONS

All temperatures are in degrees Celsius.

25 TLC refers to thin-layer chromatography.

psi refers to pounds/in².

HPLC refers to high pressure liquid chromatography.

THF refers to tetrahydrofuran.

DMF refers to dimethylformamide.

30 EDC refers to ethyl-1-(3-dimethylaminopropyl)carbodiimide or 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride.

HOBt refers to 1-hydroxy benzotriazole hydrate.

NMM refers to N-methylmorpholine.

NBS refers to N-bromosuccinimide.

TEA refers to triethylamine.

BOC refers to 1,1-dimethylethoxy carbonyl or *t*-butoxycarbonyl, -CO-O-C(CH₃)₃.

CBZ refers to benzyloxycarbonyl, -CO-O-CH₂-φ.

5 FMOC refers to 9-fluorenylmethyl carbonate.

TFA refers to trifluoroacetic acid, CF₃-COOH.

CDI refers to 1,1'-carbonyldiimidazole.

Saline refers to an aqueous saturated sodium chloride solution.

Chromatography (column and flash chromatography) refers to
10 purification/separation of compounds expressed as (support, eluent). It is understood that the appropriate fractions are pooled and concentrated to give the desired compound(s).

CMR refers to C-13 magnetic resonance spectroscopy, chemical shifts are reported in ppm (δ) downfield from TMS.

15 NMR refers to nuclear (proton) magnetic resonance spectroscopy, chemical shifts are reported in ppm (δ) downfield from TMS.

IR refers to infrared spectroscopy.

MS refers to mass spectrometry expressed as *m/e*, *m/z* or mass/charge unit.

MH⁺ refers to the positive ion of a parent plus a hydrogen atom. EI refers to
20 electron impact. CI refers to chemical ionization. FAB refers to fast atom bombardment.

HRMS refers to high resolution mass spectrometry.

Ether refers to diethyl ether.

Pharmaceutically acceptable refers to those properties and/or substances
25 which are acceptable to the patient from a pharmacological/toxicological point of view and to the manufacturing pharmaceutical chemist from a physical/chemical point of view regarding composition, formulation, stability, patient acceptance and bioavailability.

When solvent pairs are used, the ratios of solvents used are volume/volume
30 (v/v).

When the solubility of a solid in a solvent is used the ratio of the solid to the solvent is weight/volume (wt/v).

BOP refers to benzotriazol-1-yloxy-tris(dimethylamino)phosphonium hexafluorophosphate.

TBDMSCl refers to *t*-butyldimethylsilyl chloride.

TBDMSOTf refers to *t*-butyldimethylsilyl trifluorosulfonic acid ester.

5 Trisomy 21 refers to Down's Syndrome.

The following terms are used (in EXAMPLEs 321 and above) for the amide forming agent (IX):

"PHTH" refers to $(\text{CH}_3\text{-CH}_2\text{-CH}_2)_2\text{N-CO-phenyl-CO-OH}$ where the attachment to the - phenyl- ring is 1,3-;

10 "5-Me-PHTH" refers to $(\text{CH}_3\text{-CH}_2\text{-CH}_2)_2\text{N-CO-(CH}_3\text{-) phenyl -CO-OH}$ where the attachment to the - phenyl - ring is 1,3- for the carbonyl groups and 5- for the methyl group;

"3,5-pyridinyl" refers to $(\text{CH}_3\text{-CH}_2\text{-CH}_2)_2\text{N-CO-(pyridinyl)-CO-OH}$ where the attachment to the -pyridinyl- ring is 3,5- for the carbonyl groups;

15 "-SO₂-" refers to $(\text{CH}_3\text{-CH}_2\text{-CH}_2)_2\text{CH-SO}_2\text{- phenyl -CO-OH}$ where the attachment to the - phenyl - ring is 1,3-;

"5-OMe-PHTH" refers to $(\text{CH}_3\text{-CH}_2\text{-CH}_2)_2\text{N-CO-(CH}_3\text{-O-) phenyl -CO-OH}$ where the attachment to the - phenyl - ring is 1,3- for the carbonyl groups and 5- for the methoxy group;

20 "5-Cl-PHTH" refers to $(\text{CH}_3\text{-CH}_2\text{-CH}_2)_2\text{N-CO-(Cl-)phenyl-CO-OH}$ where the attachment to the -phenyl- ring is 1,3- for the carbonyl groups and 5- for the chlorine atom;

"5-F-PHTH" refers to $(\text{CH}_3\text{-CH}_2\text{-CH}_2)_2\text{N-CO-(F-)phenyl-CO-OH}$ where the attachment to the -phenyl- ring is 1,3- for the carbonyl groups and 5- for the fluorine
25 atom;

"thienyl" refers to $(\text{CH}_3\text{-CH}_2\text{-CH}_2)_2\text{N-CO-thienyl-CO-OH}$ where the attachment to the thiophene ring is -2,5;

"2,4-pyridinyl" refers to $(\text{CH}_3\text{-CH}_2\text{-CH}_2)_2\text{N-CO-(pyridinyl)-CO-OH}$ where the attachment to the -pyridinyl- ring is 2,4- for the carbonyl groups;

30 "4,6-pyrimidinyl" refers to $(\text{CH}_3\text{-CH}_2\text{-CH}_2)_2\text{N-CO-(pyrimidinyl-)phenyl-CO-OH}$ where the attachment to the -pyrimidinyl- ring is 4,6- for the carbonyl groups;

"morpholinyl" refers to morpholinyl-CO-phenyl-CO-OH where the attachment to the -phenyl- ring is 1,3 for the carbonyl groups.

APP, amyloid precursor protein, is defined as any APP polypeptide, including APP variants, mutations, and isoforms, for example, as disclosed in U.S. Patent No. 5,766,846.

5 A beta, amyloid beta peptide, is defined as any peptide resulting from beta-secretase mediated cleavage of APP, including peptides of 39, 40, 41, 42, and 43 amino acids, and extending from the beta-secretase cleavage site to amino acids 39, 40, 41, 42, or 43.

Beta-secretase (BACE1, Asp2, Memapsin 2) is an aspartyl protease that mediates cleavage of APP at the amino-terminal edge of A beta. Human beta-secretase is described, for example, in WO00/17369.

A therapeutically effective amount is defined as an amount effective to reduce or lessen at least one symptom of the disease being treated or to reduce or delay onset of one or more clinical markers or symptoms of the disease.

The present invention provides compounds, compositions, and methods for
15 inhibiting beta-secretase enzyme activity and A beta peptide production. Inhibition of beta-secretase enzyme activity halts or reduces the production of A beta from APP and reduces or eliminates the formation of beta-amyloid deposits in the brain.


CHEMICAL EXAMPLES

20


Exemplary Compounds of the Invention

Examples of compounds that are within the invention include but are not limited to those depicted below.

25 Example 1, N-[1-(3,5-Difluoro-benzyl)-2-hydroxy-3-(N'-methyl-N'-phenyl-hydrazino)-propyl]-5-methyl-N', N'-dipropyl-isophthalamide



Example 2, N-{1-(3,5-Difluoro-benzyl)-2-hydroxy-3-[N'-methyl-N'-(4-methyl-pentanoyl)-hydrazino]-propyl}-5-methyl-N',N'-dipropyl-isophthalamide



Example 3, N-[1-(3,5-Difluoro-benzyl)-2-hydroxy-3-phenoxyamino-propyl]-
5 5-methyl-N',N'-dipropyl-isophthalamide




CHART A





CHART A' (continued)




110
CHART B

CHART C








CHART C' (continued)

I





XI

HCl




XII



XV

CHART D

CHART D



BIOLOGICAL EXAMPLES**Example A****Enzyme Inhibition Assay**

5 The compounds of the invention are analyzed for inhibitory activity by use of the MBP-C125 assay. This assay determines the relative inhibition of beta-secretase cleavage of a model APP substrate, MBP-C125SW, by the compounds assayed as compared with an untreated control. A detailed description of the assay parameters can be found, for example, in U.S. Patent No. 5,942,400. Briefly, the
10 substrate is a fusion peptide formed of maltose binding protein (MBP) and the carboxy terminal 125 amino acids of APP-SW, the Swedish mutation. The beta-secretase enzyme is derived from human brain tissue as described in Sinha et.al, 1999, *Nature* 40:537-540) or recombinantly produced as the full-length enzyme (amino acids 1-501), and can be prepared, for example, from 293 cells expressing
15 the recombinant cDNA, as described in WO00/47618.

 Inhibition of the enzyme is analyzed, for example, by immunoassay of the enzyme's cleavage products. One exemplary ELISA uses an anti-MBP capture antibody that is deposited on precoated and blocked 96-well high binding plates, followed by incubation with diluted enzyme reaction supernatant, incubation with a
20 specific reporter antibody, for example, biotinylated anti-SW192 reporter antibody, and further incubation with streptavidin/alkaline phosphatase. In the assay, cleavage of the intact MBP-C125SW fusion protein results in the generation of a truncated amino-terminal fragment, exposing a new SW-192 antibody-positive epitope at the carboxy terminus. Detection is effected by a fluorescent substrate
25 signal on cleavage by the phosphatase. ELISA only detects cleavage following Leu 596 at the substrate's APP-SW 751 mutation site.

Specific Assay Procedure:

 Compounds are diluted in a 1:1 dilution series to a six-point concentration
30 curve (two wells per concentration) in one 96-plate row per compound tested. Each of the test compounds is prepared in DMSO to make up a 10 millimolar stock solution. The stock solution is serially diluted in DMSO to obtain a final compound concentration of 200 micromolar at the high point of a 6-point dilution curve. Ten
(10) microliters of each dilution is added to each of two wells on row C of a

corresponding V-bottom plate to which 190 microliters of 52 millimolar NaOAc, 7.9% DMSO, pH 4.5 are pre-added. The NaOAc diluted compound plate is spun down to pellet precipitant and 20 microliters/well is transferred to a corresponding flat-bottom plate to which 30 microliters of ice-cold enzyme-substrate mixture (2.5 microliters MBP-C125SW substrate, 0.03 microliters enzyme and 24.5 microliters ice cold 0.09% TX100 per 30 microliters) is added. The final reaction mixture of 200 micromolar compound at the highest curve point is in 5% DMSO, 20 millimolar NaAc, 0.06% TX100, at pH 4.5.

Warming the plates to 37 degrees C starts the enzyme reaction. After 90 minutes at 37 degrees C, 200 microliters/well cold specimen diluent is added to stop the reaction and 20 microliters/well is transferred to a corresponding anti-MBP antibody coated ELISA plate for capture, containing 80 microliters/well specimen diluent. This reaction is incubated overnight at 4 degrees C and the ELISA is developed the next day after a 2 hour incubation with anti-192SW antibody, followed by Streptavidin-AP conjugate and fluorescent substrate. The signal is read on a fluorescent plate reader.

Relative compound inhibition potency is determined by calculating the concentration of compound that showed a fifty percent reduction in detected signal (IC_{50}) compared to the enzyme reaction signal in the control wells with no added compound. In this assay, the compounds of the invention exhibited an IC_{50} of less than 50 micromolar.

Example B

Cell Free Inhibition Assay utilizing a Synthetic APP Substrate

A synthetic APP substrate that can be cleaved by beta-secretase and having N-terminal biotin and made fluorescent by the covalent attachment of oregon green at the Cys residue is used to assay beta-secretase activity in the presence or absence of the inhibitory compounds of the invention. Useful substrates include the following:

30

Biotin-SEVNL-DAEFR[oregon green]KK	[SEQ ID NO: 1]
Biotin-SEVKM-DAEFR[oregon green]KK	[SEQ ID NO: 2]
Biotin-GLNIKTEEISEISY-EVEFRC[oregon green]KK	[SEQ ID NO: 3]

Biotin-ADRGLTTRPGSGLTNIKTEEISEVNL-DAEF[oregon green]KK [SEQ ID NO:4]

Biotin-FVNQHLCoxGSHLVEALY-LVCoxGERGFFYTPKA[oregon green]KK [SEQ ID NO: 5]

5 The enzyme (0.1 nanomolar) and test compounds (0.001 - 100 micromolar) are incubated in pre-blocked, low affinity, black plates (384 well) at 37 degrees C for 30 minutes. The reaction is initiated by addition of 150 millimolar substrate to a final volume of 30 microliter per well. The final assay conditions are: 0.001 - 100 micromolar compound inhibitor; 0.1 molar sodium acetate (pH 4.5); 150 nanomolar
10 substrate; 0.1 nanomolar soluble beta-secretase; 0.001% Tween 20, and 2% DMSO. The assay mixture is incubated for 3 hours at 37 °C, and the reaction is terminated by the addition of a saturating concentration of immunopure streptavidin. After incubation with streptavidin at room temperature for 15 minutes, fluorescence polarization is measured, for example, using a LJL Acquest (Ex485 nm/ Em530
15 nm). The activity of the beta-secretase enzyme is detected by changes in the fluorescence polarization that occur when the substrate is cleaved by the enzyme. Incubation in the presence or absence of compound inhibitor demonstrates specific inhibition of beta-secretase enzymatic cleavage of its synthetic APP substrate. In this assay, compounds of the invention exhibited an IC₅₀ of less than 50 micromolar.

20

Example C

Beta-secretase inhibition: P26-P4'SW assay

Synthetic substrates containing the beta-secretase cleavage site of APP are used to
25 assay beta-secretase activity, using the methods described, for example, in published PCT application WO00/47618. The P26-P4'SW substrate is a peptide of the sequence: (biotin)CGGADRGLTTRPGSGLTNIKTEEISEVNLDAEF [SEQ ID NO: 6]

The P26-P1 standard has the sequence:

30 (biotin)CGGADRGLTTRPGSGLTNIKTEEISEVNL [SEQ ID NO: 7]

Briefly, the biotin-coupled synthetic substrates are incubated at a concentration of from about 0 to about 200 micromolar in this assay. When testing inhibitory compounds, a substrate concentration of about 1.0 micromolar is preferred. Test compounds diluted in DMSO are added to the reaction mixture, with

a final DMSO concentration of 5%. Controls also contain a final DMSO concentration of 5%. The concentration of beta secretase enzyme in the reaction is varied, to give product concentrations with the linear range of the ELISA assay, about 125 to 2000 picomolar, after dilution.

5 The reaction mixture also includes 20 millimolar sodium acetate, pH 4.5, 0.06% Triton X100, and is incubated at 37 degrees C for about 1 to 3 hours. Samples are then diluted in assay buffer (for example, 145.4 nanomolar sodium chloride, 9.51 millimolar sodium phosphate, 7.7 millimolar sodium azide, 0.05% Triton X405, 6g/liter bovine serum albumin, pH 7.4) to quench the reaction, then
10 diluted further for immunoassay of the cleavage products.

Cleavage products can be assayed by ELISA. Diluted samples and standards are incubated in assay plates coated with capture antibody, for example, SW192, for about 24 hours at 4 degrees C. After washing in TTBS buffer (150 millimolar sodium chloride, 25 millimolar Tris, 0.05% Tween 20, pH 7.5), the samples are
15 incubated with strepavidin-AP according to the manufacturer's instructions. After a one hour incubation at room temperature, the samples are washed in TTBS and incubated with fluorescent substrate solution A (31.2 g/liter 2-amino-2-methyl-1-propanol, 30 mg/liter, pH 9.5). Reaction with streptavidin-alkaline phosphate permits detection by fluorescence. Compounds that are effective inhibitors of beta-
20 secretase activity demonstrate reduced cleavage of the substrate as compared to a control.

Example D

Assays using Synthetic Oligopeptide-Substrates

25 Synthetic oligopeptides are prepared that incorporate the known cleavage site of beta-secretase, and optionally detectable tags, such as fluorescent or chromogenic moieties. Examples of such peptides, as well as their production and detection methods are described in U.S. Patent No: 5,942,400, herein
30 incorporated by reference. Cleavage products can be detected using high performance liquid chromatography, or fluorescent or chromogenic detection methods appropriate to the peptide to be detected, according to methods well known in the art.

By way of example, one such peptide has the sequence SEVNL-DAEF [SEQ ID NO: 8], and the cleavage site is between residues 5 and 6. Another preferred substrate has the sequence ADRGLTTRPGSGLTNIKTEEISEVNL-DAEF [SEQ ID NO: 9], and the cleavage site is between residues 26 and 27.

5 These synthetic APP substrates are incubated in the presence of beta-secretase under conditions sufficient to result in beta-secretase mediated cleavage of the substrate. Comparison of the cleavage results in the presence of the compound inhibitor to control results provides a measure of the compound's inhibitory activity.

10 **Example E**

Inhibition of beta-secretase activity - cellular assay

 An exemplary assay for the analysis of inhibition of beta-secretase activity utilizes the human embryonic kidney cell line HEKp293 (ATCC Accession No. 15 CRL-1573) transfected with APP751 containing the naturally occurring double mutation Lys651Met52 to Asn651Leu652 (numbered for APP751), commonly called the Swedish mutation and shown to overproduce A beta (Citron et.al., 1992, *Nature* 360:672-674), as described in USPN 5,604,102.

 The cells are incubated in the presence/absence of the inhibitory compound 20 (diluted in DMSO) at the desired concentration, generally up to 10 micrograms/ml. At the end of the treatment period, conditioned media is analyzed for beta-secretase activity, for example, by analysis of cleavage fragments. A beta can be analyzed by immunoassay, using specific detection antibodies. The enzymatic activity is measured in the presence and absence of the compound inhibitors to demonstrate 25 specific inhibition of beta-secretase mediated cleavage of APP substrate.

Example F

Inhibition of Beta-Secretase in Animal Models of AD

30 Various animal models can be used to screen for inhibition of beta-secretase activity. Examples of animal models useful in the invention include, but are not limited to, mouse, guinea pig, dog, and the like. The animals used can be wild type, transgenic, or knockout models. In addition, mammalian models can express mutations in APP, such as APP695-SW and the like described herein. Examples of

transgenic non-human mammalian models are described in U.S. Patent Nos. 5,604,102, 5,912,410 and 5,811,633.

PDAPP mice, prepared as described in Games et.al., 1995, *Nature* 373:523-527 are useful to analyze *in vivo* suppression of A beta release in the presence of putative inhibitory compounds. As described in USPN 6,191,166, 4 month old PDAPP mice are administered compound formulated in vehicle, such as corn oil. The mice are dosed with compound (1-30 mg/ml; preferably 1-10 mg/ml). After time, e.g., 3-10 hours, the animals are sacrificed, and brains removed for analysis.

Transgenic animals are administered an amount of the compound inhibitor formulated in a carrier suitable for the chosen mode of administration. Control animals are untreated, treated with vehicle, or treated with an inactive compound. Administration can be acute, i.e., single dose or multiple doses in one day, or can be chouronic, i.e., dosing is repeated daily for a period of days. Beginning at time 0, brain tissue or cerebral fluid is obtained from selected animals and analyzed for the presence of APP cleavage peptides, including A beta, for example, by immunoassay using specific antibodies for A beta detection. At the end of the test period, animals are sacrificed and brain tissue or cerebral fluid is analyzed for the presence of A beta and/or beta-amyloid plaques. The tissue is also analyzed for necrosis.

Animals administered the compound inhibitors of the invention are expected to demonstrate reduced A beta in brain tissues or cerebral fluids and reduced beta amyloid plaques in brain tissue, as compared with non-treated controls.

Example G

Inhibition of A beta production in human patients

Patients suffering from Alzheimer's Disease (AD) demonstrate an increased amount of A beta in the brain. AD patients are administered an amount of the compound inhibitor formulated in a carrier suitable for the chosen mode of administration. Administration is repeated daily for the duration of the test period. Beginning on day 0, cognitive and memory tests are performed, for example, once per month.

Patients administered the compound inhibitors are expected to demonstrate slowing or stabilization of disease progression as analyzed by changes in one or more of the following disease parameters: A beta present in CSF or plasma; brain or

hippocampal volume; A beta deposits in the brain; amyloid plaque in the brain; and scores for cognitive and memory function, as compared with control, non-treated patients.

5 **Example H**

Prevention of A beta production in patients at risk for AD

Patients predisposed or at risk for developing AD are identified either by recognition of a familial inheritance pattern, for example, presence of the Swedish
10 Mutation, and/or by monitoring diagnostic parameters. Patients identified as predisposed or at risk for developing AD are administered an amount of the compound inhibitor formulated in a carrier suitable for the chosen mode of administration. Administration is repeated daily for the duration of the test period. Beginning on day 0, cognitive and memory tests are performed, for example, once
15 per month.

Patients administered the compound inhibitors are expected to demonstrate slowing or stabilization of disease progression as analyzed by changes in one or more of the following disease parameters: A beta present in CSF or plasma; brain or hippocampal volume; amyloid plaque in the brain; and scores for cognitive and
20 memory function, as compared with control, non-treated patients.

It should be noted that, as used in this specification and the appended claims, the singular forms "a," "an," and "the" include plural referents unless the content clearly dictates otherwise. Thus, for example, reference to a composition containing "a compound" includes a mixture of two or more compounds. It should also be
25 noted that the term "or" is generally employed in its sense including "and/or" unless the content clearly dictates otherwise.

Unless defined otherwise, all scientific and technical terms used herein have the same meaning as commonly understood by one of skill in the art to which this invention belongs.


30 All patents and publications referred to herein are hereby incorporated by reference for all purposes.

The invention has been described with reference to various specific and preferred embodiments and techniques. However, it should be understood that many

variations and modifications may be made while remaining within the spirit and scope of the invention.

WE CLAIM:

1. A substituted amine of formula (XV)



where R_1 is:

- (I) C_1 - C_6 alkyl, optionally substituted with one, two or three substituents selected from the group consisting of C_1 - C_3 alkyl, C_1 - C_7 alkyl (optionally substituted with C_1 - C_3 alkyl and C_1 - C_3 alkoxy), -F, -Cl, -Br, -I, -OH, -SH, -C \equiv N, -CF $_3$, C_1 - C_3 alkoxy, -NR $_{1-a}$ R $_{1-b}$ where R $_{1-a}$ and R $_{1-b}$ are -H or C_1 - C_6 alkyl, and -OC=O NR $_{1-a}$ R $_{1-b}$ where R $_{1-a}$ and R $_{1-b}$ are as defined above,
- (II) -CH $_2$ -S(O) $_{0-2}$ -(C_1 - C_6 alkyl),
- (III) -CH $_2$ -CH $_2$ -S(O) $_{0-2}$ -(C_1 - C_6 alkyl),
- (IV) C_2 - C_6 alkenyl with one or two double bonds, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, -C \equiv N, -CF $_3$, C_1 - C_3 alkoxy, and -NR $_{1-a}$ R $_{1-b}$ where R $_{1-a}$ and R $_{1-b}$ are -H or C_1 - C_6 alkyl,
- (V) C_2 - C_6 alkynyl with one or two triple bonds, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, -C \equiv N, -CF $_3$, C_1 - C_3 alkoxy, and -NR $_{1-a}$ R $_{1-b}$ where R $_{1-a}$ and R $_{1-b}$ are -H or C_1 - C_6 alkyl,
- (VI) -(CH $_2$) $_{n_1}$ -(R $_{1-aryl}$) where n_1 is zero or one and where R $_{1-aryl}$ is phenyl, 1-naphthyl, 2-naphthyl and indanyl, indenyl, dihydronaphthalyl, or tetralinyl optionally substituted with one, two, three or four of the following substituents on the aryl ring:
- (A) C_1 - C_6 alkyl optionally substituted with one, two or three substituents selected from the group consisting of C_1 - C_3 alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C \equiv N, -CF $_3$, and C_1 - C_3 alkoxy, and -NR $_{1-a}$ R $_{1-b}$ where R $_{1-a}$ and R $_{1-b}$ are as defined above,

(B) C₂-C₆ alkenyl with one or two double bonds, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, -C≡N, -CF₃, C₁-C₃ alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are -H or C₁-C₆ alkyl,

5 (C) C₂-C₆ alkynyl with one or two triple bonds, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, -C≡N, -CF₃, C₁-C₃ alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are -H or C₁-C₆ alkyl,

(D) -F, Cl, -Br or -I,

10 (F) -C₁-C₆ alkoxy optionally substituted with one, two or three of -F,

(G) -NR_{N-2}R_{N-3} where R_{N-2} and R_{N-3} are as defined below,

(H) -OH,

(I) -C≡N,

15 (J) C₃-C₇ cycloalkyl, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, -C≡N, -CF₃, C₁-C₃ alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are -H or C₁-C₆ alkyl,

(K) -CO-(C₁-C₄ alkyl),

(L) -SO₂-NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

20 (M) -CO-NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,
or

(N) -SO₂-(C₁-C₄ alkyl),

(VII) -(CH₂)_{n1}-(R_{1-heteroaryl}) where n₁ is as defined above and where R_{1-heteroaryl} is selected from the group consisting of:

25 pyridinyl,
pyrimidinyl,
quinolinyl,
benzothienyl,
indolyl,
30 indolinyl,
pyridazinyl,
pyrazinyl,
isoquinolyl,

	quinazolinyl,
	quinoxalinyl,
	phthalazinyl,
	imidazolyl,
5	isoxazolyl,
	pyrazolyl,
	oxazolyl,
	thiazolyl,
	indolizinyll,
10	indazolyl,
	benzothiazolyl,
	benzimidazolyl,
	benzofuranyl,
	furanyl,
15	thienyl,
	pyrrolyl,
	oxadiazolyl,
	thiadiazolyl,
	triazolyl,
20	tetrazolyl,
	oxazolopyridinyl,
	imidazopyridinyl,
	isothiazolyl,
	naphthyridinyl,
25	cinnolinyl,
	carbazolyl,
	beta-carbolinyl,
	isochromanyl,
	chromanyl,
30	tetrahydroisoquinolinyl,
	isoindolinyl,
	isobenzotetrahydrofuranyl,
	isobenzotetrahydrothienyl,
	isobenzothieryl,

	benzoxazolyl,
	pyridopyridinyl,
	benzotetrahydrofuranyl,
	benzotetrahydrothienyl,
5	purinyl,
	benzodioxolyl,
	triazinyl,
	phenoxazinyl,
	phenothiazinyl,
10	pteridinyl,
	benzothiazolyl,
	imidazopyridinyl,
	imidazothiazolyl,
	dihydrobenzisoaxazinyl,
15	benzisoaxazinyl,
	benzoxazinyl,
	dihydrobenzisothiazinyl,
	benzopyranyl,
	benzothiopyranyl,
20	coumarinyl,
	isocoumarinyl,
	chromonyl,
	chromanonyl,
	pyridinyl-N-oxide,
25	tetrahydroquinolinyl
	dihydroquinolinyl
	dihydroquinolinonyl
	dihydroisoquinolinonyl
	dihydrocoumarinyl
30	dihydroisocoumarinyl
	isoindolinonyl
	benzodioxanyl
	benzoxazolinonyl
	pyrrolyl N-oxide,

- 5 pyrimidinyl N-oxide,
 pyridazinyl N-oxide,
 pyrazinyl N-oxide,
 quinolinyl N-oxide,
 indolyl N-oxide,
 indolinyl N-oxide,
 isoquinolyl N-oxide,
 quinazolinyl N-oxide,
 quinoxalinyl N-oxide,
 10 phthalazinyl N-oxide,
 imidazolyl N-oxide,
 isoxazolyl N-oxide,
 oxazolyl N-oxide,
 thiazolyl N-oxide,
 15 indoliziny N-oxide,
 indazolyl N-oxide,
 benzothiazolyl N-oxide,
 benzimidazolyl N-oxide,
 pyrrolyl N-oxide,
 20 oxadiazolyl N-oxide,
 thiadiazolyl N-oxide,
 triazolyl N-oxide,
 tetrazolyl N-oxide,
 benzothiopyranyl S-oxide, and
 25 benzothiopyranyl S,S-dioxide,

where the $R_{1\text{-heteroaryl}}$ group is bonded to $-(CH_2)_{n1}-$ by any ring atom
 of the parent $R_{1\text{-heteroaryl}}$ group substituted by hydrogen such that the new bond to the
 $R_{1\text{-heteroaryl}}$ group replaces the hydrogen atom and its bond, where heteroaryl is
 optionally substituted with one, two, three or four of:

- 30 (1) C_1-C_6 alkyl optionally substituted with one, two or
 three substituents selected from the group consisting of C_1-C_3 alkyl, -F, -Cl, -Br, -I,
 -OH, -SH, $-C\equiv N$, $-CF_3$, C_1-C_3 alkoxy, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as
 defined above,

- (2) C_2-C_6 alkenyl with one or two double bonds,
optionally substituted with one, two or three substituents selected from the group
consisting of -F, -Cl, -OH, -SH, $-C\equiv N$, $-CF_3$, C_1-C_3 alkoxy, and $-NR_{1-a}R_{1-b}$ where R_{1-a}
and R_{1-b} are -H or C_1-C_6 alkyl,
- 5 (3) C_2-C_6 alkynyl with one or two triple bonds,
optionally substituted with one, two or three substituents selected from the group
consisting of -F, -Cl, -OH, -SH, $-C\equiv N$, $-CF_3$, C_1-C_3 alkoxy, and $-NR_{1-a}R_{1-b}$ where R_{1-a}
and R_{1-b} are -H or C_1-C_6 alkyl,
- (4) -F, Cl, -Br or -I,
- 10 (6) $-C_1-C_6$ alkoxy optionally substituted with one, two,
or three of -F,
- (7) $-NR_{N-2}R_{N-3}$ where R_{N-2} and R_{N-3} are as defined
below,
- (8) -OH,
- 15 (9) $-C\equiv N$,
- (10) C_3-C_7 cycloalkyl, optionally substituted with one,
two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, -
 $C\equiv N$, $-CF_3$, C_1-C_3 alkoxy, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are -H or C_1-C_6 alkyl,
- (11) $-CO-(C_1-C_4 \text{ alkyl})$,
- 20 (12) $-SO_2-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined
above,
- (13) $-CO-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined
above, or
- (14) $-SO_2-(C_1-C_4 \text{ alkyl})$, with the proviso that when n_1
25 is zero $R_{1-heteroaryl}$ is not bonded to the carbon chain by nitrogen; or
(VIII) $-(CH_2)_{n_1}-(R_{1-heterocycle})$ where n_1 is as defined above and $R_{1-heterocycle}$
is selected from the group consisting of:
- morpholinyl,
- thiomorpholinyl,
- 30 thiomorpholinyl S-oxide,
- thiomorpholinyl S,S-dioxide,
- piperazinyl,
- homopiperazinyl,

pyrrolidinyl,
 pyrrolinyl,
 tetrahydropyranyl,
 piperidinyl,
 5 tetrahydrofuranyl,
 tetrahydrothienyl,
 homopiperidinyl,
 homomorpholinyl,
 homothiomorpholinyl,
 10 homothiomorpholinyl S,S-dioxide,
 oxazolidinonyl,
 dihydropyrazolyl,
 dihydropyrrolyl,
 dihydropyrazinyl,
 15 dihydropyridinyl,
 dihydropyrimidinyl,
 dihydrofuryl,
 dihydropyranyl,
 tetrahydrothienyl S-oxide,
 20 tetrahydrothienyl S,S-dioxide, and
 homothiomorpholinyl S-oxide,

where the $R_{1\text{-heterocycle}}$ group is bonded by any atom of the parent $R_{1\text{-heterocycle}}$ group substituted by hydrogen such that the new bond to the $R_{1\text{-heterocycle}}$ group replaces the hydrogen atom and its bond, where heterocycle is optionally substituted with one, two, three or four:

(1) $C_1\text{-}C_6$ alkyl optionally substituted with one, two or three substituents selected from the group consisting of $C_1\text{-}C_3$ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C \equiv N, -CF₃, $C_1\text{-}C_3$ alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(2) $C_2\text{-}C_6$ alkenyl with one or two double bonds, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, -C \equiv N, -CF₃, $C_1\text{-}C_3$ alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are -H or $C_1\text{-}C_6$ alkyl,

- (3) C₂-C₆ alkynyl with one or two triple bonds,
optionally substituted with one, two or three substituents selected from the group
consisting of -F, -Cl, -OH, -SH, -C≡N, -CF₃, C₁-C₃ alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a}
and R_{1-b} are -H or C₁-C₆ alkyl,
- 5 (4) -F, Cl, -Br or -I,
(5) C₁-C₆ alkoxy,
(6) -C₁-C₆ alkoxy optionally substituted with one,
two, or three -F,
(7) -NR_{N-2}R_{N-3} where R_{N-2} and R_{N-3} are as defined
- 10 below,
(8) -OH,
(9) -C≡N,
(10) C₃-C₇ cycloalkyl, optionally substituted with one,
two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, -
15 C≡N, -CF₃, C₁-C₃ alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are -H or C₁-C₆ alkyl,
(11) -CO-(C₁-C₄ alkyl),
(12) -SO₂-NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined
above,
(13) -CO-NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined
- 20 above,
(14) -SO₂-(C₁-C₄ alkyl), or
(15) =O, with the proviso that when n₁ is zero R₁₋
heterocycle is not bonded to the carbon chain by nitrogen;

25 where R₂ is:

- (I)-H,
(II) C₁-C₆ alkyl, optionally substituted with one, two or three
substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH,
-SH, -C≡N, -CF₃, C₁-C₃ alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined
- 30 above,
(III) -(CH₂)₀₋₄-R₂₋₁ where R₂₋₁ is R_{1-aryl} or R_{1-heteroaryl} where R_{1-aryl} and
R_{1-heteroaryl} are as defined above;

(IV) C₂-C₆ alkenyl with one or two double bonds, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, -C≡N, -CF₃, C₁-C₃ alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are -H or C₁-C₆ alkyl,

5 (V) C₂-C₆ alkynyl with one or two triple bonds, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, -C≡N, -CF₃, C₁-C₃ alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are -H or C₁-C₆ alkyl, or

10 (VI) -(CH₂)₀₋₄- C₃-C₇ cycloalkyl, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, -C≡N, -CF₃, C₁-C₃ alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are -H or C₁-C₆ alkyl;

where R₃ is:

(I)-H,

15 (II) C₁-C₆ alkyl, optionally substituted with one, two or three substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₃ alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

20 (III) -(CH₂)₀₋₄-R₂₋₁ where R₂₋₁ is R_{1-aryl} or R_{1-heteroaryl} where R_{1-aryl} and R_{1-heteroaryl} are as defined above

(IV) C₂-C₆ alkenyl with one or two double bonds,

(V) C₂-C₆ alkynyl with one or two triple bonds; or

25 (VI) -(CH₂)₀₋₄- C₃-C₇ cycloalkyl, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, -C≡N, -CF₃, C₁-C₃ alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are -H or C₁-C₆ alkyl,

and where R₂ and R₃ are taken together with the carbon to which they are attached to form a carbocycle of three, four, five, six, or seven carbon atoms, optionally where one carbon atom is replaced by a heteroatom selected from the group consisting of -O-, -S-, -SO₂-, and -NR_{N-2}-, where R_{N-2} is selected from the group consisting of:

(a) -H,

(b) -C₁-C₆ alkyl optionally substituted with one substituent selected from the group consisting of:

131

(i) -OH, and

(ii) -NH₂,(c) -C₁-C₆ alkyl optionally substituted with

one, two, or three -F, -Cl, -Br, or -I,

5

(d) -C₃-C₇ cycloalkyl,(e) -(C₁-C₂ alkyl)-(C₃-C₇ cycloalkyl),(f) -(C₁-C₆ alkyl)-O-(C₁-C₃ alkyl),(g) -C₂-C₆ alkenyl with one or two double

bonds,

10

(h) -C₂-C₆ alkynyl with one or two triple

bonds,

(i) -C₁-C₆ alkyl chain with one double bond

and one triple bond,

(j) -R_{1-aryl} where R_{1-aryl} is as defined above, and

15

(k) -R_{1-heteroaryl} where R_{1-heteroaryl} is as defined

above;

where R_N is:(I) R_{N-1}-X_N- where X_N is selected from the group consisting of:

20

(A) -CO-,

(B) -SO₂-,(C) -(CR'R'')₁₋₆ where R' and R'' are the same or different and are -H and C₁-C₄ alkyl,(D) -CO-(CR'R'')₁₋₆-X_{N-1} where X_{N-1} is selected from the

25 group consisting of -O-, -S- and -NR'- and where R' and R'' are as defined above, and

(E) a single bond;

where R_{N-1} is selected from the group consisting of:(A) R_{N-aryl} where R_{N-aryl} is phenyl, 1-naphthyl, 2-naphthyl,

30 tetralinyl, indanyl, 6,7,8,9-tetrahydro-5H-benzo[a]cycloheptenyl, or dihydronaphthyl optionally substituted with one, two or three of the following substituents which can be the same or different and are:

(1) C₁-C₆ alkyl, optionally substituted with one, two or three substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I,

-OH, -SH, -C≡N, -CF₃, C₁-C₃ alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(2) -OH,

(3) -NO₂,

5 (4) -F, -Cl, -Br, -I,

(5) -CO-OH,

(6) -C≡N,

(7) -(CH₂)₀₋₄-CO-NR_{N-2}R_{N-3} where R_{N-2} and R_{N-3} are the same or different and are selected from the group consisting of:

10 (a) -H,

(b) -C₁-C₆ alkyl optionally substituted with one substituent selected from the group consisting of:

(i) -OH, and

(ii) -NH₂,

15 (c) -C₁-C₆ alkyl optionally substituted with one, two, or three -F, -Cl, -Br, or -I,

(d) -C₃-C₇ cycloalkyl,

(e) -(C₁-C₂ alkyl)-(C₃-C₇ cycloalkyl),

(f) -(C₁-C₆ alkyl)-O-(C₁-C₃ alkyl),

20 (g) -C₂-C₆ alkenyl with one or two double bonds,

(h) -C₂-C₆ alkynyl with one or two triple

bonds,

(i) -C₁-C₆ alkyl chain with one double bond

25 and one triple bond,

(j) -R_{1-aryl} where R_{1-aryl} is as defined above, and

(k) -R_{1-heteroaryl} where R_{1-heteroaryl} is as defined

above,

(8) -(CH₂)₀₋₄-CO-(C₁-C₁₂ alkyl),

30 (9) -(CH₂)₀₋₄-CO-(C₂-C₁₂ alkenyl with one, two or three double bonds),

(10) -(CH₂)₀₋₄-CO-(C₂-C₁₂ alkynyl with one, two or

three triple bonds),

(11) $-(\text{CH}_2)_{0-4}-\text{CO}-(\text{C}_3-\text{C}_7 \text{ cycloalkyl})$,

(12) $-(\text{CH}_2)_{0-4}-\text{CO}-\text{R}_{1-\text{aryl}}$ where $\text{R}_{1-\text{aryl}}$ is as defined

above,

(13) $-(\text{CH}_2)_{0-4}-\text{CO}-\text{R}_{1-\text{heteroaryl}}$ where $\text{R}_{1-\text{heteroaryl}}$ is as

5 defined above,

(14) $-(\text{CH}_2)_{0-4}-\text{CO}-\text{R}_{1-\text{heterocycle}}$ where $\text{R}_{1-\text{heterocycle}}$ is as

defined above,

(15) $-(\text{CH}_2)_{0-4}-\text{CO}-\text{R}_{\text{N}-4}$ where $\text{R}_{\text{N}-4}$ is selected from

the group consisting of morpholinyl, thiomorpholinyl, piperazinyl, piperidinyl,

10 homomorpholinyl, homothiomorpholinyl, homothiomorpholinyl S-oxide,

homothiomorpholinyl S,S-dioxide, pyrrolinyl and pyrrolidinyl where each group is optionally substituted with one, two, three, or four of C_1-C_6 alkyl,

(16) $-(\text{CH}_2)_{0-4}-\text{CO}-\text{O}-\text{R}_{\text{N}-5}$ where $\text{R}_{\text{N}-5}$ is

selected from the group consisting of:

15

(a) C_1-C_6 alkyl,

(b) $-(\text{CH}_2)_{0-2}-(\text{R}_{1-\text{aryl}})$ where $\text{R}_{1-\text{aryl}}$ is as defined

above,

(c) C_2-C_6 alkenyl containing one or two double

bonds,

20

(d) C_2-C_6 alkynyl containing one or two triple

bonds,

(e) C_3-C_7 cycloalkyl, and

(f) $-(\text{CH}_2)_{0-2}-(\text{R}_{1-\text{heteroaryl}})$ where $\text{R}_{1-\text{heteroaryl}}$ is as

defined above,

25

(17) $-(\text{CH}_2)_{0-4}-\text{SO}_2-\text{NR}_{\text{N}-2}\text{R}_{\text{N}-3}$ where $\text{R}_{\text{N}-2}$ and $\text{R}_{\text{N}-3}$ are

as defined above,

(18) $-(\text{CH}_2)_{0-4}-\text{SO}-(\text{C}_1-\text{C}_8 \text{ alkyl})$,

(19) $-(\text{CH}_2)_{0-4}-\text{SO}_2-(\text{C}_1-\text{C}_{12} \text{ alkyl})$,

(20) $-(\text{CH}_2)_{0-4}-\text{SO}_2-(\text{C}_3-\text{C}_7 \text{ cycloalkyl})$,

30

(21) $-(\text{CH}_2)_{0-4}-\text{N}(\text{H or } \text{R}_{\text{N}-5})-\text{CO}-\text{O}-\text{R}_{\text{N}-5}$ where $\text{R}_{\text{N}-5}$

can be the same or different and is as defined above,

(22) $-(\text{CH}_2)_{0-4}-\text{N}(\text{H or } \text{R}_{\text{N}-5})-\text{CO}-\text{N}(\text{R}_{\text{N}-5})_2$, where $\text{R}_{\text{N}-5}$

can be the same or different and is as defined above,

(23) $-(CH_2)_{0-4}-N-CS-N(R_{N-5})_2$, where R_{N-5} can be the same or different and is as defined above,

(24) $-(CH_2)_{0-4}-N(-H \text{ or } R_{N-5})-CO-R_{N-2}$ where R_{N-5} and R_{N-2} can be the same or different and are as defined above,

5 (25) $-(CH_2)_{0-4}-NR_{N-2}R_{N-3}$ where R_{N-2} and R_{N-3} can be the same or different and are as defined above,

(26) $-(CH_2)_{0-4}-R_{N-4}$ where R_{N-4} is as defined above,

(27) $-(CH_2)_{0-4}-O-CO-(C_1-C_6 \text{ alkyl})$,

(28) $-(CH_2)_{0-4}-O-P(O)-(OR_{N-aryl-1})_2$ where $R_{N-aryl-1}$ is --
10 H or C_1-C_4 alkyl,

(29) $-(CH_2)_{0-4}-O-CO-N(R_{N-5})_2$ where R_{N-5} is as defined above,

(30) $-(CH_2)_{0-4}-O-CS-N(R_{N-5})_2$ where R_{N-5} is as defined above,

15 (31) $-(CH_2)_{0-4}-O-(R_{N-5})_2$ where R_{N-5} is as defined above,

(32) $-(CH_2)_{0-4}-O-(R_{N-5})_2-COOH$ where R_{N-5} is as defined above,

(33) $-(CH_2)_{0-4}-S-(R_{N-5})_2$ where R_{N-5} is as defined
20 above,

(34) $-(CH_2)_{0-4}-O-(C_1-C_6 \text{ alkyl optionally substituted with one, two, three, four, or five of } -F)$,

(35) C_3-C_7 cycloalkyl,

(36) C_2-C_6 alkenyl with one or two double bonds
25 optionally substituted with C_1-C_3 alkyl, $-F$, $-Cl$, $-Br$, $-I$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, C_1-C_3 alkoxy, or $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

(37) C_2-C_6 alkynyl with one or two triple bonds
optionally substituted with C_1-C_3 alkyl, $-F$, $-Cl$, $-Br$, $-I$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, C_1-C_3 alkoxy, or $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

30 (38) $-(CH_2)_{0-4}-N(-H \text{ or } R_{N-5})-SO_2-R_{N-2}$ where R_{N-5} and R_{N-2} can be the same or different and are as described above, or

(39) $-(CH_2)_{0-4}-C_3-C_7$ cycloalkyl,

(B) -R_{N-heteroaryl} where R_{N-heteroaryl} is selected from the group

consisting of:

- | | |
|----|-------------------|
| | pyridinyl, |
| | pyrimidinyl, |
| 5 | quinolinyl, |
| | benzothienyl, |
| | indolyl, |
| | indolinyl, |
| | pyridazinyl, |
| 10 | pyrazinyl, |
| | isoindolyl, |
| | isoquinolyl, |
| | quinazolinyl, |
| | quinoxaliny, |
| 15 | phthalazinyl, |
| | imidazolyl, |
| | isoxazolyl, |
| | pyrazolyl, |
| | oxazolyl, |
| 20 | thiazolyl, |
| | indoliziny, |
| | indazolyl, |
| | benzothiazolyl, |
| | benzimidazolyl, |
| 25 | benzofuranyl, |
| | furanyl, |
| | thienyl, |
| | pyrroly, |
| | oxadiazolyl, |
| 30 | thiadiazolyl, |
| | triazolyl, |
| | tetrazolyl, |
| | oxazolopyridinyl, |
| | imidazopyridinyl, |

	isothiazolyl,
	naphthyridinyl,
	cinnoliny,
	carbazolyl,
5	beta-carbolinyl,
	isochromanyl,
	chromanyl,
	tetrahydroisoquinolinyl,
	isoindolinyl,
10	isobenzotetrahydrofuranyl,
	isobenzotetrahydrothienyl,
	isobenzothienyl,
	benzoxazolyl,
	pyridopyridinyl,
15	benzotetrahydrofuranyl,
	benzotetrahydrothienyl,
	purinyl,
	benzodioxolyl,
	triazinyl,
20	phenoxazinyl,
	phenothiazinyl,
	pteridinyl,
	benzothiazolyl,
	imidazopyridinyl,
25	imidazothiazolyl,
	dihydrobenzisoaxazinyl,
	benzisoaxazinyl,
	benzoxazinyl,
	dihydrobenzisothiazinyl,
30	benzopyranyl,
	benzothiopyranyl,
	coumarinyl,
	isocoumarinyl,
	chromonyl,

5 chromanonyl,
pyridinyl-N-oxide,
tetrahydroquinoliny1,
dihydroquinoliny1,
dihydroquinolinonyl,
dihydroisoquinolinonyl,
dihydrocoumariny1,
dihydroisocoumariny1,
10 isoindolinonyl,
benzodioxany1,
benzoxazolinonyl,
pyrrolyl N-oxide,
pyrimidinyl N-oxide,
pyridazinyl N-oxide,
15 pyrazinyl N-oxide,
quinoliny1 N-oxide,
indolyl N-oxide,
indoliny1 N-oxide,
isoquinolyl N-oxide,
20 quinazoliny1 N-oxide,
quinoxaliny1 N-oxide,
phthalazinyl N-oxide,
imidazolyl N-oxide,
isoxazolyl N-oxide,
25 oxazolyl N-oxide,
thiazolyl N-oxide,
indoliziny1 N-oxide,
indazolyl N-oxide,
benzothiazolyl N-oxide,
30 benzimidazolyl N-oxide,
pyrrolyl N-oxide,
oxadiazolyl N-oxide,
thiadiazolyl N-oxide,
triazolyl N-oxide,

tetrazolyl N-oxide,

benzothiopyranyl S-oxide, and

benzothiopyranyl S,S-dioxide

where the $R_{N\text{-heteroaryl}}$ group is bonded by any atom of the

5 parent $R_{N\text{-heteroaryl}}$ group substituted by hydrogen such that the new bond to the $R_{N\text{-heteroaryl}}$ group replaces the hydrogen atom and its bond, where heteroaryl is optionally substituted with one, two, three, or four of:

(1) $C_1\text{-}C_6$ alkyl, optionally substituted with one, two or three substituents selected from the group consisting of $C_1\text{-}C_3$ alkyl, -F, -Cl, -Br, -I, 10 -OH, -SH, $-C\equiv N$, $-CF_3$, $C_1\text{-}C_3$ alkoxy, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

(2) -OH,

(3) $-NO_2$,

(4) -F, -Cl, -Br, or -I,

15 (5) -CO-OH,

(6) $-C\equiv N$,

(7) $-(CH_2)_{0-4}\text{-CO-NR}_{N-2}R_{N-3}$ where R_{N-2} and R_{N-3} are the same or different and are selected from the group consisting of:

(a) -H,

20 (b) $C_1\text{-}C_6$ alkyl optionally substituted with one substituent selected from the group consisting of:

(i) -OH, and

(ii) $-NH_2$,

25 (c) $C_1\text{-}C_6$ alkyl optionally substituted with one, two, or three -F, -Cl, -Br, -I,

(d) $C_3\text{-}C_7$ cycloalkyl,

(e) $-(C_1\text{-}C_2\text{ alkyl})\text{-}(C_3\text{-}C_7\text{ cycloalkyl})$,

(f) $-(C_1\text{-}C_6\text{ alkyl})\text{-O}\text{-}(C_1\text{-}C_3\text{ alkyl})$,

30 (g) $C_2\text{-}C_6$ alkenyl with one or two double bonds,

(h) $C_2\text{-}C_6$ alkynyl with one or two triple

bonds,

(i) $-C_1-C_6$ alkyl chain with one double bond

and one triple bond,

(j) $-R_{1-aryl}$ where R_{1-aryl} is as defined above,

(k) $-R_{1-heteroaryl}$ where $R_{1-heteroaryl}$ is as defined

5 above,

(8) $-(CH_2)_{0-4}-CO-(C_1-C_{12}$ alkyl),

(9) $-(CH_2)_{0-4}-CO-(C_2-C_{12}$ alkenyl with one, two or
three double bonds),

10 three triple bonds),

(10) $-(CH_2)_{0-4}-CO-(C_2-C_{12}$ alkynyl with one, two or

(11) $-(CH_2)_{0-4}-CO-(C_3-C_7$ cycloalkyl),

(12) $-(CH_2)_{0-4}-CO-R_{1-aryl}$ where R_{1-aryl} is as defined

above,

(13) $-(CH_2)_{0-4}-CO-R_{1-heteroaryl}$ where $R_{1-heteroaryl}$ is as

15 defined above,

(14) $-(CH_2)_{0-4}-CO-R_{1-heterocycle}$ where $R_{1-heterocycle}$ is as

defined above,

(15) $-(CH_2)_{0-4}-CO-R_{N-4}$ where R_{N-4} is selected from

the group consisting of morpholinyl, thiomorpholinyl, piperazinyl, piperidinyl,

20 homomorpholinyl, homothiomorpholinyl, homothiomorpholinyl S-oxide,

homothiomorpholinyl S,S-dioxide, pyrrolinyl and pyrrolidinyl where each group is
optionally substituted with one, two, three, or four of C_1-C_6 alkyl,

(16) $-(CH_2)_{0-4}-CO-O-R_{N-5}$ where R_{N-5} is selected from

the group consisting of:

25

(a) C_1-C_6 alkyl,

(b) $-(CH_2)_{0-2}-(R_{1-aryl})$ where R_{1-aryl} is as defined

above,

(c) C_2-C_6 alkenyl containing one or two double

bonds,

30

(d) C_2-C_6 alkynyl containing one or two triple

bonds,

(e) C_3-C_7 cycloalkyl, and

(f) $-(CH_2)_{0-2}-(R_{1-heteroaryl})$ where $R_{1-heteroaryl}$ is as

defined above,

- (17) $-(\text{CH}_2)_{0-4}-\text{SO}_2-\text{NR}_{\text{N}-2}\text{R}_{\text{N}-3}$ where $\text{R}_{\text{N}-2}$ and $\text{R}_{\text{N}-3}$ are as defined above,
- (18) $-(\text{CH}_2)_{0-4}-\text{SO}-(\text{C}_1-\text{C}_8 \text{ alkyl})$,
- (19) $-(\text{CH}_2)_{0-4}-\text{SO}_2-(\text{C}_1-\text{C}_{12} \text{ alkyl})$,
- 5 (20) $-(\text{CH}_2)_{0-4}-\text{SO}_2-(\text{C}_3-\text{C}_7 \text{ cycloalkyl})$,
- (21) $-(\text{CH}_2)_{0-4}-\text{N}(\text{H or } \text{R}_{\text{N}-5})-\text{CO}-\text{O}-\text{R}_{\text{N}-5}$ where $\text{R}_{\text{N}-5}$ can be the same or different and is as defined above,
- (22) $-(\text{CH}_2)_{0-4}-\text{N}(\text{H or } \text{R}_{\text{N}-5})-\text{CO}-\text{N}(\text{R}_{\text{N}-5})_2$, where $\text{R}_{\text{N}-5}$ can be the same or different and is as defined above,
- 10 (23) $-(\text{CH}_2)_{0-4}-\text{N}-\text{CS}-\text{N}(\text{R}_{\text{N}-5})_2$, where $\text{R}_{\text{N}-5}$ can be the same or different and is as defined above,
- (24) $-(\text{CH}_2)_{0-4}-\text{N}(-\text{H or } \text{R}_{\text{N}-5})-\text{CO}-\text{R}_{\text{N}-2}$ where $\text{R}_{\text{N}-5}$ and $\text{R}_{\text{N}-2}$ can be the same or different and are as defined above,
- (25) $-(\text{CH}_2)_{0-4}-\text{NR}_{\text{N}-2}\text{R}_{\text{N}-3}$ where $\text{R}_{\text{N}-2}$ and $\text{R}_{\text{N}-3}$ can be the same or different and are as defined above,
- 15 (26) $-(\text{CH}_2)_{0-4}-\text{R}_{\text{N}-4}$ where $\text{R}_{\text{N}-4}$ is as defined above,
- (27) $-(\text{CH}_2)_{0-4}-\text{O}-\text{CO}-(\text{C}_1-\text{C}_6 \text{ alkyl})$,
- (28) $-(\text{CH}_2)_{0-4}-\text{O}-\text{P}(\text{O})-(\text{OR}_{\text{N-aryl-1}})_2$ where $\text{R}_{\text{N-aryl-1}}$ is $-\text{H or } \text{C}_1-\text{C}_4 \text{ alkyl}$,
- 20 (29) $-(\text{CH}_2)_{0-4}-\text{O}-\text{CO}-\text{N}(\text{R}_{\text{N}-5})_2$ where $\text{R}_{\text{N}-5}$ is as defined above,
- (30) $-(\text{CH}_2)_{0-4}-\text{O}-\text{CS}-\text{N}(\text{R}_{\text{N}-5})_2$ where $\text{R}_{\text{N}-5}$ is as defined above,
- (31) $-(\text{CH}_2)_{0-4}-\text{O}-(\text{R}_{\text{N}-5})_2$ where $\text{R}_{\text{N}-5}$ is as defined above,
- 25 (32) $-(\text{CH}_2)_{0-4}-\text{O}-(\text{R}_{\text{N}-5})_2-\text{COOH}$ where $\text{R}_{\text{N}-5}$ is as defined above,
- (33) $-(\text{CH}_2)_{0-4}-\text{S}-(\text{R}_{\text{N}-5})_2$ where $\text{R}_{\text{N}-5}$ is as defined above,
- 30 (34) $-(\text{CH}_2)_{0-4}-\text{O}-(\text{C}_1-\text{C}_6 \text{ alkyl optionally substituted with one, two, three, four, or five of } -\text{F})$,
- (35) $\text{C}_3-\text{C}_7 \text{ cycloalkyl}$,

- (36) C₂-C₆ alkenyl with one or two double bonds
optionally substituted with C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-
C₃ alkoxy, or -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,
- (37) C₂-C₆ alkynyl with one or two triple bonds
5 optionally substituted with C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-
C₃ alkoxy, or -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,
- (38) -(CH₂)₀₋₄-N(-H or R_{N-5})-SO₂-R_{N-2} where R_{N-5} and
R_{N-2} can be the same or different and are as defined above, or
- (39) -(CH₂)₀₋₄- C₃-C₇ cycloalkyl,
- 10 (C) R_{N-aryl}-W-R_{N-aryl}, where R_{N-aryl} can be the same or
different,
- (D) R_{N-aryl}-W-R_{N-heteroaryl},
- (E) R_{N-aryl}-W-R_{N-1-heterocycle}, wherein R_{N-1-heterocycle} is the same
as R_{1-heterocycle}, and R_{1-heterocycle} is as defined above
- 15 (F) R_{N-heteroaryl}-W-R_{N-aryl},
- (G) R_{N-heteroaryl}-W-R_{N-heteroaryl},
- (H) R_{N-heteroaryl}-W-R_{N-1-heterocycle},
- (I) R_{N-heterocycle}-W-R_{N-aryl}, wherein R_{N-heterocycle} is the same as
R_{1-heterocycle}, and R_{1-heterocycle} is as defined above, and R_{N-aryl} is as defined above,
- 20 (J) R_{N-heterocycle}-W-R_{N-heteroaryl}, and
- (K) R_{N-heterocycle}-W-R_{N-1-heterocycle},
- where W is
- (1) -(CH₂)₀₋₄-,
- (2) -O-,
- 25 (3) -S(O)₀₋₂-,
- (4) -N(R_{N-5})- where R_{N-5} is as defined above, or
- (5) -CO-;
- (II) -CO-(C₁-C₁₀ alkyl) where alkyl is optionally substituted with one
three substituents selected from the group consisting of:
- 30 (A) -OH,
- (B) -C₁-C₆ alkoxy,
- (C) -C₁-C₆ thioalkoxy,
- (D) -CO-O-R_{N-8} where R_{N-8} is -H, C₁-C₆ alkyl or -phenyl,

(E) $-\text{CO}-\text{NR}_{\text{N-2}}\text{R}_{\text{N-3}}$ where $\text{R}_{\text{N-2}}$ and $\text{R}_{\text{N-3}}$ are the same or different and are as defined above,

(F) $-\text{CO}-\text{R}_{\text{N-4}}$ where $\text{R}_{\text{N-4}}$ is as defined above,

(G) $-\text{SO}_2-(\text{C}_1-\text{C}_8 \text{ alkyl})$,

5 (H) $-\text{SO}_2-\text{NR}_{\text{N-2}}\text{R}_{\text{N-3}}$ where $\text{R}_{\text{N-2}}$ and $\text{R}_{\text{N-3}}$ are the same or different and are as defined above,

(I) $-\text{NH}-\text{CO}-(\text{C}_1-\text{C}_6 \text{ alkyl})$,

(J) $-\text{NH}-\text{CO}-\text{O}-\text{R}_{\text{N-8}}$ where $\text{R}_{\text{N-8}}$ is as defined above,

(K) $-\text{NR}_{\text{N-2}}\text{R}_{\text{N-3}}$ where $\text{R}_{\text{N-2}}$ and $\text{R}_{\text{N-3}}$ are the same or different
10 and are as defined above,

(L) $-\text{R}_{\text{N-4}}$ where $\text{R}_{\text{N-4}}$ is as defined above,

(M) $-\text{O}-\text{CO}-(\text{C}_1-\text{C}_6 \text{ alkyl})$,

(N) $-\text{O}-\text{CO}-\text{NR}_{\text{N-8}}\text{R}_{\text{N-8}}$ where $\text{R}_{\text{N-8}}$ are the same or different
and are as defined above,

15 (O) $-\text{O}-(\text{C}_1-\text{C}_5 \text{ alkyl})-\text{COOH}$,

(P) $-\text{O}-(\text{C}_1-\text{C}_6 \text{ alkyl})$ optionally substituted with one, two, or three of -F, -Cl, -Br, or -I),

(Q) $-\text{NH}-\text{SO}_2-(\text{C}_1-\text{C}_6 \text{ alkyl})$, and

(R) -F, or -Cl,

20 (III) $-\text{CO}-(\text{C}_1-\text{C}_6 \text{ alkyl})-\text{O}-(\text{C}_1-\text{C}_6 \text{ alkyl})$ where alkyl is optionally substituted with one, two, or three substituents selected from the group consisting of:

(A) -OH,

(B) $-\text{C}_1-\text{C}_6 \text{ alkoxy}$,

25 (C) $-\text{C}_1-\text{C}_6 \text{ thioalkoxy}$,

(D) $-\text{CO}-\text{O}-\text{R}_{\text{N-8}}$ where $\text{R}_{\text{N-8}}$ is -H, $\text{C}_1-\text{C}_6 \text{ alkyl}$ or -phenyl,

(E) $-\text{CO}-\text{NR}_{\text{N-2}}\text{R}_{\text{N-3}}$ where $\text{R}_{\text{N-2}}$ and $\text{R}_{\text{N-3}}$ are the same or different and are as defined above,

(F) $-\text{CO}-\text{R}_{\text{N-4}}$ where $\text{R}_{\text{N-4}}$ is as defined above,

30 (G) $-\text{SO}_2-(\text{C}_1-\text{C}_8 \text{ alkyl})$,

(H) $-\text{SO}_2-\text{NR}_{\text{N-2}}\text{R}_{\text{N-3}}$ where $\text{R}_{\text{N-2}}$ and $\text{R}_{\text{N-3}}$ are the same or different and are as defined above,

(I) $-\text{NH}-\text{CO}-(\text{C}_1-\text{C}_6 \text{ alkyl})$,

(J) -NH-CO-O-R_{N-8} where R_{N-8} is as defined above,

(K) $\text{-NR}_{N-2}\text{R}_{N-3}$ where R_{N-2} and R_{N-3} are the same or different

and are as defined above,

(L) -R_{N-4} where R_{N-4} is as defined above,

5 (M) $\text{-O-CO-(C}_1\text{-C}_6\text{ alkyl)}$,

(N) $\text{-O-CO-NR}_{N-8}\text{R}_{N-8}$ where R_{N-8} are the same or different

and are as defined above,

(O) $\text{-O-(C}_1\text{-C}_5\text{ alkyl)-COOH}$,

(P) $\text{-O-(C}_1\text{-C}_6\text{ alkyl optionally substituted with one, two, or}$

10 three of -F , -Cl , -Br , or -I),

(Q) $\text{-NH-SO}_2\text{-(C}_1\text{-C}_6\text{ alkyl)}$, and

(R) -F , or -Cl ,

(IV) $\text{-CO-(C}_1\text{-C}_6\text{ alkyl)-S-(C}_1\text{-C}_6\text{ alkyl)}$ where alkyl is optionally

substituted with one, two, or three of substituents selected from the group consisting

15 of:

(A) -OH ,

(B) $\text{-C}_1\text{-C}_6\text{ alkoxy}$,

(C) $\text{-C}_1\text{-C}_6\text{ thioalkoxy}$,

(D) -CO-O-R_{N-8} where R_{N-8} is as defined above,

20 (E) $\text{-CO-NR}_{N-2}\text{R}_{N-3}$ where R_{N-2} and R_{N-3} are the same or

different and are as defined above,

(F) -CO-R_{N-4} where R_{N-4} is as defined above,

(G) $\text{-SO}_2\text{-(C}_1\text{-C}_8\text{ alkyl)}$,

(H) $\text{-SO}_2\text{-NR}_{N-2}\text{R}_{N-3}$ where R_{N-2} and R_{N-3} are the same or

25 different and are as defined above,

(I) $\text{-NH-CO-(C}_1\text{-C}_6\text{ alkyl)}$,

(J) -NH-CO-O-R_{N-8} where R_{N-8} is as defined above,

(K) $\text{-NR}_{N-2}\text{R}_{N-3}$ where R_{N-2} and R_{N-3} are the same or different

and are as defined above,

30 (L) -R_{N-4} where R_{N-4} is as defined above,

(M) $\text{-O-CO-(C}_1\text{-C}_6\text{ alkyl)}$,

(N) $\text{-O-CO-NR}_{N-8}\text{R}_{N-8}$ where R_{N-8} are the same or different

and are as defined above,

(O) -O-(C₁-C₅ alkyl)-COOH,

(P) -O-(C₁-C₆ alkyl optionally substituted with one, two, or three of -F, -Cl, -Br, or -I),

(Q) -NH-SO₂-(C₁-C₆ alkyl), and

5 (R) -F, or -Cl,

(V) -CO-CH(-(CH₂)₀₋₂-O-R_{N-10})-(CH₂)₀₋₂-R_{N-aryl}/R_{N-heteroaryl}) where R_{N-aryl} and R_{N-heteroaryl} are as defined above, where R_{N-10} is selected from the group consisting of:

(A) -H,

10 (B) C₁-C₆ alkyl,

(C) C₃-C₇ cycloalkyl,

(D) C₂-C₆ alkenyl with one double bond,

(E) C₂-C₆ alkynyl with one triple bond,

(F) R_{1-aryl} where R_{1-aryl} is as defined above, and

15 (G) R_{N-heteroaryl} where R_{N-heteroaryl} is as defined above, or

(VI) -CO-(C₃-C₈ cycloalkyl) where alkyl is optionally substituted with one or two substituents selected from the group consisting of:

(A) -(CH₂)₀₋₄-OH,

(B) -(CH₂)₀₋₄-C₁-C₆ alkoxy,

20 (C) -(CH₂)₀₋₄-C₁-C₆ thioalkoxy,

(D) -(CH₂)₀₋₄-CO-O-R_{N-8} where R_{N-8} is -H, C₁-C₆ alkyl or phenyl,

(E) -(CH₂)₀₋₄-CO-NR_{N-2}R_{N-3} where R_{N-2} and R_{N-3} are the same or different and are as defined above,

25 (F) -(CH₂)₀₋₄-CO-R_{N-4} where R_{N-4} is as defined above,

(G) -(CH₂)₀₋₄-SO₂-(C₁-C₈ alkyl),

(H) -(CH₂)₀₋₄-SO₂-NR_{N-2}R_{N-3} where R_{N-2} and R_{N-3} are the same or different and are as defined above,

(I) -(CH₂)₀₋₄-NH-CO-(C₁-C₆ alkyl),

30 (J) -NH-CO-O-R_{N-8} where R_{N-8} is as defined above,

(K) -(CH₂)₀₋₄-NR_{N-2}R_{N-3} where R_{N-2} and R_{N-3} are the same or different and are as defined above,

(L) $-(\text{CH}_2)_{0-4}-\text{R}_{\text{N}-4}$ where $\text{R}_{\text{N}-4}$ is as defined above,

(M) $-\text{O}-\text{CO}-(\text{C}_1-\text{C}_6 \text{ alkyl})$,

(N) $-\text{O}-\text{CO}-\text{NR}_{\text{N}-8}\text{R}_{\text{N}-8}$ where $\text{R}_{\text{N}-8}$ are the same or different

and are as defined above,

5 (O) $-\text{O}-(\text{C}_1-\text{C}_5 \text{ alkyl})-\text{COOH}$,

(P) $-\text{O}-(\text{C}_1-\text{C}_6 \text{ alkyl})$ optionally substituted with one, two, or three of $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, or $-\text{I}$,

(Q) $-\text{NH}-\text{SO}_2-(\text{C}_1-\text{C}_6 \text{ alkyl})$, and

(R) $-\text{F}$, or $-\text{Cl}$;

10

where R_A is:

(I) C_1-C_{10} alkyl optionally substituted with one, two or three substituents selected from the group consisting of C_1-C_3 alkyl, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, $-\text{OH}$, $-\text{SH}$, $-\text{C}\equiv\text{N}$, $-\text{CF}_3$, C_1-C_6 alkoxy, $-\text{O}$ -phenyl, $-\text{NR}_{1-\text{a}}\text{R}_{1-\text{b}}$ where $\text{R}_{1-\text{a}}$ and $\text{R}_{1-\text{b}}$ are as defined above, $-\text{OC}=\text{O NR}_{1-\text{a}}\text{R}_{1-\text{b}}$ where $\text{R}_{1-\text{a}}$ and $\text{R}_{1-\text{b}}$ are as defined above, $-\text{S}(=\text{O})_{0-2}$ $\text{R}_{1-\text{a}}$ where $\text{R}_{1-\text{a}}$ is as defined above, $-\text{NR}_{1-\text{a}}\text{C}=\text{O NR}_{1-\text{a}}\text{R}_{1-\text{b}}$ where $\text{R}_{1-\text{a}}$ and $\text{R}_{1-\text{b}}$ are as defined above, $-\text{C}=\text{O NR}_{1-\text{a}}\text{R}_{1-\text{b}}$ where $\text{R}_{1-\text{a}}$ and $\text{R}_{1-\text{b}}$ are as defined above, and $-\text{S}(=\text{O})_2 \text{NR}_{1-\text{a}}\text{R}_{1-\text{b}}$ where $\text{R}_{1-\text{a}}$ and $\text{R}_{1-\text{b}}$ are as defined above,

(II) $-(\text{CH}_2)_{0-3}-(\text{C}_3-\text{C}_8)$ cycloalkyl where cycloalkyl can be optionally substituted with one, two or three substituents selected from the group consisting of C_1-C_3 alkyl, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, $-\text{OH}$, $-\text{SH}$, $-\text{C}\equiv\text{N}$, $-\text{CF}_3$, C_1-C_6 alkoxy, $-\text{O}$ -phenyl, $-\text{CO}-\text{OH}$, $-\text{CO}-\text{O}-(\text{C}_1-\text{C}_4 \text{ alkyl})$, and $-\text{NR}_{1-\text{a}}\text{R}_{1-\text{b}}$ where $\text{R}_{1-\text{a}}$ and $\text{R}_{1-\text{b}}$ are as defined above,

(III) $-(\text{CR}_{\text{A}-\text{x}}\text{R}_{\text{A}-\text{y}})_{0-4}-\text{R}_{\text{A}-\text{aryl}}$ where $\text{R}_{\text{A}-\text{x}}$ and $\text{R}_{\text{A}-\text{y}}$ are

25 (A) $-\text{H}$,

(B) C_1-C_4 alkyl optionally substituted with one or two $-\text{OH}$,

(C) C_1-C_4 alkoxy optionally substituted with one, two, or three of $-\text{F}$,

(D) $-(\text{CH}_2)_{0-4}-\text{C}_3-\text{C}_7$ cycloalkyl,

30 (E) C_2-C_6 alkenyl containing one or two double bonds,

(F) C_2-C_6 alkynyl containing one or two triple bonds, or

(G) phenyl,

and where R_{A-x} and R_{A-y} are taken together with the carbon to which they are attached to form a carbocycle of three, four, five, six or seven carbon atoms, optionally where one carbon atom is replaced by a heteroatom selected from the group consisting of -O-, -S-, -SO₂-, and -NR_{N-2}- and R_{A-aryl} is the same as R_{N-aryl} ,

5 (IV) $-(CR_{A-x}R_{A-y})_{0-4}-R_{A-heteroaryl}$ where $R_{A-heteroaryl}$ is the same as $R_{N-heteroaryl}$ and R_{A-x} and R_{A-y} are as defined above,

(V) $-(CR_{A-x}R_{A-y})_{0-4}-R_{A-aryl}-R_{A-aryl}$ where R_{A-aryl} , R_{A-x} and R_{A-y} are as defined above,

10 (VI) $-(CR_{A-x}R_{A-y})_{0-4}-R_{A-aryl}-R_{A-heteroaryl}$ where R_{A-aryl} , $R_{A-heteroaryl}$, R_{A-x} and R_{A-y} are as defined above,

(VII) $-(CR_{A-x}R_{A-y})_{0-4}-R_{A-heteroaryl}-R_{A-aryl}$ where $R_{A-heteroaryl}$, R_{A-aryl} , R_{A-x} and R_{A-y} are as defined above,

(VIII) $-(CR_{A-x}R_{A-y})_{0-4}-R_{A-heteroaryl}-R_{A-heteroaryl}$ where $R_{A-heteroaryl}$, R_{A-x} and R_{A-y} are as defined above,

15 (IX) $-(CR_{A-x}R_{A-y})_{0-4}-R_{A-aryl}-R_{A-heterocycle}$ where $R_{A-heterocycle}$ is defined as $R_{1-heterocycle}$, and where R_{A-aryl} , R_{A-x} and R_{A-y} are as defined above,

(X) $-(CR_{A-x}R_{A-y})_{0-4}-R_{A-heteroaryl}-R_{A-heterocycle}$ where $R_{A-heteroaryl}$, $R_{A-heterocycle}$, R_{A-x} and R_{A-y} are as defined above,

20 (XI) $-(CR_{A-x}R_{A-y})_{0-4}-R_{A-heterocycle}-R_{A-aryl}$ where $R_{A-heterocycle}$, R_{A-aryl} , R_{A-x} and R_{A-y} are as defined above,

(XII) $-(CR_{A-x}R_{A-y})_{0-4}-R_{A-heterocycle}-R_{A-heteroaryl}$ where $R_{A-heterocycle}$, $R_{A-heteroaryl}$, R_{A-x} and R_{A-y} are as defined above,

(XIII) $-(CR_{A-x}R_{A-y})_{0-4}-R_{A-heterocycle}-R_{A-heterocycle}$ where $R_{A-heterocycle}$, R_{A-x} and R_{A-y} are as defined above,

25 (XIV) $-(CR_{A-x}R_{A-y})_{0-4}-R_{A-heterocycle}$ where $R_{A-heterocycle}$, R_{A-x} and R_{A-y} are as defined above,

(XV) $-[C(R_{A-1})(R_{A-2})]_{1-3}-CO-N(R_{A-3})_2$ where R_{A-1} and R_{A-2} are the same or different and are selected from the group consisting of:

(A) -H,

30 (B) -C₁-C₆ alkyl, optionally substituted with one, two or three substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(C) C₂-C₆ alkenyl with one or two double bonds, optionally substituted with one, two or three substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

5 (D) C₂-C₆ alkynyl with one or two triple bonds, optionally substituted with one, two or three substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(E) -(CH₂)₁₋₂-S(O)₀₋₂-(C₁-C₆ alkyl),

10 (F) -(CH₂)₀₋₄-C₃-C₇ cycloalkyl, optionally substituted with one, two or three substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(G) -(C₁-C₄ alkyl)-R_{A'-aryl} where R_{A'-aryl} is as defined for R₁.

15 aryl,

(H) -(C₁-C₄ alkyl)-R_{A-heteroaryl} where R_{A-heteroaryl} is as defined

above,

(I) -(C₁-C₄ alkyl)-R_{A-heterocycle} where R_{A-heterocycle} is as defined

above,

20 (J) -R_{A-heteroaryl} where R_{A-heteroaryl} is as defined above,

(K) -R_{A-heterocycle} where R_{A-heterocycle} is as defined above,

(M) -(CH₂)₁₋₄-R_{A-4}-(CH₂)₀₋₄-R_{A'-aryl} where R_{A-4} is -O-, -S- or -NR_{A-5}- where R_{A-5} is C₁-C₆ alkyl, and where R_{A'-aryl} is defined above,

25 (N) -(CH₂)₁₋₄-R_{A-4}-(CH₂)₀₋₄-R_{A-heteroaryl} where R_{A-4} and R_{A-heteroaryl} are as defined above, and

(O) -R_{A'-aryl} where R_{A'-aryl} is as defined above,

and where R_{A-3} is the same or different and is:

(A) -H,

30 (B) -C₁-C₆ alkyl optionally substituted with one, two or three substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(C) C₂-C₆ alkenyl with one or two double bonds, optionally substituted with one, two or three substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

5 (D) C₂-C₆ alkynyl with one or two triple bonds, optionally substituted with one, two or three substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(E) -(CH₂)₀₋₄-C₃-C₇ cycloalkyl, optionally substituted with
10 one, two or three substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(F) -R_{A'-aryl} where R_{A'-aryl} is as defined above,

(G) -R_{A-heteroaryl} where R_{A-heteroaryl} is as defined above,

15 (H) -R_{A-heterocycle} where R_{A-heterocycle} is as defined above,

(I) -(C₁-C₄ alkyl)-R_{A'-aryl} where R_{A'-aryl} is as defined
above,

(J) -(C₁-C₄ alkyl)-R_{A-heteroaryl} where R_{A-heteroaryl} is as defined
above,

20 (K) -(C₁-C₄ alkyl)-R_{A-heterocycle} where R_{A-heterocycle} is as defined
above, or

(XVI) -CH(R_{A-aryl})₂ where R_{A-aryl} are the same or different and are as
defined above,

(XVII) -CH(R_{A-heteroaryl})₂ where R_{A-heteroaryl} are the same or different
25 and are as defined above,

(XVIII) -CH(R_{A-aryl})(R_{A-heteroaryl}) where R_{A-aryl} and R_{A-heteroaryl} are as
defined above,

(XIX) -cyclopentyl, -cyclohexyl, or -cycloheptyl ring fused to R_{A-aryl},
R_{A-heteroaryl}, R_{A-heterocycle} where R_{A-aryl} or R_{A-heteroaryl} or R_{A-heterocycle} are as defined above
30 where one carbon of cyclopentyl, cyclohexyl, or -cycloheptyl is optionally replaced
with NH, NR_{N-5}, O, or S(=O)₀₋₂, and where cyclopentyl, cyclohexyl, or -cycloheptyl
can be optionally substituted with one or two -C₁-C₃ alkyl, -F, -OH, -SH, -C≡N, -
CF₃, C₁-C₆ alkoxy, =O, or -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(XX) C₂-C₁₀ alkenyl containing one or two double bonds optionally substituted with one, two or three substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

5 (XXI) C₂-C₁₀ alkynyl containing one or two triple bonds optionally substituted with one, two or three substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(XXI) -(CH₂)₀₋₁-CHR_{A-6}-(CH₂)₀₋₁-R_{A-aryl} where R_{A-aryl} is as defined
10 above and R_{A-6} is -(CH₂)₀₋₆-OH,

(XXII) -(CH₂)₀₋₁-CHR_{A-6}-(CH₂)₀₋₁-R_{A-heteroaryl} where R_{A-heteroaryl} and R_{A-6} is as defined above,

(XXIII) -CH(-R_{A-aryl} or R_{A-heteroaryl})-CO-O(C₁-C₄ alkyl) where R_{A-aryl} and R_{A-heteroaryl} are as defined above,

15 (XXIV) -CH(-CH₂-OH)-CH(-OH)-micro-NO₂,

(XXV) (C₁-C₆ alkyl)-O-(C₁-C₆ alkyl)-OH,

(XXVII) -CH₂-NH-CH₂-CH(-O-CH₂-CH₃)₂,

(XXVIII) -H,

(XXIX) -(CH₂)₀₋₆-C(=NR_{1-a})(NR_{1-a}R_{1-b}) where R_{1-a} and R_{1-b} are as
20 defined above; or

(XXX)

-C=OC(HR₆)NHR₇, where R₆ and R₇ are as defined below,

-C=OR₇, where R₇ is as defined below,

-C=OOR₇, where R₇ is as defined below, or

25 -SOOR₇ where R₇ is as defined below,

wherein R₆ is:

hydrogen,

C₁ - C₃ alkyl,

phenyl,

30 thioalkoxyalkyl,

alkyl substituted aryl,

cycloalkyl,

cycloalkylalkyl,

hydroxyalkyl,
alkoxyalkyl,
aryloxyalkyl,
haloalkyl,
5 carboxyalkyl,
alkoxycarbonylalkyl,
aminoalkyl,
(N-protected)aminoalkyl,
alkylaminoalkyl,
10 ((N-protected)(alkyl)amino)alkyl,
dialkylaminoalkyl,
guanidinoalkyl,
lower alkenyl,
heterocyclic,
15 (heterocyclic)alkyl),
arylthioalkyl,
arylsulfonylalkyl,
(heterocyclic)thioalkyl,
(heterocyclic)sulfonylalkyl,
20 (heterocyclic)oxyalkyl,
arylalkoxyalkyl,
arylthioalkoxyalkyl,
arylalkylsulfonylalkyl,
(heterocyclic))alkoxyalkyl,
25 (heterocyclic)thioalkoxyalkyl,
(heterocyclic)alkylsulfonylalkyl,
cycloalkyloxyalkyl,
cycloalkylthioalkyl,
cycloalkylsulfonylalkyl,
30 cycloalkylalkoxyalkyl,
cycloalkylthioalkoxyalkyl,
cycloalkylalkylsulfonylalkyl,
aminocarbonyl,
alkylaminocarbonyl,

- dialkylaminocarbonyl,
aroylalkyl,
(heterocyclic)carbonylalkyl,
polyhydroxyalkyl,
5 aminocarbonylalkyl,
alkylaminocarbonylalkyl,
dialkylaminocarbonylalkyl,
aryloxyalkyl, or
alkylsulfonylalkyl,
- 10 wherein heterocyclic is pyridyl, thiazolyl, isothiazolyl,
oxazolyl, isoxazolyl, furanyl, thienyl, tetrahydrofuranyl, tetrahydrothienyl and
tetrahydro[2H]pyranyl and wherein the heterocycle is unsubstituted or substituted
with one to three substituents independently selected from hydroxy, halo, amino,
alkylamino, dialkylamino, alkoxy, polyalkoxy, haloalkyl, cycloalkyl,
15 cycloalkylalkyl, aryl, arylalkyl, COOH, -SO₃H, lower alkenyl or lower alkyl;
wherein R₇ is:
C₁ - C₃ alkyl,
phenyl,
thioalkoxyalkyl,
20 (aryl)alkyl,
cycloalkyl,
cycloalkylalkyl,
hydroxyalkyl,
alkoxyalkyl,
25 aryloxyalkyl,
haloalkyl,
carboxyalkyl,
alkoxycarbonylalkyl,
aminoalkyl,
30 (N-protected)aminocalkyl,
alkylaminoalkyl,
(N-protected)(alkyl)amino)alkyl,
dialkylaminoalkyl,
guanidinoalkyl,

- lower alkenyl,
heterocyclic,
(heterocyclic)alkyl),
arylthioalkyl,
5 arylsulfonylalkyl,
(heterocyclic)thioalkyl,
(heterocyclic)sulfonylalkyl,
(heterocyclic)oxyalkyl,
arylalkoxyalkyl,
10 arylthioalkoxyalkyl,
arylalkylsulfonylalkyl,
(heterocyclic)alkoxyalkyl,
(heterocyclic)thioalkoxyalkyl,
(heterocyclic)alkylsulfonylalkyl,
15 cycloalkyloxyalkyl,
cycloalkylthioalkyl,
cycloalkylsulfonylalkyl,
cycloalkylalkoxyalkyl,
cycloalkylthioalkoxyalkyl,
20 cycloalkylalkylsulfonylalkyl,
aminocarbonyl,
alkylaminocarbonyl,
dialkylaminocarbonyl,
aroylalkyl,
25 (heterocyclic)carbonylalkyl,
polyhydroxyalkyl,
aminocarbonylalkyl,
alkylaminocarbonylalkyl,
dialkylaminocarbonylalkyl,
30 aryloxyalkyl, or
alkylsulfonylalkyl,
wherein heterocyclic is pyridyl, thiazolyl, isothiazolyl,
oxazolyl, isoxazolyl, furanyl, thienyl, tetrahydrofuranyl, tetrahydrothienyl and
tetrahydro[2H]pyranyl and wherein the heterocycle is unsubstituted or substituted

with one to three substituents independently selected from hydroxy, halo, amino, alkylamino, dialkylamino, alkoxy, polyalkoxy, haloalkyl, cycloalkyl, cycloalkylalkyl, aryl, arylalkyl, COOH, -SO₃H, lower alkenyl or lower alkyl;

5 where X is -N, or -O, with the proviso that when X is O, R_B is absent;
and when X is N,

R_B is:

(I)-C₁-C₁₀ alkyl optionally substituted with one, two or three
substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH,
10 -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as
defined above, -OC=O NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above, -S(=O)₀₋₂
R_{1-a} where R_{1-a} is as defined above, -NR_{1-a}C=O NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as
defined above, -C=O NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above, and -
S(=O)₂ NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

15 (II) -(CH₂)₀₋₃-(C₃-C₈) cycloalkyl where cycloalkyl can be optionally
substituted with one, two or three substituents selected from the group consisting of
C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, -CO-
OH, -CO-O-(C₁-C₄ alkyl), and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(III) -(CR_{B-x}R_{B-y})₀₋₄-R_{B-aryl} where R_{B-x} and R_{B-y} are

20 (A) -H,
 (B) C₁-C₄ alkyl optionally substituted with one or two -OH,
 (C) C₁-C₄ alkoxy optionally substituted with one, two, or
three of -F,

 (D) -(CH₂)₀₋₄-C₃-C₇ cycloalkyl,
25 (E) C₂-C₆ alkenyl containing one or two double bonds,
 (F) C₂-C₆ alkynyl containing one or two triple bonds, or
 (G) phenyl,

 and where R_{B-x} and R_{B-y} are taken together with the carbon to
which they are attached to form a carbocycle of three, four, five, six or seven carbon
30 atoms, optionally where one carbon atom is replaced by a heteroatom selected from
the group consisting of -O-, -S-, -SO₂-, and -NR_{N-2} where R_{N-2} is as defined above,
and R_{B-aryl} is the same as R_{N-aryl} and is defined above

(IV) $-(\text{CR}_{\text{B-x}}\text{R}_{\text{B-y}})_{0-4}-\text{R}_{\text{B-heteroaryl}}$ where $\text{R}_{\text{B-heteroaryl}}$ is the same as $\text{R}_{\text{N-heteroaryl}}$, $\text{R}_{\text{B-x}}$, and $\text{R}_{\text{B-y}}$ are as defined above,

(V) $-(\text{CR}_{\text{B-x}}\text{R}_{\text{B-y}})_{0-4}-\text{R}_{\text{B-aryl}}-\text{R}_{\text{B-aryl}}$ where $\text{R}_{\text{B-aryl}}$, $\text{R}_{\text{B-x}}$, and $\text{R}_{\text{B-y}}$ are as defined above,

5 (VI) $-(\text{CR}_{\text{B-x}}\text{R}_{\text{B-y}})_{0-4}-\text{R}_{\text{B-aryl}}-\text{R}_{\text{B-heteroaryl}}$ where $\text{R}_{\text{B-aryl}}$, $\text{R}_{\text{B-heteroaryl}}$, $\text{R}_{\text{B-x}}$ and $\text{R}_{\text{B-y}}$ are as defined above,

(VII) $-(\text{CR}_{\text{B-x}}\text{R}_{\text{B-y}})_{0-4}-\text{R}_{\text{B-heteroaryl}}-\text{R}_{\text{B-aryl}}$ where $\text{R}_{\text{B-heteroaryl}}$, $\text{R}_{\text{B-aryl}}$, $\text{R}_{\text{B-x}}$ and $\text{R}_{\text{B-y}}$ are as defined above,

10 (VIII) $-(\text{CR}_{\text{B-x}}\text{R}_{\text{B-y}})_{0-4}-\text{R}_{\text{B-heteroaryl}}-\text{R}_{\text{B-heteroaryl}}$ where $\text{R}_{\text{B-heteroaryl}}$, $\text{R}_{\text{B-x}}$ and $\text{R}_{\text{B-y}}$ are as defined above,

(IX) $-(\text{CR}_{\text{B-x}}\text{R}_{\text{B-y}})_{0-4}-\text{R}_{\text{B-aryl}}-\text{R}_{\text{B-heterocycle}}$ where $\text{R}_{\text{B-heterocycle}}$ is defined as $\text{R}_{1\text{-heterocycle}}$, and where $\text{R}_{\text{B-aryl}}$, $\text{R}_{\text{B-x}}$ and $\text{R}_{\text{B-y}}$ are as defined above,

(X) $-(\text{CR}_{\text{B-x}}\text{R}_{\text{B-y}})_{0-4}-\text{R}_{\text{B-heteroaryl}}-\text{R}_{\text{B-heterocycle}}$ where $\text{R}_{\text{B-heteroaryl}}$, $\text{R}_{\text{B-heterocycle}}$, $\text{R}_{\text{B-x}}$ and $\text{R}_{\text{B-y}}$ are as defined above,

15 (XI) $-(\text{CR}_{\text{B-x}}\text{R}_{\text{B-y}})_{0-4}-\text{R}_{\text{B-heterocycle}}-\text{R}_{\text{B-aryl}}$ where $\text{R}_{\text{B-heterocycle}}$, $\text{R}_{\text{B-aryl}}$, $\text{R}_{\text{B-x}}$ and $\text{R}_{\text{B-y}}$ are as defined above,

(XII) $-(\text{CR}_{\text{B-x}}\text{R}_{\text{B-y}})_{0-4}-\text{R}_{\text{B-heterocycle}}-\text{R}_{\text{B-heteroaryl}}$ where $\text{R}_{\text{B-heterocycle}}$, $\text{R}_{\text{B-heteroaryl}}$, $\text{R}_{\text{B-x}}$ and $\text{R}_{\text{B-y}}$ are as defined above,

20 (XIII) $-(\text{CR}_{\text{B-x}}\text{R}_{\text{B-y}})_{0-4}-\text{R}_{\text{B-heterocycle}}-\text{R}_{\text{B-heterocycle}}$ where $\text{R}_{\text{B-heterocycle}}$, $\text{R}_{\text{B-x}}$ and $\text{R}_{\text{B-y}}$ are as defined above,

(XIV) $-(\text{CR}_{\text{B-x}}\text{R}_{\text{B-y}})_{0-4}-\text{R}_{\text{B-heterocycle}}$ where $\text{R}_{\text{B-heterocycle}}$, $\text{R}_{\text{B-x}}$ and $\text{R}_{\text{B-y}}$ are as defined above,

(XV) $-[\text{C}(\text{R}_{\text{B-1}})(\text{R}_{\text{B-2}})]_{1-3}-\text{CO}-\text{N}(\text{R}_{\text{B-3}})_2$ where $\text{R}_{\text{B-1}}$ and $\text{R}_{\text{B-2}}$ are the same or different and are selected from the group consisting of:

25 (A) -H,

(B) $-\text{C}_1-\text{C}_6$ alkyl, optionally substituted with one, two or three substituents selected from the group consisting of C_1-C_3 alkyl, -F, -Cl, -Br, -I, -OH, -SH, $-\text{C}\equiv\text{N}$, $-\text{CF}_3$, C_1-C_6 alkoxy, -O-phenyl, and $-\text{NR}_{1\text{-a}}\text{R}_{1\text{-b}}$ where $\text{R}_{1\text{-a}}$ and $\text{R}_{1\text{-b}}$ are as defined above,

30 (C) C_2-C_6 alkenyl with one or two double bonds, optionally substituted with one, two or three substituents selected from the group consisting of C_1-C_3 alkyl, -F, -Cl, -Br, -I, -OH, -SH, $-\text{C}\equiv\text{N}$, $-\text{CF}_3$, C_1-C_6 alkoxy, -O-phenyl, and $-\text{NR}_{1\text{-a}}\text{R}_{1\text{-b}}$ where $\text{R}_{1\text{-a}}$ and $\text{R}_{1\text{-b}}$ are as defined above,

- (D) C₂-C₆ alkynyl with one or two triple bonds, optionally substituted with one, two or three substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,
- 5 (E) -(CH₂)₁₋₂-S(O)₀₋₂-(C₁-C₆ alkyl),
 (F) -(CH₂)₀₋₄-C₃-C₇ cycloalkyl, optionally substituted with one, two or three substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,
- 10 (G) -(C₁-C₄ alkyl)-R_{B'-aryl} where R_{B'-aryl} is as defined above for R_{1-aryl},
 (H) -(C₁-C₄ alkyl)-R_{B-heteroaryl} where R_{B-heteroaryl} is as defined above,
 (I) -(C₁-C₄ alkyl)-R_{B-heterocycle} where R_{B-heterocycle} is as defined
 15 above,
- (J) -R_{B-heteroaryl} where R_{B-heteroaryl} is as defined above,
 (K) -R_{B-heterocycle} where R_{B-heterocycle} is as defined above,
 (M) -(CH₂)₁₋₄-R_{B-4}-(CH₂)₀₋₄-R_{B'-aryl} where R_{B-4} is -O-, -S- or -NR_{B-5}- where R_{B-5} is C₁-C₆ alkyl, and where R_{B'-aryl} is defined above,
- 20 (N) -(CH₂)₁₋₄-R_{B-4}-(CH₂)₀₋₄-R_{B-heteroaryl} where R_{B-4} and R_{B-heteroaryl} are as defined above, and
 (O) -R_{B'-aryl} where R_{B'-aryl} is as defined above,
 and where R_{B-3} is the same or different and is:
- (A) -H,
 25 (B) -C₁-C₆ alkyl optionally substituted with one, two or three substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,
- (C) C₂-C₆ alkenyl with one or two double bonds, optionally
 30 substituted with one, two or three substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(D) C₂-C₆ alkynyl with one or two triple bonds, optionally substituted with one, two or three substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

5 (E) -(CH₂)₀₋₄-C₃-C₇ cycloalkyl, optionally substituted with one, two or three substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(F) -R_{B'-aryl} where R_{B'-aryl} is as defined above,

10 (G) -R_{B-heteroaryl} where R_{B-heteroaryl} is as defined above,

(H) -R_{B-heterocycle} where R_{B-heterocycle} is as defined above,

(I) -(C₁-C₄ alkyl)-R_{B'-aryl} where R_{B'-aryl} is as defined above,

(J) -(C₁-C₄ alkyl)-R_{B-heteroaryl} where R_{B-heteroaryl} is as defined above,

(K) -(C₁-C₄ alkyl)-R_{B-heterocycle} where R_{B-heterocycle} is as defined above, or

(XVI) -CH(R_{B-aryl})₂ where R_{B-aryl} are the same or different and are as defined above,

20 (XVII) -CH(R_{B-heteroaryl})₂ where R_{B-heteroaryl} are the same or different and are as defined above,

(XVIII) -CH(R_{B-aryl})(R_{B-heteroaryl}) where R_{B-aryl} and R_{B-heteroaryl} are as defined above,

(XIX) -cyclopentyl, -cyclohexyl, or -cycloheptyl ring fused to R_{B-aryl} or R_{B-heteroaryl} or R_{B-heterocycle} where R_{B-aryl} or R_{B-heteroaryl} or R_{B-heterocycle} are as defined above where one carbon of cyclopentyl, cyclohexyl, or -cycloheptyl is optionally replaced with NH, NR_{N-5}, O, or S(=O)₀₋₂, and where cyclopentyl, cyclohexyl, or -cycloheptyl can be optionally substituted with one or two -C₁-C₃ alkyl, -F, -OH, -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, =O, or -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(XX) C₂-C₁₀ alkenyl containing one or two double bonds optionally substituted with one, two or three substituents selected from the group consisting of

C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(XXI) C₂-C₁₀ alkynyl containing one or two triple bonds optionally substituted with one, two or three substituents selected from the group consisting of

5 C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(XXI) -(CH₂)₀₋₁-CHR_{C-6}-(CH₂)₀₋₁-RB_{B-aryl} where RB_{B-aryl} is as defined above and R_{C-6} is -(CH₂)₀₋₆-OH,

(XXII) -(CH₂)₀₋₁-CHR_{B-6}-(CH₂)₀₋₁-RB_{B-heteroaryl} where RB_{B-heteroaryl} and
10 R_{C-6} is as defined above,

(XXIII) -CH(-RB_{B-aryl} or RB_{B-heteroaryl})-CO-O(C₁-C₄ alkyl) where RB_{B-aryl} and RB_{B-heteroaryl} are as defined above,

(XXIV) -CH(-CH₂-OH)-CH(-OH)-micro-NO₂,

(XXV) (C₁-C₆ alkyl)-O-(C₁-C₆ alkyl)-OH,

15 (XXVII) -CH₂-NH-CH₂-CH(-O-CH₂-CH₃)₂,

(XXVIII) -H, or

(XXIX) -(CH₂)₀₋₆-C(=NR_{1-a})(NR_{1-a}R_{1-b}) where R_{1-a} and R_{1-b} are as defined above;

or a pharmaceutically acceptable salt thereof.

20

2. A substituted amine according to claim 1

where R₁ is:

-(CH₂)₀₋₁-(R_{1-aryl}), or

-(CH₂)_{n1}-(R_{1-heteroaryl});

25 where R_N is:

R_{N-1}-X_N-, where X_N is selected from the group consisting of:

-CO-, and

-SO₂-,

where R_{N-1} is selected from the group consisting of:

30 -R_{N-aryl}, and

-R_{N-heteroaryl}, or

-CO-CH(-(CH₂)₀₋₂-O-R_{N-10})-(CH₂)₀₋₂-R_{N-aryl}/R_{N-heteroaryl});

where R_A is:

- C₁-C₈ alkyl,
 -(CH₂)₀₋₃-(C₃-C₇) cycloalkyl,
 -(CR_{A-x}R_{A-y})₀₋₄-R_{A-aryl},
 -(CR_{A-x}R_{A-y})₀₋₄-R_{A-heteroaryl},
 5 -(CR_{A-x}R_{A-y})₀₋₄-R_{A-heterocycle},
 -cyclopentyl or -cyclohexyl ring fused to R_{A-aryl} or R_{A-heteroaryl} or R_{A-}
 heterocycle;

where X is -N or -O, with the proviso that when X is O, R_B is absent;

and when X is N,

- 10 R_B is:
 -C₁-C₈ alkyl,
 -(CH₂)₀₋₃-(C₃-C₇) cycloalkyl,
 -(CR_{A-x}R_{A-y})₀₋₄-R_{A-aryl},
 -(CR_{A-x}R_{A-y})₀₋₄-R_{A-heteroaryl},
 15 -(CR_{A-x}R_{A-y})₀₋₄-R_{A-heterocycle},
 -cyclopentyl or -cyclohexyl ring fused to R_{A-aryl} or R_{A-heteroaryl} or R_{A-}
 heterocycle.

3. A substituted amine according to claim 2

- 20 where R₁ is:
 -(CH₂)-(R_{1-aryl}), or
 -(CH₂)-(R_{1-heteroaryl});

where R₂ is -H;

where R₃ is -H;

- 25 where R_N is:
 R_{N-1}-X_N where X_N is:

-CO-,

where R_{N-1} is selected from the group consisting of:

-R_{N-aryl}, and

- 30 -R_{N-heteroaryl};

where R_A is:

- C₁-C₈ alkyl,
 -(CH₂)₀₋₃-(C₃-C₇) cycloalkyl,
 -(CR_{A-x}R_{A-y})₀₋₄-R_{A-aryl},

$-(\text{CR}_{\text{A-x}}\text{R}_{\text{A-y}})_{0-4}-\text{R}_{\text{A-heteroaryl}}$,
 $-(\text{CR}_{\text{A-x}}\text{R}_{\text{A-y}})_{0-4}-\text{R}_{\text{A-heterocycle}}$,
 -cyclopentyl or -cyclohexyl ring fused to $\text{R}_{\text{A-aryl}}$ or $\text{R}_{\text{A-heteroaryl}}$ or $\text{R}_{\text{A-heterocycle}}$;

5 where X is -N or -O, with the proviso that when X is O, R_{B} is absent;
 and when X is N,

R_{B} is:

$-\text{C}_1-\text{C}_8$ alkyl,
 $-(\text{CH}_2)_{0-3}-(\text{C}_3-\text{C}_7)$ cycloalkyl,
 10 $-(\text{CR}_{\text{B-x}}\text{R}_{\text{B-y}})_{0-4}-\text{R}_{\text{B-aryl}}$,
 $-(\text{CR}_{\text{B-x}}\text{R}_{\text{B-y}})_{0-4}-\text{R}_{\text{B-heteroaryl}}$,
 $-(\text{CR}_{\text{B-x}}\text{R}_{\text{B-y}})_{0-4}-\text{R}_{\text{B-heterocycle}}$,
 -cyclopentyl or -cyclohexyl ring fused to $\text{R}_{\text{B-aryl}}$ or $\text{R}_{\text{B-heteroaryl}}$ or $\text{R}_{\text{B-heterocycle}}$.

15

4. A substituted amine according to claim 3,

where R_{A} is:

$-(\text{CR}_{\text{A-x}}\text{R}_{\text{A-y}})_{0-4}-\text{R}_{\text{A-aryl}}$,
 $-(\text{CR}_{\text{A-x}}\text{R}_{\text{A-y}})_{0-4}-\text{R}_{\text{A-heteroaryl}}$, or
 20 -cyclopentyl or -cyclohexyl ring fused to a $\text{R}_{\text{A-aryl}}$ or $\text{R}_{\text{A-heteroaryl}}$ or $\text{R}_{\text{A-heterocycle}}$; and

where R_{B} is:

$-(\text{CR}_{\text{B-x}}\text{R}_{\text{B-y}})_{0-4}-\text{R}_{\text{B-aryl}}$,
 $-(\text{CR}_{\text{B-x}}\text{R}_{\text{B-y}})_{0-4}-\text{R}_{\text{B-heteroaryl}}$, or
 25 -cyclopentyl or -cyclohexyl ring fused to $\text{R}_{\text{B-aryl}}$ or $\text{R}_{\text{B-heteroaryl}}$ or $\text{R}_{\text{B-heterocycle}}$.

5. A substituted amine according to claim 1 where R_1 is

$-(\text{CH}_2)-(\text{R}_{1\text{-aryl}})$ where $\text{R}_{1\text{-aryl}}$ is phenyl.

30

6. A substituted amine according to claim 1 where R_1 is

$-(\text{CH}_2)-(\text{R}_{1\text{-aryl}})$ where $\text{R}_{1\text{-aryl}}$ is phenyl substituted with two -F.

7. A substituted amine according to claim 6 where the -F substitution is 3,5-difluorobenzyl.
8. A substituted amine according to claim 1 where R_2 is -H.
- 5 9. A substituted amine according to claim 1 where R_3 is -H.
- 10 10. A substituted amine according to claim 1 where R_N is $R_{N-1}-X_N-$, where X_N is -CO-, where R_{N-1} is R_{N-aryl} where R_{N-aryl} is phenyl substituted with one -CO-NR_{N-2}R_{N-3} where the substitution on phenyl is 1,3-.
11. A substituted amine according to claim 10 where R_{N-2} and R_{N-3} are the same and are C₃ alkyl.
- 15 12. A substituted amine according to claim 1 where R_N is $R_{N-1}-X_N-$ where X_N is -CO-, and where R_{N-1} is R_{N-aryl} where R_{N-aryl} is phenyl substituted with one C₁ alkyl and with one -CO-NR_{N-2}R_{N-3} where the substitution on the phenyl is 1,3,5-.
- 20 13. A substituted amine according to claim 12 where R_{N-2} and R_{N-3} are the same and are C₃ alkyl.
14. A substituted amine according to claim 1 where R_N is $R_{N-1}-X_N-$, where X_N is -CO-, and where R_{N-1} is $R_{N-heteroaryl}$ where $R_{N-heteroaryl}$ is substituted with one -CO-NR_{N-2}R_{N-3}.
- 25 15. A substituted amine according to claim 14 where R_{N-2} and R_{N-3} are the same and are -C₃ alkyl.
- 30 16. A substituted amine according to claim 1 where R_A is:
 -(CR_{A-x}R_{A-y})₀₋₄-R_{A-aryl} where R_{A-aryl} is phenyl,
 -(CR_{A-x}R_{A-y})₀₋₄-R_{A-heteroaryl},
 -cyclopentyl or -cyclohexyl ring fused to a R_{A-aryl} or R_{A-heteroaryl} or R_{A-heterocycle}.

17. A substituted amine according to claim 16 where R_A is:
 $-(CR_{A-x}R_{A-y})_{0-4}-R_{A-aryl}$ where R_{A-aryl} is phenyl.
- 5 18. A substituted amine according to claim 17 where phenyl is substituted in the 3-position or 3,5-positions.
19. A substituted amine according to claim 16 where R_A is
 $-(CH_2)-R_{A-heteroaryl}$.
- 10 20. A substituted amine according to claim 16 where R_A is:
 $-(CH_2)-R_{A-heterocycle}$.
21. A substituted amine according to claim 16 where R_A is:
15 $-\text{cyclohexyl ring fused to a phenyl ring}$.
22. A substituted amine according to claim 1 where R_B is:
 $-(CR_{B-x}R_{B-y})_{0-4}-R_{B-aryl}$ where R_{B-aryl} is phenyl,
 $-(CR_{B-x}R_{B-y})_{0-4}-R_{B-heteroaryl}$,
20 $-\text{cyclopentyl or -cyclohexyl ring fused to a } R_{B-aryl} \text{ or } R_{B-heteroaryl} \text{ or } R_{B-}$
heterocycle.
23. A substituted amine according to claim 22 where R_B is:
 $-(CR_{B-x}R_{B-y})_{0-4}-R_{B-aryl}$ where R_{B-aryl} is phenyl.
- 25 24. A substituted amine according to claim 23 where phenyl is substituted in the 3-position or 3,5-positions.
25. A substituted amine according to claim 22 where R_B is:
30 $-(CH_2)-R_{B-heteroaryl}$.
26. A substituted amine according to claim 22 where R_B is:
 $-(CH_2)-R_{B-heterocycle}$.

27. A substituted amine according to claim 22 where R_B is:

-cyclohexyl ring fused to a phenyl ring.

28. A substituted amine according to claim 1, where R_B is absent.

5


29. A substituted amine according to claim 1 chosen from the group consisting of:

N-[1-(3,5-Difluoro-benzyl)-2-hydroxy-3-(N'-methyl-N'-phenyl-hydrazino)-propyl]-5-methyl-N', N'-dipropyl-isophthalamide,

10 N-{1-(3,5-Difluoro-benzyl)-2-hydroxy-3-[N'-methyl-N'-(4-methyl-pentanoyl)-hydrazino]-propyl}-5-methyl-N',N'-dipropyl-isophthalamide, and
N-[1-(3,5-Difluoro-benzyl)-2-hydroxy-3-phenoxyamino-propyl]-5-methyl-N',N'-dipropyl-isophthalamide.

30. A substituted amine according to claim 1 where the pharmaceutically acceptable
15 salt is selected from the group consisting of salts of the following acids acetic, aspartic, benzenesulfonic, benzoic, bicarbonic, bisulfuric, bitartaric, butyric, calcium edetate, camsyllic, carbonic, chlorobenzoic, citric, edetic, edisylic, estolic, esyl, esylic, formic, fumaric, gluceptic, gluconic, glutamic, glycolylarsanilic, hexamic, hexylresorcinoic, hydrabamic, hydrobromic, hydrochloric, hydroiodic,
20 hydroxynaphthoic, isethionic, lactic, lactobionic, maleic, malic, malonic, mandelic, methanesulfonic, methylnitric, methylsulfuric, mucic, muconic, napsylic, nitric, oxalic, p-nitromethanesulfonic, pamoic, pantothenic, phosphoric, monohydrogen phosphoric, dihydrogen phosphoric, phthalic, polygalactouronic, propionic, salicylic, stearic, succinic, sulfamic, sulfanilic, sulfonic, sulfuric, tannic, tartaric,
25 teoclic and toluenesulfonic.

31. A protected compound of the formula (II)



II

30

where R_1 is:

(I) C₁-C₆ alkyl, optionally substituted with one, two or three substituents selected from the group consisting of C₁-C₃ alkyl, C₁-C₇ alkyl (optionally substituted with C₁-C₃ alkyl and C₁-C₃ alkoxy), -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₃ alkoxy, -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are -H or C₁-C₆ alkyl, and -OC=O NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(II) -CH₂-S(O)₀₋₂-(C₁-C₆ alkyl),

(III) -CH₂-CH₂-S(O)₀₋₂-(C₁-C₆ alkyl),

(IV) C₂-C₆ alkenyl with one or two double bonds, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, -C≡N, -CF₃, C₁-C₃ alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are -H or C₁-C₆ alkyl,

(V) C₂-C₆ alkynyl with one or two triple bonds, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, -C≡N, -CF₃, C₁-C₃ alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are -H or C₁-C₆ alkyl,

(VI) -(CH₂)_{n1}-(R_{1-aryl}) where n₁ is zero or one and where R_{1-aryl} is phenyl, 1-naphthyl, 2-naphthyl and indanyl, indenyl, dihydronaphthalyl, or tetralinyl optionally substituted with one, two, three or four of the following substituents on the aryl ring:

(A) C₁-C₆ alkyl optionally substituted with one, two or three substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, and C₁-C₃ alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(B) C₂-C₆ alkenyl with one or two double bonds, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, -C≡N, -CF₃, C₁-C₃ alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are -H or C₁-C₆ alkyl,

(C) C₂-C₆ alkynyl with one or two triple bonds, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, -C≡N, -CF₃, C₁-C₃ alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are -H or C₁-C₆ alkyl,

(D) -F, Cl, -Br or -I,

(F) -C₁-C₆ alkoxy optionally substituted with one, two or three of -F,

(G) -NR_{N-2}R_{N-3} where R_{N-2} and R_{N-3} are as defined below,

(H) -OH,

5 (I) -C≡N,

(J) C₃-C₇ cycloalkyl, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, -C≡N, -CF₃, C₁-C₃ alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are -H or C₁-C₆ alkyl,

(K) -CO-(C₁-C₄ alkyl),

10 (L) -SO₂-NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(M) -CO-NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

or

(N) -SO₂-(C₁-C₄ alkyl),

(VII) -(CH₂)_{n₁}-(R_{1-heteroaryl}) where n₁ is as defined above and where

15 R_{1-heteroaryl} is selected from the group consisting of:

pyridinyl,

pyrimidinyl,

quinolinyl,

benzothienyl,

20 indolyl,

indolinyl,

pyridazinyl,

pyrazinyl,

isoquinolyl,

25 quinazolinyl,

quinoxalinyl,

phthalazinyl,

imidazolyl,

isoxazolyl,

30 pyrazolyl,

oxazolyl,

thiazolyl,

indolizinyl,

	indazolyl,
	benzothiazolyl,
	benzimidazolyl,
	benzofuranyl,
5	furanyl,
	thienyl,
	pyrrolyl,
	oxadiazolyl,
	thiadiazolyl,
10	triazolyl,
	tetrazolyl,
	oxazolopyridinyl,
	imidazopyridinyl,
	isothiazolyl,
15	naphthyridinyl,
	cinnolinyl,
	carbazolyl,
	beta-carbolinyl,
	isochromanyl,
20	chromanyl,
	tetrahydroisoquinolinyl,
	isoindolinyl,
	isobenzotetrahydrofuranyl,
	isobenzotetrahydrothienyl,
25	isobenzothieryl,
	benzoxazolyl,
	pyridopyridinyl,
	benzotetrahydrofuranyl,
	benzotetrahydrothienyl,
30	purinyl,
	benzodioxolyl,
	triazinyl,
	phenoxazinyl,
	phenothiazinyl,

	pteridinyl,
	benzothiazolyl,
	imidazopyridinyl,
	imidazothiazolyl,
5	dihydrobenzisoxazinyl,
	benzisoxazinyl,
	benzoxazinyl,
	dihydrobenzisoithiazinyl,
	benzopyranyl,
10	benzothiopyranyl,
	coumarinyl,
	isocoumarinyl,
	chromonyl,
	chromanonyl,
15	pyridinyl-N-oxide,
	tetrahydroquinolinyl
	dihydroquinolinyl
	dihydroquinolinonyl
	dihydroisoquinolinonyl
20	dihydrocoumarinyl
	dihydroisocoumarinyl
	isoindolinonyl
	benzodioxanyl
	benzoxazolinonyl
25	pyrrolyl N-oxide,
	pyrimidinyl N-oxide,
	pyridazinyl N-oxide,
	pyrazinyl N-oxide,
	quinolinyl N-oxide,
30	indolyl N-oxide,
	indolinyl N-oxide,
	isoquinolyl N-oxide,
	quinazolinyl N-oxide,
	quinoxalinyl N-oxide,

phthalaziny N-oxide,
 imidazolyl N-oxide,
 isoxazolyl N-oxide,
 oxazolyl N-oxide,
 5 thiazolyl N-oxide,
 indoliziny N-oxide,
 indazolyl N-oxide,
 benzothiazolyl N-oxide,
 benzimidazolyl N-oxide,
 10 pyrrolyl N-oxide,
 oxadiazolyl N-oxide,
 thiadiazolyl N-oxide,
 triazolyl N-oxide,
 tetrazolyl N-oxide,
 15 benzothiopyranyl S-oxide, and
 benzothiopyranyl S,S-dioxide,

where the $R_{1\text{-heteroaryl}}$ group is bonded to $-(CH_2)_{n1}-$ by any ring atom
 of the parent $R_{1\text{-heteroaryl}}$ group substituted by hydrogen such that the new bond to the
 $R_{1\text{-heteroaryl}}$ group replaces the hydrogen atom and its bond, where heteroaryl is
 20 optionally substituted with one, two, three or four of:

(1) C_1-C_6 alkyl optionally substituted with one, two or
 three substituents selected from the group consisting of C_1-C_3 alkyl, $-F$, $-Cl$, $-Br$, $-I$,
 $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, C_1-C_3 alkoxy, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as
 defined above,

25 (2) C_2-C_6 alkenyl with one or two double bonds,
 optionally substituted with one, two or three substituents selected from the group
 consisting of $-F$, $-Cl$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, C_1-C_3 alkoxy, and $-NR_{1-a}R_{1-b}$ where R_{1-}
 a and R_{1-b} are $-H$ or C_1-C_6 alkyl,

(3) C_2-C_6 alkynyl with one or two triple bonds,
 30 optionally substituted with one, two or three substituents selected from the group
 consisting of $-F$, $-Cl$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, C_1-C_3 alkoxy, and $-NR_{1-a}R_{1-b}$ where R_{1-}
 a and R_{1-b} are $-H$ or C_1-C_6 alkyl,

(4) $-F$, Cl , $-Br$ or $-I$,

(6) -C₁-C₆ alkoxy optionally substituted with one, two,
or three of -F,

(7) -NR_{N-2}R_{N-3} where R_{N-2} and R_{N-3} are as defined
below,

5 (8) -OH,

(9) -C≡N,

(10) C₃-C₇ cycloalkyl, optionally substituted with one,
two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, -
C≡N, -CF₃, C₁-C₃ alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are -H or C₁-C₆ alkyl,

10 (11) -CO-(C₁-C₄ alkyl),

(12) -SO₂-NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined
above,

(13) -CO-NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined
above, or

15 (14) -SO₂-(C₁-C₄ alkyl), with the proviso that when n₁
is zero R_{1-heteroaryl} is not bonded to the carbon chain by nitrogen; or

(VIII) -(CH₂)_{n1}-(R_{1-heterocycle}) where n₁ is as defined above and R<sub>1-
heterocycle</sub> is selected from the group consisting of:

20 morpholinyl,
thiomorpholinyl,
thiomorpholinyl S-oxide,
thiomorpholinyl S,S-dioxide,
piperazinyl,
homopiperazinyl,
25 pyrrolidinyl,
pyrrolinyl,
tetrahydropyranyl,
piperidinyl,
tetrahydrofuranyl,
30 tetrahydrothienyl,
homopiperidinyl,
homomorpholinyl,
homothiomorpholinyl,

homothiomorpholinyl S,S-dioxide,
oxazolidinonyl,
dihydropyrazolyl,
dihydropyrrolyl,
5 dihydropyrazinyl,
dihydropyridinyl,
dihydropyrimidinyl,
dihydrofuryl,
dihydropyranyl,
10 tetrahydrothienyl S-oxide,
tetrahydrothienyl S,S-dioxide, and
homothiomorpholinyl S-oxide,

where the $R_{1\text{-heterocycle}}$ group is bonded by any atom of the parent $R_{1\text{-heterocycle}}$ group substituted by hydrogen such that the new bond to the $R_{1\text{-heterocycle}}$ group replaces the hydrogen atom and its bond, where heterocycle is optionally substituted with one, two, three or four:

(1) $C_1\text{-}C_6$ alkyl optionally substituted with one, two or three substituents selected from the group consisting of $C_1\text{-}C_3$ alkyl, -F, -Cl, -Br, -I, -OH, -SH, $-C\equiv N$, $-CF_3$, $C_1\text{-}C_3$ alkoxy, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

(2) $C_2\text{-}C_6$ alkenyl with one or two double bonds, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, $-C\equiv N$, $-CF_3$, $C_1\text{-}C_3$ alkoxy, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are -H or $C_1\text{-}C_6$ alkyl,

(3) $C_2\text{-}C_6$ alkynyl with one or two triple bonds, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, $-C\equiv N$, $-CF_3$, $C_1\text{-}C_3$ alkoxy, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are -H or $C_1\text{-}C_6$ alkyl,

(4) -F, Cl, -Br or -I,

(5) $C_1\text{-}C_6$ alkoxy,

(6) $-C_1\text{-}C_6$ alkoxy optionally substituted with one, two, or three -F,

(7) $-NR_{N-2}R_{N-3}$ where R_{N-2} and R_{N-3} are as defined

below,

(8) $-OH$,

(9) $-C\equiv N$,

5 (10) C_3-C_7 cycloalkyl, optionally substituted with one, two or three substituents selected from the group consisting of $-F$, $-Cl$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, C_1-C_3 alkoxy, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are $-H$ or C_1-C_6 alkyl,

(11) $-CO-(C_1-C_4 \text{ alkyl})$,

(12) $-SO_2-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined

10 above,

(13) $-CO-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined

above,

(14) $-SO_2-(C_1-C_4 \text{ alkyl})$, or

(15) $=O$, with the proviso that when n_1 is zero R_1 .

15 heterocycle is not bonded to the carbon chain by nitrogen;

where R_2 is:

(I) $-H$,

(II) C_1-C_6 alkyl, optionally substituted with one, two or three
20 substituents selected from the group consisting of C_1-C_3 alkyl, $-F$, $-Cl$, $-Br$, $-I$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, C_1-C_3 alkoxy, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

(III) $-(CH_2)_{0-4}-R_{2-1}$ where R_{2-1} is R_{1-aryl} or $R_{1-heteroaryl}$ where R_{1-aryl} and $R_{1-heteroaryl}$ are as defined above;

25 (IV) C_2-C_6 alkenyl with one or two double bonds, optionally substituted with one, two or three substituents selected from the group consisting of $-F$, $-Cl$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, C_1-C_3 alkoxy, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are $-H$ or C_1-C_6 alkyl,

(V) C_2-C_6 alkynyl with one or two triple bonds, optionally substituted
30 with one, two or three substituents selected from the group consisting of $-F$, $-Cl$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, C_1-C_3 alkoxy, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are $-H$ or C_1-C_6 alkyl, or

(VI) $-(CH_2)_{0-4}-C_3-C_7$ cycloalkyl, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, $-C\equiv N$, $-CF_3$, C_1-C_3 alkoxy, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are -H or C_1-C_6 alkyl;

5 where R_3 is:

(I)-H,

(II) C_1-C_6 alkyl, optionally substituted with one, two or three substituents selected from the group consisting of C_1-C_3 alkyl, -F, -Cl, -Br, -I, -OH, -SH, $-C\equiv N$, $-CF_3$, C_1-C_3 alkoxy, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined

10 above,

(III) $-(CH_2)_{0-4}-R_{2-1}$ where R_{2-1} is R_{1-aryl} or $R_{1-heteroaryl}$ where R_{1-aryl} and $R_{1-heteroaryl}$ are as defined above

(IV) C_2-C_6 alkenyl with one or two double bonds,

(V) C_2-C_6 alkynyl with one or two triple bonds; or

15 (VI) $-(CH_2)_{0-4}-C_3-C_7$ cycloalkyl, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, $-C\equiv N$, $-CF_3$, C_1-C_3 alkoxy, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are -H or C_1-C_6 alkyl,

and where R_2 and R_3 are taken together with the carbon to which they are attached to form a carbocycle of three, four, five, six, or seven carbon atoms,

20 optionally where one carbon atom is replaced by a heteroatom selected from the group consisting of -O-, -S-, $-SO_2-$, and $-NR_{N-2}-$, where R_{N-2} is selected from the group consisting of:

(a) -H,

(b) C_1-C_6 alkyl optionally substituted with one

25 substituent selected from the group consisting of:

(i) -OH, and

(ii) $-NH_2$,

(c) C_1-C_6 alkyl optionally substituted with

one, two, or three -F, -Cl, -Br, or -I,

30

(d) C_3-C_7 cycloalkyl,

(e) $-(C_1-C_2 \text{ alkyl})-(C_3-C_7 \text{ cycloalkyl})$,

(f) $-(C_1-C_6 \text{ alkyl})-O-(C_1-C_3 \text{ alkyl})$,

(g) $-C_2-C_6$ alkenyl with one or two double bonds,

(h) $-C_2-C_6$ alkynyl with one or two triple bonds,

5 (i) $-C_1-C_6$ alkyl chain with one double bond and one triple bond,

(j) $-R_{1-aryl}$ where R_{1-aryl} is as defined above, and

(k) $-R_{1-heteroaryl}$ where $R_{1-heteroaryl}$ is as defined above;

10

where R_N is:

(I) $R_{N-1}-X_N$ where X_N is selected from the group consisting of:

(A) $-CO-$,

(B) $-SO_2-$,

15 (C) $-(CR'R'')_{1-6}$ where R' and R'' are the same or different and are $-H$ and C_1-C_4 alkyl,

(D) $-CO-(CR'R'')_{1-6}-X_{N-1}$ where X_{N-1} is selected from the group consisting of $-O-$, $-S-$ and $-NR'-$ and where R' and R'' are as defined above, and

20 (E) a single bond;

where R_{N-1} is selected from the group consisting of:

(A) R_{N-aryl} where R_{N-aryl} is phenyl, 1-naphthyl, 2-naphthyl, tetralinyl, indanyl, 6,7,8,9-tetrahydro-5H-benzo[a]cycloheptenyl, or dihydronaphthyl optionally substituted with one, two or three of the following substituents which can
25 be the same or different and are:

(1) C_1-C_6 alkyl, optionally substituted with one, two or three substituents selected from the group consisting of C_1-C_3 alkyl, $-F$, $-Cl$, $-Br$, $-I$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, C_1-C_3 alkoxy, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

30

(2) $-OH$,

(3) $-NO_2$,

(4) $-F$, $-Cl$, $-Br$, $-I$,

(5) $-CO-OH$,

(6) $-\text{C}\equiv\text{N}$,

(7) $-(\text{CH}_2)_{0-4}-\text{CO}-\text{NR}_{\text{N}-2}\text{R}_{\text{N}-3}$ where $\text{R}_{\text{N}-2}$ and $\text{R}_{\text{N}-3}$ are the same or different and are selected from the group consisting of:

(a) $-\text{H}$,

5 (b) $-\text{C}_1-\text{C}_6$ alkyl optionally substituted with one substituent selected from the group consisting of:

(i) $-\text{OH}$, and

(ii) $-\text{NH}_2$,

10 (c) $-\text{C}_1-\text{C}_6$ alkyl optionally substituted with one, two, or three $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, or $-\text{I}$,

(d) $-\text{C}_3-\text{C}_7$ cycloalkyl,

(e) $-(\text{C}_1-\text{C}_2 \text{ alkyl})-(\text{C}_3-\text{C}_7 \text{ cycloalkyl})$,

(f) $-(\text{C}_1-\text{C}_6 \text{ alkyl})-\text{O}-(\text{C}_1-\text{C}_3 \text{ alkyl})$,

15 (g) $-\text{C}_2-\text{C}_6$ alkenyl with one or two double bonds,

(h) $-\text{C}_2-\text{C}_6$ alkynyl with one or two triple

bonds,

(i) $-\text{C}_1-\text{C}_6$ alkyl chain with one double bond

and one triple bond,

20 (j) $-\text{R}_1\text{-aryl}$ where $\text{R}_1\text{-aryl}$ is as defined above, and

(k) $-\text{R}_1\text{-heteroaryl}$ where $\text{R}_1\text{-heteroaryl}$ is as defined

above,

(8) $-(\text{CH}_2)_{0-4}-\text{CO}-(\text{C}_1-\text{C}_{12} \text{ alkyl})$,

25 (9) $-(\text{CH}_2)_{0-4}-\text{CO}-(\text{C}_2-\text{C}_{12} \text{ alkenyl with one, two or three double bonds})$,

(10) $-(\text{CH}_2)_{0-4}-\text{CO}-(\text{C}_2-\text{C}_{12} \text{ alkynyl with one, two or$

three triple bonds),

(11) $-(\text{CH}_2)_{0-4}-\text{CO}-(\text{C}_3-\text{C}_7 \text{ cycloalkyl})$,

(12) $-(\text{CH}_2)_{0-4}-\text{CO}-\text{R}_1\text{-aryl}$ where $\text{R}_1\text{-aryl}$ is as defined

30 above,

(13) $-(\text{CH}_2)_{0-4}-\text{CO}-\text{R}_1\text{-heteroaryl}$ where $\text{R}_1\text{-heteroaryl}$ is as

defined above,

(14) $-(CH_2)_{0-4}-CO-R_{1\text{-heterocycle}}$ where $R_{1\text{-heterocycle}}$ is as defined above,

(15) $-(CH_2)_{0-4}-CO-R_{N-4}$ where R_{N-4} is selected from the group consisting of morpholinyl, thiomorpholinyl, piperazinyl, piperidinyl, homomorpholinyl, homothiomorpholinyl, homothiomorpholinyl S-oxide, homothiomorpholinyl S,S-dioxide, pyrrolinyl and pyrrolidinyl where each group is optionally substituted with one, two, three, or four of C_1-C_6 alkyl,

(16) $-(CH_2)_{0-4}-CO-O-R_{N-5}$ where R_{N-5} is selected from the group consisting of:

- (a) C_1-C_6 alkyl,
 (b) $-(CH_2)_{0-2}-(R_{1\text{-aryl}})$ where $R_{1\text{-aryl}}$ is as defined above,
 (c) C_2-C_6 alkenyl containing one or two double bonds,
 (d) C_2-C_6 alkynyl containing one or two triple bonds,
 (e) C_3-C_7 cycloalkyl, and
 (f) $-(CH_2)_{0-2}-(R_{1\text{-heteroaryl}})$ where $R_{1\text{-heteroaryl}}$ is as defined above,

(17) $-(CH_2)_{0-4}-SO_2-NR_{N-2}R_{N-3}$ where R_{N-2} and R_{N-3} are as defined above,

(18) $-(CH_2)_{0-4}-SO-(C_1-C_8 \text{ alkyl})$,

(19) $-(CH_2)_{0-4}-SO_2-(C_1-C_{12} \text{ alkyl})$,

(20) $-(CH_2)_{0-4}-SO_2-(C_3-C_7 \text{ cycloalkyl})$,

(21) $-(CH_2)_{0-4}-N(H \text{ or } R_{N-5})-CO-O-R_{N-5}$ where R_{N-5} can be the same or different and is as defined above,

(22) $-(CH_2)_{0-4}-N(H \text{ or } R_{N-5})-CO-N(R_{N-5})_2$, where R_{N-5} can be the same or different and is as defined above,

(23) $-(CH_2)_{0-4}-N-CS-N(R_{N-5})_2$, where R_{N-5} can be the same or different and is as defined above,

(24) $-(CH_2)_{0-4}-N(-H \text{ or } R_{N-5})-CO-R_{N-2}$ where R_{N-5} and R_{N-2} can be the same or different and are as defined above,

(25) $-(CH_2)_{0-4}-NR_{N-2}R_{N-3}$ where R_{N-2} and R_{N-3} can be the same or different and are as defined above,

- (26) $-(\text{CH}_2)_{0-4}-\text{R}_{\text{N}-4}$ where $\text{R}_{\text{N}-4}$ is as defined above,
- (27) $-(\text{CH}_2)_{0-4}-\text{O}-\text{CO}-(\text{C}_1-\text{C}_6 \text{ alkyl})$,
- (28) $-(\text{CH}_2)_{0-4}-\text{O}-\text{P}(\text{O})-(\text{OR}_{\text{N-aryl-1}})_2$ where $\text{R}_{\text{N-aryl-1}}$ is –
 H or C_1-C_4 alkyl,
- 5 (29) $-(\text{CH}_2)_{0-4}-\text{O}-\text{CO}-\text{N}(\text{R}_{\text{N-5}})_2$ where $\text{R}_{\text{N-5}}$ is as
 defined above,
- (30) $-(\text{CH}_2)_{0-4}-\text{O}-\text{CS}-\text{N}(\text{R}_{\text{N-5}})_2$ where $\text{R}_{\text{N-5}}$ is as defined
 above,
- (31) $-(\text{CH}_2)_{0-4}-\text{O}-(\text{R}_{\text{N-5}})_2$ where $\text{R}_{\text{N-5}}$ is as defined
 10 above,
- (32) $-(\text{CH}_2)_{0-4}-\text{O}-(\text{R}_{\text{N-5}})_2-\text{COOH}$ where $\text{R}_{\text{N-5}}$ is as
 defined above,
- (33) $-(\text{CH}_2)_{0-4}-\text{S}-(\text{R}_{\text{N-5}})_2$ where $\text{R}_{\text{N-5}}$ is as defined
 above,
- 15 (34) $-(\text{CH}_2)_{0-4}-\text{O}-(\text{C}_1-\text{C}_6 \text{ alkyl optionally substituted}$
 with one, two, three, four, or five of $-\text{F}$),
- (35) C_3-C_7 cycloalkyl,
- (36) C_2-C_6 alkenyl with one or two double bonds
 optionally substituted with C_1-C_3 alkyl, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, $-\text{OH}$, $-\text{SH}$, $-\text{C}\equiv\text{N}$, $-\text{CF}_3$, C_1-
 20 C_3 alkoxy, or $-\text{NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,
- (37) C_2-C_6 alkynyl with one or two triple bonds
 optionally substituted with C_1-C_3 alkyl, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, $-\text{OH}$, $-\text{SH}$, $-\text{C}\equiv\text{N}$, $-\text{CF}_3$, C_1-
 C_3 alkoxy, or $-\text{NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,
- (38) $-(\text{CH}_2)_{0-4}-\text{N}(-\text{H} \text{ or } \text{R}_{\text{N-5}})-\text{SO}_2-\text{R}_{\text{N-2}}$ where $\text{R}_{\text{N-5}}$ and
 25 $\text{R}_{\text{N-2}}$ can be the same or different and are as described above, or
- (39) $-(\text{CH}_2)_{0-4}-\text{C}_3-\text{C}_7$ cycloalkyl,
- (B) $-\text{R}_{\text{N-heteroaryl}}$ where $\text{R}_{\text{N-heteroaryl}}$ is selected from the group
 consisting of:
- 30 pyridinyl,
 pyrimidinyl,
 quinolinyl,
 benzothienyl,
 indolyl,

	indolinyI,
	pyridazinyI,
	pyrazinyI,
	isoindolyI,
5	isoquinolyI,
	quinazolinyI,
	quinoxalinyI,
	phthalazinyI,
	imidazolyl,
10	isoxazolyl,
	pyrazolyl,
	oxazolyl,
	thiazolyl,
	indolizinyI,
15	indazolyl,
	benzothiazolyl,
	benzimidazolyl,
	benzofuranyl,
	furanyl,
20	thienyl,
	pyrrolyl,
	oxadiazolyl,
	thiadiazolyl,
	triazolyl,
25	tetrazolyl,
	oxazolopyridinyI,
	imidazopyridinyI,
	isothiazolyl,
	naphthyridinyI,
30	cinnolinyI,
	carbazolyl,
	beta-carbolinyI,
	isochromanyI,
	chromanyI,

	tetrahydroisoquinolinyI,
	isoindolinyI,
	isobenzotetrahydrofuranyI,
	isobenzotetrahydrothienyI,
5	isobenzothiényI,
	benzoxazolyl,
	pyridopyridinyI,
	benzotetrahydrofuranyI,
	benzotetrahydrothienyI,
10	purinyI,
	benzodioxolyl,
	triazinyI,
	phenoxazinyI,
	phenothiazinyI,
15	pteridinyI,
	benzothiazolyl,
	imidazopyridinyI,
	imidazothiazolyl,
	dihydrobenzisoXazinyI,
20	benzisoXazinyI,
	benzoxazinyI,
	dihydrobenzisothiazinyI,
	benzopyranyI,
	benzothiopyranyI,
25	coumarinyI,
	isocoumarinyI,
	chromonyI,
	chromanonyI,
	pyridinyI-N-oxide,
30	tetrahydroquinolinyI,
	dihydroquinolinyI,
	dihydroquinolinonyI,
	dihydroisoquinolinonyI,
	dihydrocoumarinyI,

	dihydroisocoumarinyl,
	isoindolinonyl,
	benzodioxanyl,
	benzoxazolinonyl,
5	pyrrolyl N-oxide,
	pyrimidinyl N-oxide,
	pyridazinyl N-oxide,
	pyrazinyl N-oxide,
	quinolinyl N-oxide,
10	indolyl N-oxide,
	indolinyl N-oxide,
	isoquinolyl N-oxide,
	quinazolinyl N-oxide,
	quinoxalinyl N-oxide,
15	phthalazinyl N-oxide,
	imidazolyl N-oxide,
	isoxazolyl N-oxide,
	oxazolyl N-oxide,
	thiazolyl N-oxide,
20	indoliziny N-oxide,
	indazolyl N-oxide,
	benzothiazolyl N-oxide,
	benzimidazolyl N-oxide,
	pyrrolyl N-oxide,
25	oxadiazolyl N-oxide,
	thiadiazolyl N-oxide,
	triazolyl N-oxide,
	tetrazolyl N-oxide,
	benzothiopyranyl S-oxide, and
30	benzothiopyranyl S,S-dioxide

where the $R_{N\text{-heteroaryl}}$ group is bonded by any atom of the parent $R_{N\text{-heteroaryl}}$ group substituted by hydrogen such that the new bond to the $R_{N\text{-heteroaryl}}$ group replaces the hydrogen atom and its bond, where heteroaryl is optionally substituted with one, two, three, or four of:

- (1) C₁-C₆ alkyl, optionally substituted with one, two or three substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₃ alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,
- 5 (2) -OH,
 (3) -NO₂,
 (4) -F, -Cl, -Br, or -I,
 (5) -CO-OH,
 (6) -C≡N,
- 10 (7) -(CH₂)₀₋₄-CO-NR_{N-2}R_{N-3} where R_{N-2} and R_{N-3} are the same or different and are selected from the group consisting of:
- (a) -H,
 (b) -C₁-C₆ alkyl optionally substituted with one substituent selected from the group consisting of:
- 15 (i) -OH, and
 (ii) -NH₂,
 (c) -C₁-C₆ alkyl optionally substituted with one, two, or three -F, -Cl, -Br, -I,
 (d) -C₃-C₇ cycloalkyl,
 (e) -(C₁-C₂ alkyl)-(C₃-C₇ cycloalkyl),
 (f) -(C₁-C₆ alkyl)-O-(C₁-C₃ alkyl),
 (g) -C₂-C₆ alkenyl with one or two double bonds,
 (h) -C₂-C₆ alkynyl with one or two triple bonds,
 (i) -C₁-C₆ alkyl chain with one double bond and one triple bond,
 (j) -R_{1-aryl} where R_{1-aryl} is as defined above,
 (k) -R_{1-heteroaryl} where R_{1-heteroaryl} is as defined above,
- 20
 25
 30
- (8) -(CH₂)₀₋₄-CO-(C₁-C₁₂ alkyl),
 (9) -(CH₂)₀₋₄-CO-(C₂-C₁₂ alkenyl with one, two or three double bonds),

(10) $-(\text{CH}_2)_{0-4}-\text{CO}-(\text{C}_2-\text{C}_{12}$ alkynyl with one, two or three triple bonds),

(11) $-(\text{CH}_2)_{0-4}-\text{CO}-(\text{C}_3-\text{C}_7$ cycloalkyl),

(12) $-(\text{CH}_2)_{0-4}-\text{CO}-\text{R}_{1-\text{aryl}}$ where $\text{R}_{1-\text{aryl}}$ is as defined

5 above,

(13) $-(\text{CH}_2)_{0-4}-\text{CO}-\text{R}_{1-\text{heteroaryl}}$ where $\text{R}_{1-\text{heteroaryl}}$ is as

defined above,

(14) $-(\text{CH}_2)_{0-4}-\text{CO}-\text{R}_{1-\text{heterocycle}}$ where $\text{R}_{1-\text{heterocycle}}$ is as

defined above,

10 (15) $-(\text{CH}_2)_{0-4}-\text{CO}-\text{R}_{\text{N-4}}$ where $\text{R}_{\text{N-4}}$ is selected from the group consisting of morpholinyl, thiomorpholinyl, piperazinyl, piperidinyl, homomorpholinyl, homothiomorpholinyl, homothiomorpholinyl S-oxide, homothiomorpholinyl S,S-dioxide, pyrrolinyl and pyrrolidinyl where each group is optionally substituted with one, two, three, or four of C_1-C_6 alkyl,

15 (16) $-(\text{CH}_2)_{0-4}-\text{CO}-\text{O}-\text{R}_{\text{N-5}}$ where $\text{R}_{\text{N-5}}$ is selected from the group consisting of:

(a) C_1-C_6 alkyl,

(b) $-(\text{CH}_2)_{0-2}-(\text{R}_{1-\text{aryl}})$ where $\text{R}_{1-\text{aryl}}$ is as defined

above,

20 (c) C_2-C_6 alkenyl containing one or two double bonds,

(d) C_2-C_6 alkynyl containing one or two triple

bonds,

(e) C_3-C_7 cycloalkyl, and

25 (f) $-(\text{CH}_2)_{0-2}-(\text{R}_{1-\text{heteroaryl}})$ where $\text{R}_{1-\text{heteroaryl}}$ is as defined above,

(17) $-(\text{CH}_2)_{0-4}-\text{SO}_2-\text{NR}_{\text{N-2}}\text{R}_{\text{N-3}}$ where $\text{R}_{\text{N-2}}$ and $\text{R}_{\text{N-3}}$ are as defined above,

(18) $-(\text{CH}_2)_{0-4}-\text{SO}-(\text{C}_1-\text{C}_8$ alkyl),

30 (19) $-(\text{CH}_2)_{0-4}-\text{SO}_2-(\text{C}_1-\text{C}_{12}$ alkyl),

(20) $-(\text{CH}_2)_{0-4}-\text{SO}_2-(\text{C}_3-\text{C}_7$ cycloalkyl),

(21) $-(\text{CH}_2)_{0-4}-\text{N}(\text{H or } \text{R}_{\text{N-5}})-\text{CO}-\text{O}-\text{R}_{\text{N-5}}$ where $\text{R}_{\text{N-5}}$

can be the same or different and is as defined above,

- (22) $-(\text{CH}_2)_{0-4}-\text{N}(\text{H or } \text{R}_{\text{N-5}})-\text{CO}-\text{N}(\text{R}_{\text{N-5}})_2$, where $\text{R}_{\text{N-5}}$ can be the same or different and is as defined above,
- (23) $-(\text{CH}_2)_{0-4}-\text{N}-\text{CS}-\text{N}(\text{R}_{\text{N-5}})_2$, where $\text{R}_{\text{N-5}}$ can be the same or different and is as defined above,
- 5 (24) $-(\text{CH}_2)_{0-4}-\text{N}(\text{H or } \text{R}_{\text{N-5}})-\text{CO}-\text{R}_{\text{N-2}}$ where $\text{R}_{\text{N-5}}$ and $\text{R}_{\text{N-2}}$ can be the same or different and are as defined above,
- (25) $-(\text{CH}_2)_{0-4}-\text{NR}_{\text{N-2}}\text{R}_{\text{N-3}}$ where $\text{R}_{\text{N-2}}$ and $\text{R}_{\text{N-3}}$ can be the same or different and are as defined above,
- (26) $-(\text{CH}_2)_{0-4}-\text{R}_{\text{N-4}}$ where $\text{R}_{\text{N-4}}$ is as defined above,
- 10 (27) $-(\text{CH}_2)_{0-4}-\text{O}-\text{CO}-(\text{C}_1-\text{C}_6 \text{ alkyl})$,
- (28) $-(\text{CH}_2)_{0-4}-\text{O}-\text{P}(\text{O})-(\text{OR}_{\text{N-aryl-1}})_2$ where $\text{R}_{\text{N-aryl-1}}$ is $-\text{H}$ or C_1-C_4 alkyl,
- (29) $-(\text{CH}_2)_{0-4}-\text{O}-\text{CO}-\text{N}(\text{R}_{\text{N-5}})_2$ where $\text{R}_{\text{N-5}}$ is as defined above,
- 15 (30) $-(\text{CH}_2)_{0-4}-\text{O}-\text{CS}-\text{N}(\text{R}_{\text{N-5}})_2$ where $\text{R}_{\text{N-5}}$ is as defined above,
- (31) $-(\text{CH}_2)_{0-4}-\text{O}-(\text{R}_{\text{N-5}})_2$ where $\text{R}_{\text{N-5}}$ is as defined above,
- (32) $-(\text{CH}_2)_{0-4}-\text{O}-(\text{R}_{\text{N-5}})_2-\text{COOH}$ where $\text{R}_{\text{N-5}}$ is as defined above,
- 20 (33) $-(\text{CH}_2)_{0-4}-\text{S}-(\text{R}_{\text{N-5}})_2$ where $\text{R}_{\text{N-5}}$ is as defined above,
- (34) $-(\text{CH}_2)_{0-4}-\text{O}-(\text{C}_1-\text{C}_6 \text{ alkyl optionally substituted with one, two, three, four, or five of } -\text{F})$,
- 25 (35) C_3-C_7 cycloalkyl,
- (36) C_2-C_6 alkenyl with one or two double bonds optionally substituted with C_1-C_3 alkyl, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, $-\text{OH}$, $-\text{SH}$, $-\text{C}\equiv\text{N}$, $-\text{CF}_3$, C_1-C_3 alkoxy, or $-\text{NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,
- (37) C_2-C_6 alkynyl with one or two triple bonds optionally substituted with C_1-C_3 alkyl, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, $-\text{OH}$, $-\text{SH}$, $-\text{C}\equiv\text{N}$, $-\text{CF}_3$, C_1-C_3 alkoxy, or $-\text{NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,
- 30 (38) $-(\text{CH}_2)_{0-4}-\text{N}(\text{H or } \text{R}_{\text{N-5}})-\text{SO}_2-\text{R}_{\text{N-2}}$ where $\text{R}_{\text{N-5}}$ and $\text{R}_{\text{N-2}}$ can be the same or different and are as defined above, or

(39) $-(\text{CH}_2)_{0-4}-\text{C}_3-\text{C}_7$ cycloalkyl,

(C) $\text{R}_{\text{N-aryl}}-\text{W}-\text{R}_{\text{N-aryl}}$, where $\text{R}_{\text{N-aryl}}$ can be the same or different,

(D) $\text{R}_{\text{N-aryl}}-\text{W}-\text{R}_{\text{N-heteroaryl}}$,

5 (E) $\text{R}_{\text{N-aryl}}-\text{W}-\text{R}_{\text{N-1-heterocycle}}$, wherein $\text{R}_{\text{N-1-heterocycle}}$ is the same as $\text{R}_{1-\text{heterocycle}}$, and $\text{R}_{1-\text{heterocycle}}$ is as defined above

(F) $\text{R}_{\text{N-heteroaryl}}-\text{W}-\text{R}_{\text{N-aryl}}$,

(G) $\text{R}_{\text{N-heteroaryl}}-\text{W}-\text{R}_{\text{N-heteroaryl}}$,

(H) $\text{R}_{\text{N-heteroaryl}}-\text{W}-\text{R}_{\text{N-1-heterocycle}}$,

10 (I) $\text{R}_{\text{N-heterocycle}}-\text{W}-\text{R}_{\text{N-aryl}}$, wherein $\text{R}_{\text{N-heterocycle}}$ is the same as $\text{R}_{1-\text{heterocycle}}$, and $\text{R}_{1-\text{heterocycle}}$ is as defined above, and $\text{R}_{\text{N-aryl}}$ is as defined above,

(J) $\text{R}_{\text{N-heterocycle}}-\text{W}-\text{R}_{\text{N-heteroaryl}}$, and

(K) $\text{R}_{\text{N-heterocycle}}-\text{W}-\text{R}_{\text{N-1-heterocycle}}$,

where W is

15 (5) $-(\text{CH}_2)_{0-4}-$,

(6) $-\text{O}-$,

(7) $-\text{S}(\text{O})_{0-2}-$,

(8) $-\text{N}(\text{R}_{\text{N-5}})-$ where $\text{R}_{\text{N-5}}$ is as defined above, or

(5) $-\text{CO}-$;

20 (II) $-\text{CO}-(\text{C}_1-\text{C}_{10} \text{ alkyl})$ where alkyl is optionally substituted with one three substituents selected from the group consisting of:

(A) $-\text{OH}$,

(B) $-\text{C}_1-\text{C}_6$ alkoxy,

(C) $-\text{C}_1-\text{C}_6$ thioalkoxy,

25 (D) $-\text{CO}-\text{O}-\text{R}_{\text{N-8}}$ where $\text{R}_{\text{N-8}}$ is $-\text{H}$, C_1-C_6 alkyl or $-\text{phenyl}$,

(E) $-\text{CO}-\text{NR}_{\text{N-2}}\text{R}_{\text{N-3}}$ where $\text{R}_{\text{N-2}}$ and $\text{R}_{\text{N-3}}$ are the same or

different and are as defined above,

(F) $-\text{CO}-\text{R}_{\text{N-4}}$ where $\text{R}_{\text{N-4}}$ is as defined above,

(G) $-\text{SO}_2-(\text{C}_1-\text{C}_8 \text{ alkyl})$,

30 (H) $-\text{SO}_2-\text{NR}_{\text{N-2}}\text{R}_{\text{N-3}}$ where $\text{R}_{\text{N-2}}$ and $\text{R}_{\text{N-3}}$ are the same or different and are as defined above,

(I) $-\text{NH}-\text{CO}-(\text{C}_1-\text{C}_6 \text{ alkyl})$,

(J) $-\text{NH}-\text{CO}-\text{O}-\text{R}_{\text{N-8}}$ where $\text{R}_{\text{N-8}}$ is as defined above,

(K) $-NR_{N-2}R_{N-3}$ where R_{N-2} and R_{N-3} are the same or different and are as defined above,

(L) $-R_{N-4}$ where R_{N-4} is as defined above,

(M) $-O-CO-(C_1-C_6 \text{ alkyl})$,

5 (N) $-O-CO-NR_{N-8}R_{N-8}$ where R_{N-8} are the same or different and are as defined above,

(O) $-O-(C_1-C_5 \text{ alkyl})-COOH$,

(P) $-O-(C_1-C_6 \text{ alkyl})$ optionally substituted with one, two, or three of $-F$, $-Cl$, $-Br$, or $-I$,

10 (Q) $-NH-SO_2-(C_1-C_6 \text{ alkyl})$, and

(R) $-F$, or $-Cl$,

(III) $-CO-(C_1-C_6 \text{ alkyl})-O-(C_1-C_6 \text{ alkyl})$ where alkyl is optionally substituted with one, two, or three substituents selected from the group consisting of:

15 (A) $-OH$,

(B) $-C_1-C_6 \text{ alkoxy}$,

(C) $-C_1-C_6 \text{ thioalkoxy}$,

(D) $-CO-O-R_{N-8}$ where R_{N-8} is $-H$, $C_1-C_6 \text{ alkyl}$ or $-phenyl$,

(E) $-CO-NR_{N-2}R_{N-3}$ where R_{N-2} and R_{N-3} are the same or
20 different and are as defined above,

(F) $-CO-R_{N-4}$ where R_{N-4} is as defined above,

(G) $-SO_2-(C_1-C_8 \text{ alkyl})$,

(H) $-SO_2-NR_{N-2}R_{N-3}$ where R_{N-2} and R_{N-3} are the same or different and are as defined above,

25 (I) $-NH-CO-(C_1-C_6 \text{ alkyl})$,

(J) $-NH-CO-O-R_{N-8}$ where R_{N-8} is as defined above,

(K) $-NR_{N-2}R_{N-3}$ where R_{N-2} and R_{N-3} are the same or different and are as defined above,

(L) $-R_{N-4}$ where R_{N-4} is as defined above,

30 (M) $-O-CO-(C_1-C_6 \text{ alkyl})$,

(N) $-O-CO-NR_{N-8}R_{N-8}$ where R_{N-8} are the same or different and are as defined above,

(O) $-O-(C_1-C_5 \text{ alkyl})-COOH$,

(P) $-O-(C_1-C_6 \text{ alkyl optionally substituted with one, two, or three of } -F, -Cl, -Br, \text{ or } -I),$

(Q) $-NH-SO_2-(C_1-C_6 \text{ alkyl}),$ and

(R) $-F, \text{ or } -Cl,$

5 (IV) $-CO-(C_1-C_6 \text{ alkyl})-S-(C_1-C_6 \text{ alkyl})$ where alkyl is optionally substituted with one, two, or three of substituents selected from the group consisting of:

(A) $-OH,$

(B) $-C_1-C_6 \text{ alkoxy},$

10 (C) $-C_1-C_6 \text{ thioalkoxy},$

(D) $-CO-O-R_{N-8}$ where R_{N-8} is as defined above,

(E) $-CO-NR_{N-2}R_{N-3}$ where R_{N-2} and R_{N-3} are the same or different and are as defined above,

(F) $-CO-R_{N-4}$ where R_{N-4} is as defined above,

15 (G) $-SO_2-(C_1-C_8 \text{ alkyl}),$

(H) $-SO_2-NR_{N-2}R_{N-3}$ where R_{N-2} and R_{N-3} are the same or different and are as defined above,

(I) $-NH-CO-(C_1-C_6 \text{ alkyl}),$

(J) $-NH-CO-O-R_{N-8}$ where R_{N-8} is as defined above,

20 (K) $-NR_{N-2}R_{N-3}$ where R_{N-2} and R_{N-3} are the same or different and are as defined above,

(L) $-R_{N-4}$ where R_{N-4} is as defined above,

(M) $-O-CO-(C_1-C_6 \text{ alkyl}),$

(N) $-O-CO-NR_{N-8}R_{N-8}$ where R_{N-8} are the same or different

25 and are as defined above,

(O) $-O-(C_1-C_5 \text{ alkyl})-COOH,$

(P) $-O-(C_1-C_6 \text{ alkyl optionally substituted with one, two, or three of } -F, -Cl, -Br, \text{ or } -I),$

(Q) $-NH-SO_2-(C_1-C_6 \text{ alkyl}),$ and

30 (R) $-F, \text{ or } -Cl,$

(V) $-CO-CH(-(CH_2)_{0-2}-O-R_{N-10})-(CH_2)_{0-2}-R_{N-aryl}/R_{N-heteroaryl})$ where R_{N-aryl} and $R_{N-heteroaryl}$ are as defined above, where R_{N-10} is selected from the group consisting of:

- (A) -H,
 (B) C₁-C₆ alkyl,
 (C) C₃-C₇ cycloalkyl,
 (D) C₂-C₆ alkenyl with one double bond,
 5 (E) C₂-C₆ alkynyl with one triple bond,
 (F) R_{1-aryl} where R_{1-aryl} is as defined above, and
 (G) R_{N-heteroaryl} where R_{N-heteroaryl} is as defined above, or
 (VI) -CO-(C₃-C₈ cycloalkyl) where alkyl is optionally substituted
 with one or two substituents selected from the group consisting of:
- 10 (A) -(CH₂)₀₋₄-OH,
 (B) -(CH₂)₀₋₄-C₁-C₆ alkoxy,
 (C) -(CH₂)₀₋₄-C₁-C₆ thioalkoxy,
 (D) -(CH₂)₀₋₄-CO-O-R_{N-8} where R_{N-8} is -H, C₁-C₆ alkyl or
 phenyl,
 15 (E) -(CH₂)₀₋₄-CO-NR_{N-2}R_{N-3} where R_{N-2} and R_{N-3} are the
 same or different and are as defined above,
 (F) -(CH₂)₀₋₄-CO-R_{N-4} where R_{N-4} is as defined above,
 (G) -(CH₂)₀₋₄-SO₂-(C₁-C₈ alkyl),
 (H) -(CH₂)₀₋₄-SO₂-NR_{N-2}R_{N-3} where R_{N-2} and R_{N-3} are the
 20 same or different and are as defined above,
 (I) -(CH₂)₀₋₄-NH-CO-(C₁-C₆ alkyl),
 (J) -NH-CO-O-R_{N-8} where R_{N-8} is as defined above,
 (K) -(CH₂)₀₋₄-NR_{N-2}R_{N-3} where R_{N-2} and R_{N-3} are the same or
 different and are as defined above,
 25 (L) -(CH₂)₀₋₄-R_{N-4} where R_{N-4} is as defined above,
 (M) -O-CO-(C₁-C₆ alkyl),
 (N) -O-CO-NR_{N-8}R_{N-8} where R_{N-8} are the same or different
 and are as defined above,
 (O) -O-(C₁-C₅ alkyl)-COOH,
 30 (P) -O-(C₁-C₆ alkyl optionally substituted with one, two, or
 three of -F, -Cl, -Br, or -I),
 (Q) -NH-SO₂-(C₁-C₆ alkyl), and

(R) -F, or -Cl;

where R_A is:

- (I)- C_1 - C_{10} alkyl optionally substituted with one, two or three
 5 substituents selected from the group consisting of C_1 - C_3 alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C \equiv N, -CF₃, C_1 - C_6 alkoxy, -O-phenyl, -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above, -OC=O NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above, -S(=O)₀₋₂ R_{1-a} where R_{1-a} is as defined above, -NR_{1-a}C=O NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above, -C=O NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above, and -
 10 S(=O)₂ NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

- (II) -(CH₂)₀₋₃-(C₃-C₈) cycloalkyl where cycloalkyl can be optionally substituted with one, two or three substituents selected from the group consisting of C_1 - C_3 alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C \equiv N, -CF₃, C_1 - C_6 alkoxy, -O-phenyl, -CO-OH, -CO-O-(C_1 - C_4 alkyl), and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined
 15 above,

(III) -(CR_{A-x}R_{A-y})₀₋₄-R_{A-aryl} where R_{A-x} and R_{A-y} are

- (A) -H,
 (B) C_1 - C_4 alkyl optionally substituted with one or two -OH,
 (C) C_1 - C_4 alkoxy optionally substituted with one, two, or
 20 three of -F,
 (D) -(CH₂)₀₋₄- C_3 - C_7 cycloalkyl,
 (E) C_2 - C_6 alkenyl containing one or two double bonds,
 (F) C_2 - C_6 alkynyl containing one or two triple bonds, or
 (G) phenyl,

- 25 and where R_{A-x} and R_{A-y} are taken together with the carbon to which they are attached to form a carbocycle of three, four, five, six or seven carbon atoms, optionally where one carbon atom is replaced by a heteroatom selected from the group consisting of -O-, -S-, -SO₂-, and -NR_{N-2}- and R_{A-aryl} is the same as R_{N-aryl},

- (IV) -(CR_{A-x}R_{A-y})₀₋₄-R_{A-heteroaryl} where R_{A-heteroaryl} is the same as R_{N-heteroaryl}
 30 heteroaryl and R_{A-x} and R_{A-y} are as defined above,

(V) -(CR_{A-x}R_{A-y})₀₋₄-R_{A-aryl}-R_{A-aryl} where R_{A-aryl}, R_{A-x} and R_{A-y} are as defined above,

(VI) $-(\text{CR}_{\text{A-x}}\text{R}_{\text{A-y}})_{0-4}-\text{R}_{\text{A-aryl}}-\text{R}_{\text{A-heteroaryl}}$ where $\text{R}_{\text{A-aryl}}$, $\text{R}_{\text{A-heteroaryl}}$, $\text{R}_{\text{A-x}}$ and $\text{R}_{\text{A-y}}$ are as defined above,

(VII) $-(\text{CR}_{\text{A-x}}\text{R}_{\text{A-y}})_{0-4}-\text{R}_{\text{A-heteroaryl}}-\text{R}_{\text{A-aryl}}$ where $\text{R}_{\text{A-heteroaryl}}$, $\text{R}_{\text{A-aryl}}$, $\text{R}_{\text{A-x}}$ and $\text{R}_{\text{A-y}}$ are as defined above,

5 (VIII) $-(\text{CR}_{\text{A-x}}\text{R}_{\text{A-y}})_{0-4}-\text{R}_{\text{A-heteroaryl}}-\text{R}_{\text{A-heteroaryl}}$ where $\text{R}_{\text{A-heteroaryl}}$, $\text{R}_{\text{A-x}}$ and $\text{R}_{\text{A-y}}$ are as defined above,

(IX) $-(\text{CR}_{\text{A-x}}\text{R}_{\text{A-y}})_{0-4}-\text{R}_{\text{A-aryl}}-\text{R}_{\text{A-heterocycle}}$ where $\text{R}_{\text{A-heterocycle}}$ is defined as $\text{R}_{1\text{-heterocycle}}$, and where $\text{R}_{\text{A-aryl}}$, $\text{R}_{\text{A-x}}$ and $\text{R}_{\text{A-y}}$ are as defined above,

10 (X) $-(\text{CR}_{\text{A-x}}\text{R}_{\text{A-y}})_{0-4}-\text{R}_{\text{A-heteroaryl}}-\text{R}_{\text{A-heterocycle}}$ where $\text{R}_{\text{A-heteroaryl}}$, $\text{R}_{\text{A-heterocycle}}$, $\text{R}_{\text{A-x}}$ and $\text{R}_{\text{A-y}}$ are as defined above,

(XI) $-(\text{CR}_{\text{A-x}}\text{R}_{\text{A-y}})_{0-4}-\text{R}_{\text{A-heterocycle}}-\text{R}_{\text{A-aryl}}$ where $\text{R}_{\text{A-heterocycle}}$, $\text{R}_{\text{A-aryl}}$, $\text{R}_{\text{A-x}}$ and $\text{R}_{\text{A-y}}$ are as defined above,

(XII) $-(\text{CR}_{\text{A-x}}\text{R}_{\text{A-y}})_{0-4}-\text{R}_{\text{A-heterocycle}}-\text{R}_{\text{A-heteroaryl}}$ where $\text{R}_{\text{A-heterocycle}}$, $\text{R}_{\text{A-heteroaryl}}$, $\text{R}_{\text{A-x}}$ and $\text{R}_{\text{A-y}}$ are as defined above,

15 (XIII) $-(\text{CR}_{\text{A-x}}\text{R}_{\text{A-y}})_{0-4}-\text{R}_{\text{A-heterocycle}}-\text{R}_{\text{A-heterocycle}}$ where $\text{R}_{\text{A-heterocycle}}$, $\text{R}_{\text{A-x}}$ and $\text{R}_{\text{A-y}}$ are as defined above,

(XIV) $-(\text{CR}_{\text{A-x}}\text{R}_{\text{A-y}})_{0-4}-\text{R}_{\text{A-heterocycle}}$ where $\text{R}_{\text{A-heterocycle}}$, $\text{R}_{\text{A-x}}$ and $\text{R}_{\text{A-y}}$ are as defined above,

20 (XV) $-\text{[C(R}_{\text{A-1}})(\text{R}_{\text{A-2}})]_{1-3}-\text{CO-N-(R}_{\text{A-3}})_2$ where $\text{R}_{\text{A-1}}$ and $\text{R}_{\text{A-2}}$ are the same or different and are selected from the group consisting of:

(A) -H,

(B) $\text{-C}_1\text{-C}_6$ alkyl, optionally substituted with one, two or three substituents selected from the group consisting of $\text{C}_1\text{-C}_3$ alkyl, -F, -Cl, -Br, -I, -OH, -SH, $\text{-C}\equiv\text{N}$, -CF_3 , $\text{C}_1\text{-C}_6$ alkoxy, -O-phenyl, and $\text{-NR}_{1\text{-a}}\text{R}_{1\text{-b}}$ where $\text{R}_{1\text{-a}}$ and $\text{R}_{1\text{-b}}$ are as defined above,

(C) $\text{C}_2\text{-C}_6$ alkenyl with one or two double bonds, optionally substituted with one, two or three substituents selected from the group consisting of $\text{C}_1\text{-C}_3$ alkyl, -F, -Cl, -Br, -I, -OH, -SH, $\text{-C}\equiv\text{N}$, -CF_3 , $\text{C}_1\text{-C}_6$ alkoxy, -O-phenyl, and $\text{-NR}_{1\text{-a}}\text{R}_{1\text{-b}}$ where $\text{R}_{1\text{-a}}$ and $\text{R}_{1\text{-b}}$ are as defined above,

30 (D) $\text{C}_2\text{-C}_6$ alkynyl with one or two triple bonds, optionally substituted with one, two or three substituents selected from the group consisting of $\text{C}_1\text{-C}_3$ alkyl, -F, -Cl, -Br, -I, -OH, -SH, $\text{-C}\equiv\text{N}$, -CF_3 , $\text{C}_1\text{-C}_6$ alkoxy, -O-phenyl, and $\text{-NR}_{1\text{-a}}\text{R}_{1\text{-b}}$ where $\text{R}_{1\text{-a}}$ and $\text{R}_{1\text{-b}}$ are as defined above,

(E) $-(\text{CH}_2)_{1-2}-\text{S}(\text{O})_{0-2}-(\text{C}_1-\text{C}_6 \text{ alkyl})$,

(F) $-(\text{CH}_2)_{0-4}-\text{C}_3-\text{C}_7$ cycloalkyl, optionally substituted with one, two or three substituents selected from the group consisting of C_1-C_3 alkyl, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, $-\text{OH}$, $-\text{SH}$, $-\text{C}\equiv\text{N}$, $-\text{CF}_3$, C_1-C_6 alkoxy, $-\text{O}$ -phenyl, and $-\text{NR}_{1-a}\text{R}_{1-b}$ where

5 R_{1-a} and R_{1-b} are as defined above,

(G) $-(\text{C}_1-\text{C}_4 \text{ alkyl})-\text{R}_{\text{A}'\text{-aryl}}$ where $\text{R}_{\text{A}'\text{-aryl}}$ is as defined for $\text{R}_{1\text{-aryl}}$

aryl,

(H) $-(\text{C}_1-\text{C}_4 \text{ alkyl})-\text{R}_{\text{A-heteroaryl}}$ where $\text{R}_{\text{A-heteroaryl}}$ is as defined

above,

10 (I) $-(\text{C}_1-\text{C}_4 \text{ alkyl})-\text{R}_{\text{A-heterocycle}}$ where $\text{R}_{\text{A-heterocycle}}$ is as defined above,

(J) $-\text{R}_{\text{A-heteroaryl}}$ where $\text{R}_{\text{A-heteroaryl}}$ is as defined above,

(K) $-\text{R}_{\text{A-heterocycle}}$ where $\text{R}_{\text{A-heterocycle}}$ is as defined above,

(M) $-(\text{CH}_2)_{1-4}-\text{R}_{\text{A-4}}-(\text{CH}_2)_{0-4}-\text{R}_{\text{A}'\text{-aryl}}$ where $\text{R}_{\text{A-4}}$ is $-\text{O}-$, $-\text{S}-$ or $-\text{NR}_{\text{A-5}}-$ where $\text{R}_{\text{A-5}}$ is C_1-C_6 alkyl, and where $\text{R}_{\text{A}'\text{-aryl}}$ is defined above,

15 (N) $-(\text{CH}_2)_{1-4}-\text{R}_{\text{A-4}}-(\text{CH}_2)_{0-4}-\text{R}_{\text{A-heteroaryl}}$ where $\text{R}_{\text{A-4}}$ and $\text{R}_{\text{A-heteroaryl}}$ are as defined above, and

(O) $-\text{R}_{\text{A}'\text{-aryl}}$ where $\text{R}_{\text{A}'\text{-aryl}}$ is as defined above,

and where $\text{R}_{\text{A-3}}$ is the same or different and is:

20 (A) $-\text{H}$,

(B) C_1-C_6 alkyl optionally substituted with one, two or three substituents selected from the group consisting of C_1-C_3 alkyl, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, $-\text{OH}$, $-\text{SH}$, $-\text{C}\equiv\text{N}$, $-\text{CF}_3$, C_1-C_6 alkoxy, $-\text{O}$ -phenyl, $-\text{NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

25 (C) C_2-C_6 alkenyl with one or two double bonds, optionally substituted with one, two or three substituents selected from the group consisting of C_1-C_3 alkyl, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, $-\text{OH}$, $-\text{SH}$, $-\text{C}\equiv\text{N}$, $-\text{CF}_3$, C_1-C_6 alkoxy, $-\text{O}$ -phenyl, and $-\text{NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

30 (D) C_2-C_6 alkynyl with one or two triple bonds, optionally substituted with one, two or three substituents selected from the group consisting of C_1-C_3 alkyl, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, $-\text{OH}$, $-\text{SH}$, $-\text{C}\equiv\text{N}$, $-\text{CF}_3$, C_1-C_6 alkoxy, $-\text{O}$ -phenyl, and $-\text{NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

(E) $-(CH_2)_{0-4}-C_3-C_7$ cycloalkyl, optionally substituted with one, two or three substituents selected from the group consisting of C_1-C_3 alkyl, $-F$, $-Cl$, $-Br$, $-I$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, C_1-C_6 alkoxy, $-O$ -phenyl, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

5 (F) $-R_{A'-aryl}$ where $R_{A'-aryl}$ is as defined above,

(G) $-R_{A-heteroaryl}$ where $R_{A-heteroaryl}$ is as defined above,

(H) $-R_{A-heterocycle}$ where $R_{A-heterocycle}$ is as defined above,

(I) $-(C_1-C_4 \text{ alkyl})-R_{A'-aryl}$ where $R_{A'-aryl}$ is as defined above,

10 (J) $-(C_1-C_4 \text{ alkyl})-R_{A-heteroaryl}$ where $R_{A-heteroaryl}$ is as defined above,

(K) $-(C_1-C_4 \text{ alkyl})-R_{A-heterocycle}$ where $R_{A-heterocycle}$ is as defined above, or

(XVI) $-\text{CH}(R_{A-aryl})_2$ where R_{A-aryl} are the same or different and are as defined above,

(XVII) $-\text{CH}(R_{A-heteroaryl})_2$ where $R_{A-heteroaryl}$ are the same or different and are as defined above,

(XVIII) $-\text{CH}(R_{A-aryl})(R_{A-heteroaryl})$ where R_{A-aryl} and $R_{A-heteroaryl}$ are as defined above,

20 (XIX) -cyclopentyl, -cyclohexyl, or -cycloheptyl ring fused to R_{A-aryl} , $R_{A-heteroaryl}$, $R_{A-heterocycle}$ where R_{A-aryl} or $R_{A-heteroaryl}$ or $R_{A-heterocycle}$ are as defined above where one carbon of cyclopentyl, cyclohexyl, or -cycloheptyl is optionally replaced with NH , NR_{N-5} , O , or $S(=O)_{0-2}$, and where cyclopentyl, cyclohexyl, or -cycloheptyl can be optionally substituted with one or two $-C_1-C_3$ alkyl, $-F$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, C_1-C_6 alkoxy, $=O$, or $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

(XX) C_2-C_{10} alkenyl containing one or two double bonds optionally substituted with one, two or three substituents selected from the group consisting of C_1-C_3 alkyl, $-F$, $-Cl$, $-Br$, $-I$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, C_1-C_6 alkoxy, $-O$ -phenyl, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

30 (XXI) C_2-C_{10} alkynyl containing one or two triple bonds optionally substituted with one, two or three substituents selected from the group consisting of C_1-C_3 alkyl, $-F$, $-Cl$, $-Br$, $-I$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, C_1-C_6 alkoxy, $-O$ -phenyl, $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

(XXI) $-(\text{CH}_2)_{0-1}-\text{CHR}_{\text{A-6}}-(\text{CH}_2)_{0-1}-\text{R}_{\text{A-aryl}}$ where $\text{R}_{\text{A-aryl}}$ is as defined above and $\text{R}_{\text{A-6}}$ is $-(\text{CH}_2)_{0-6}-\text{OH}$,

(XXII) $-(\text{CH}_2)_{0-1}-\text{CHR}_{\text{A-6}}-(\text{CH}_2)_{0-1}-\text{R}_{\text{A-heteroaryl}}$ where $\text{R}_{\text{A-heteroaryl}}$ and $\text{R}_{\text{A-6}}$ is as defined above,

5 (XXIII) $-\text{CH}(-\text{R}_{\text{A-aryl}}$ or $\text{R}_{\text{A-heteroaryl}})-\text{CO}-\text{O}(\text{C}_1-\text{C}_4 \text{ alkyl})$ where $\text{R}_{\text{A-aryl}}$ and $\text{R}_{\text{A-heteroaryl}}$ are as defined above,

(XXIV) $-\text{CH}(-\text{CH}_2-\text{OH})-\text{CH}(-\text{OH})-\text{micro-NO}_2$,

(XXV) $(\text{C}_1-\text{C}_6 \text{ alkyl})-\text{O}-(\text{C}_1-\text{C}_6 \text{ alkyl})-\text{OH}$,

(XXVII) $-\text{CH}_2-\text{NH}-\text{CH}_2-\text{CH}(-\text{O}-\text{CH}_2-\text{CH}_3)_2$,

10 (XXVIII) $-\text{H}$,

(XXIX) $-(\text{CH}_2)_{0-6}-\text{C}(=\text{NR}_{1-a})(\text{NR}_{1-a}\text{R}_{1-b})$ where R_{1-a} and R_{1-b} are as defined above; or

(XXX)

15 $-\text{C}=\text{OC}(\text{HR}_6)\text{NHR}_7$, where R_6 and R_7 are as defined below,
 $-\text{C}=\text{OR}_7$, where R_7 is as defined below,
 $-\text{C}=\text{OOR}_7$, where R_7 is as defined below, or
 $-\text{SOOR}_7$ where R_7 is as defined below,

wherein R_6 is:

20 hydrogen,
 $\text{C}_1 - \text{C}_3$ alkyl,
phenyl,
thioalkoxyalkyl,
alkyl substituted aryl,
cycloalkyl,
25 cycloalkylalkyl,
hydroxyalkyl,
alkoxyalkyl,
aryloxyalkyl,
haloalkyl,
30 carboxyalkyl,
alkoxycarbonylalkyl,
aminoalkyl,
(N-protected)aminoalkyl,
alkylaminoalkyl,

5 ((N-protected)(alkyl)amino)alkyl,
dialkylaminoalkyl,
guanidinoalkyl,
lower alkenyl,
heterocyclic,
(heterocyclic)alkyl),
arylthioalkyl,
arylsulfonylalkyl,
(heterocyclic)thioalkyl,
10 (heterocyclic)sulfonylalkyl,
(heterocyclic)oxyalkyl,
arylalkoxyalkyl,
arylthioalkoxyalkyl,
arylalkylsulfonylalkyl,
15 (heterocyclic))alkoxyalkyl,
(heterocyclic)thioalkoxyalkyl,
(heterocyclic)alkylsulfonylalkyl,
cycloalkyloxyalkyl,
cycloalkylthioalkyl,
20 cycloalkylsulfonylalkyl,
cycloalkylalkoxyalkyl,
cycloalkylthioalkoxyalkyl,
cycloalkylalkylsulfonylalkyl,
aminocarbonyl,
25 alkylaminocarbonyl,
dialkylaminocarbonyl,
aroylalkyl,
(heterocyclic)carbonylalkyl,
polyhydroxyalkyl,
30 aminocarbonylalkyl,
alkylaminocarbonylalkyl,
dialkylaminocarbonylalkyl,
aryloxyalkyl, or
alkylsulfonylalkyl,

wherein heterocyclic is pyridyl, thiazolyl, isothiazolyl, oxazolyl, isoxazolyl, furanyl, thienyl, tetrahydrofuranyl, tetrahydrothienyl and tetrahydro[2H]pyranyl and wherein the heterocycle is unsubstituted or substituted with one to three substituents independently selected from hydroxy, halo, amino, alkylamino, dialkylamino, alkoxy, polyalkoxy, haloalkyl, cycloalkyl, 5 cycloalkylalkyl, aryl, arylalkyl, COOH, -SO₃H, lower alkenyl or lower alkyl;

wherein R₇ is:

C₁ - C₃ alkyl,
phenyl,
10 thioalkoxyalkyl,
(aryl)alkyl,
cycloalkyl,
cycloalkylalkyl,
hydroxyalkyl,
15 alkoxyalkyl,
aryloxyalkyl,
haloalkyl,
carboxyalkyl,
alkoxycarbonylalkyl,
20 aminoalkyl,
(N-protected)aminocalkyl,
alkylaminoalkyl,
((N-protected)(alkyl)amino)alkyl,
dialkylaminoalkyl,
25 guanidinoalkyl,
lower alkenyl,
heterocyclic,
(heterocyclic)alkyl),
arylthioalkyl,
30 arylsulfonyalkyl,
(heterocyclic)thioalkyl,
(heterocyclic)sulfonylalkyl,
(heterocyclic)oxyalkyl,
arylalkoxyalkyl,

- arylthioalkoxyalkyl,
 arylalkylsulfonylalkyl,
 (heterocyclic)alkoxyalkyl,
 (heterocyclic)thioalkoxyalkyl,
 5 (heterocyclic)alkylsulfonylalkyl,
 cycloalkyloxyalkyl,
 cycloalkylthioalkyl,
 cycloalkylsulfonylalkyl,
 cycloalkylalkoxyalkyl,
 10 cycloalkylthioalkoxyalkyl,
 cycloalkylalkylsulfonylalkyl,
 aminocarbonyl,
 alkylaminocarbonyl,
 dialkylaminocarbonyl,
 15 aroylalkyl,
 (heterocyclic)carbonylalkyl,
 polyhydroxyalkyl,
 aminocarbonylalkyl,
 alkylaminocarbonylalkyl,
 20 dialkylaminocarbonylalkyl,
 aryloxyalkyl, or
 alkylsulfonylalkyl,
 wherein heterocyclic is pyridyl, thiazolyl, isothiazolyl,
 oxazolyl, isoxazolyl, furanyl, thienyl, tetrahydrofuranyl, tetrahydrothienyl and
 25 tetrahydro[2H]pyranyl and wherein the heterocycle is unsubstituted or substituted
 with one to three substituents independently selected from hydroxy, halo, amino,
 alkylamino, dialkylamino, alkoxy, polyalkoxy, haloalkyl, cycloalkyl,
 cycloalkylalkyl, aryl, arylalkyl, COOH, -SO₃H, lower alkenyl or lower alkyl;
- 30 where R_B is:
 (I)-C₁-C₁₀ alkyl optionally substituted with one, two or three
 substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH,

- SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above, -OC=O NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above, -S(=O)₀₋₂ R_{1-a} where R_{1-a} is as defined above, -NR_{1-a}C=O NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above, -C=O NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above, and -
- 5 S(=O)₂ NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(II) -(CH₂)₀₋₃-(C₃-C₈) cycloalkyl where cycloalkyl can be optionally substituted with one, two or three substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, -CO-OH, -CO-O-(C₁-C₄ alkyl), and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

- 10 (III) -(CR_{B-x}R_{B-y})₀₋₄-R_{B-aryl} where R_{B-x} and R_{B-y} are

(A) -H,

(B) C₁-C₄ alkyl optionally substituted with one or two -OH,

(C) C₁-C₄ alkoxy optionally substituted with one, two, or

three of -F,

- 15 (D) -(CH₂)₀₋₄-C₃-C₇ cycloalkyl,

(E) C₂-C₆ alkenyl containing one or two double bonds,

(F) C₂-C₆ alkynyl containing one or two triple bonds, or

(G) phenyl,

and where R_{B-x} and R_{B-y} are taken together with the carbon to

- 20 which they are attached to form a carbocycle of three, four, five, six or seven carbon atoms, optionally where one carbon atom is replaced by a heteroatom selected from the group consisting of -O-, -S-, -SO₂-, and -NR_{N-2} where R_{N-2} is as defined above, and R_{B-aryl} is the same as R_{N-aryl} and is defined above

- (IV) -(CR_{B-x}R_{B-y})₀₋₄-R_{B-heteroaryl} where R_{B-heteroaryl} is the same as R_{N-heteroaryl}, R_{B-x}, and R_{B-y} are as defined above,
- 25

(V) -(CR_{B-x}R_{B-y})₀₋₄-R_{B-aryl}-R_{B-aryl} where R_{B-aryl}, R_{B-x}, and R_{B-y} are as defined above,

(VI) -(CR_{B-x}R_{B-y})₀₋₄-R_{B-aryl}-R_{B-heteroaryl} where R_{B-aryl}, R_{B-heteroaryl}, R_{B-x} and R_{B-y} are as defined above,

- 30 (VII) -(CR_{B-x}R_{B-y})₀₋₄-R_{B-heteroaryl}-R_{B-aryl} where R_{B-heteroaryl}, R_{B-aryl}, R_{B-x} and R_{B-y} are as defined above,

(VIII) -(CR_{B-x}R_{B-y})₀₋₄-R_{B-heteroaryl}-R_{B-heteroaryl} where R_{B-heteroaryl}, R_{B-x} and R_{B-y} are as defined above,

(IX) $-(\text{CR}_{\text{B-x}}\text{R}_{\text{B-y}})_{0-4}-\text{R}_{\text{B-aryl}}-\text{R}_{\text{B-heterocycle}}$ where $\text{R}_{\text{B-heterocycle}}$ is defined as $\text{R}_{\text{1-heterocycle}}$, and where $\text{R}_{\text{B-aryl}}$, $\text{R}_{\text{B-x}}$ and $\text{R}_{\text{B-y}}$ are as defined above,

(X) $-(\text{CR}_{\text{B-x}}\text{R}_{\text{B-y}})_{0-4}-\text{R}_{\text{B-heteroaryl}}-\text{R}_{\text{B-heterocycle}}$ where $\text{R}_{\text{B-heteroaryl}}$, $\text{R}_{\text{B-heterocycle}}$, $\text{R}_{\text{B-x}}$ and $\text{R}_{\text{B-y}}$ are as defined above,

5 (XI) $-(\text{CR}_{\text{B-x}}\text{R}_{\text{B-y}})_{0-4}-\text{R}_{\text{B-heterocycle}}-\text{R}_{\text{B-aryl}}$ where $\text{R}_{\text{B-heterocycle}}$, $\text{R}_{\text{B-aryl}}$, $\text{R}_{\text{B-x}}$ and $\text{R}_{\text{B-y}}$ are as defined above,

(XII) $-(\text{CR}_{\text{B-x}}\text{R}_{\text{B-y}})_{0-4}-\text{R}_{\text{B-heterocycle}}-\text{R}_{\text{B-heteroaryl}}$ where $\text{R}_{\text{B-heterocycle}}$, $\text{R}_{\text{B-heteroaryl}}$, $\text{R}_{\text{B-x}}$ and $\text{R}_{\text{B-y}}$ are as defined above,

10 (XIII) $-(\text{CR}_{\text{B-x}}\text{R}_{\text{B-y}})_{0-4}-\text{R}_{\text{B-heterocycle}}-\text{R}_{\text{B-heterocycle}}$ where $\text{R}_{\text{B-heterocycle}}$, $\text{R}_{\text{B-x}}$ and $\text{R}_{\text{B-y}}$ are as defined above,

(XIV) $-(\text{CR}_{\text{B-x}}\text{R}_{\text{B-y}})_{0-4}-\text{R}_{\text{B-heterocycle}}$ where $\text{R}_{\text{B-heterocycle}}$, $\text{R}_{\text{B-x}}$ and $\text{R}_{\text{B-y}}$ are as defined above,

(XV) $-\text{[C(R}_{\text{B-1}}\text{)(R}_{\text{B-2}}\text{)]}_{1-3}-\text{CO-N-(R}_{\text{B-3}}\text{)}_2$ where $\text{R}_{\text{B-1}}$ and $\text{R}_{\text{B-2}}$ are the same or different and are selected from the group consisting of:

15 (A) -H,

(B) $\text{C}_1\text{-C}_6$ alkyl, optionally substituted with one, two or three substituents selected from the group consisting of $\text{C}_1\text{-C}_3$ alkyl, -F, -Cl, -Br, -I, -OH, -SH, $\text{-C}\equiv\text{N}$, -CF_3 , $\text{C}_1\text{-C}_6$ alkoxy, -O-phenyl, and $\text{-NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

20 (C) $\text{C}_2\text{-C}_6$ alkenyl with one or two double bonds, optionally substituted with one, two or three substituents selected from the group consisting of $\text{C}_1\text{-C}_3$ alkyl, -F, -Cl, -Br, -I, -OH, -SH, $\text{-C}\equiv\text{N}$, -CF_3 , $\text{C}_1\text{-C}_6$ alkoxy, -O-phenyl, and $\text{-NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

(D) $\text{C}_2\text{-C}_6$ alkynyl with one or two triple bonds, optionally substituted with one, two or three substituents selected from the group consisting of $\text{C}_1\text{-C}_3$ alkyl, -F, -Cl, -Br, -I, -OH, -SH, $\text{-C}\equiv\text{N}$, -CF_3 , $\text{C}_1\text{-C}_6$ alkoxy, -O-phenyl, and $\text{-NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

(E) $-(\text{CH}_2)_{1-2}-\text{S(O)}_{0-2}-(\text{C}_1\text{-C}_6 \text{ alkyl})$,

25 (F) $-(\text{CH}_2)_{0-4}-\text{C}_3\text{-C}_7$ cycloalkyl, optionally substituted with one, two or three substituents selected from the group consisting of $\text{C}_1\text{-C}_3$ alkyl, -F, -Cl, -Br, -I, -OH, -SH, $\text{-C}\equiv\text{N}$, -CF_3 , $\text{C}_1\text{-C}_6$ alkoxy, -O-phenyl, and $\text{-NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

30

- (G) $-(C_1-C_4 \text{ alkyl})-R_{B'-aryl}$ where $R_{B'-aryl}$ is as defined above for R_{1-aryl} ,
- (H) $-(C_1-C_4 \text{ alkyl})-R_{B-heteroaryl}$ where $R_{B-heteroaryl}$ is as defined above,
- 5 (I) $-(C_1-C_4 \text{ alkyl})-R_{B-heterocycle}$ where $R_{B-heterocycle}$ is as defined above,
- (J) $-R_{B-heteroaryl}$ where $R_{B-heteroaryl}$ is as defined above,
- (K) $-R_{B-heterocycle}$ where $R_{B-heterocycle}$ is as defined above,
- (M) $-(CH_2)_{1-4}-R_{B-4}-(CH_2)_{0-4}-R_{B'-aryl}$ where R_{B-4} is $-O-$, $-S-$ or
- 10 $-NR_{B-5}-$ where R_{B-5} is C_1-C_6 alkyl, and where $R_{B'-aryl}$ is defined above,
- (N) $-(CH_2)_{1-4}-R_{B-4}-(CH_2)_{0-4}-R_{B-heteroaryl}$ where R_{B-4} and $R_{B-heteroaryl}$ are as defined above, and
- (O) $-R_{B'-aryl}$ where $R_{B'-aryl}$ is as defined above,
- and where R_{B-3} is the same or different and is:
- 15 (A) $-H$,
- (B) $-C_1-C_6$ alkyl optionally substituted with one, two or three substituents selected from the group consisting of C_1-C_3 alkyl, $-F$, $-Cl$, $-Br$, $-I$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, C_1-C_6 alkoxy, $-O$ -phenyl, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,
- 20 (C) C_2-C_6 alkenyl with one or two double bonds, optionally substituted with one, two or three substituents selected from the group consisting of C_1-C_3 alkyl, $-F$, $-Cl$, $-Br$, $-I$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, C_1-C_6 alkoxy, $-O$ -phenyl, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,
- (D) C_2-C_6 alkynyl with one or two triple bonds, optionally
- 25 substituted with one, two or three substituents selected from the group consisting of C_1-C_3 alkyl, $-F$, $-Cl$, $-Br$, $-I$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, C_1-C_6 alkoxy, $-O$ -phenyl, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,
- (E) $-(CH_2)_{0-4}-C_3-C_7$ cycloalkyl, optionally substituted with one, two or three substituents selected from the group consisting of C_1-C_3 alkyl, $-F$,
- 30 $-Cl$, $-Br$, $-I$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, C_1-C_6 alkoxy, $-O$ -phenyl, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,
- (F) $-R_{B'-aryl}$ where $R_{B'-aryl}$ is as defined above,
- (G) $-R_{B-heteroaryl}$ where $R_{B-heteroaryl}$ is as defined above,

(H) $-R_{B\text{-heterocycle}}$ where $R_{B\text{-heterocycle}}$ is as defined above,

(I) $-(C_1-C_4 \text{ alkyl})-R_{B'\text{-aryl}}$ where $R_{B'\text{-aryl}}$ is as defined above,

(J) $-(C_1-C_4 \text{ alkyl})-R_{B\text{-heteroaryl}}$ where $R_{B\text{-heteroaryl}}$ is as defined above,

(K) $-(C_1-C_4 \text{ alkyl})-R_{B\text{-heterocycle}}$ where $R_{B\text{-heterocycle}}$ is as defined above, or

(XVI) $-\text{CH}(R_{B\text{-aryl}})_2$ where $R_{B\text{-aryl}}$ are the same or different and are as defined above,

(XVII) $-\text{CH}(R_{B\text{-heteroaryl}})_2$ where $R_{B\text{-heteroaryl}}$ are the same or different and are as defined above,

(XVIII) $-\text{CH}(R_{B\text{-aryl}})(R_{B\text{-heteroaryl}})$ where $R_{B\text{-aryl}}$ and $R_{B\text{-heteroaryl}}$ are as defined above,

(XIX) -cyclopentyl, -cyclohexyl, or -cycloheptyl ring fused to $R_{B\text{-aryl}}$ or $R_{B\text{-heteroaryl}}$ or $R_{B\text{-heterocycle}}$ where $R_{B\text{-aryl}}$ or $R_{B\text{-heteroaryl}}$ or $R_{B\text{-heterocycle}}$ are as defined above where one carbon of cyclopentyl, cyclohexyl, or -cycloheptyl is optionally replaced with NH, NR_{N-5} , O, or $\text{S}(=\text{O})_{0-2}$, and where cyclopentyl, cyclohexyl, or -cycloheptyl can be optionally substituted with one or two $-C_1-C_3$ alkyl, -F, -OH, -SH, $-\text{C}\equiv\text{N}$, $-\text{CF}_3$, C_1-C_6 alkoxy, =O, or $-\text{NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

(XX) C_2-C_{10} alkenyl containing one or two double bonds optionally substituted with one, two or three substituents selected from the group consisting of C_1-C_3 alkyl, -F, -Cl, -Br, -I, -OH, -SH, $-\text{C}\equiv\text{N}$, $-\text{CF}_3$, C_1-C_6 alkoxy, -O-phenyl, and $-\text{NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

(XXI) C_2-C_{10} alkynyl containing one or two triple bonds optionally substituted with one, two or three substituents selected from the group consisting of C_1-C_3 alkyl, -F, -Cl, -Br, -I, -OH, -SH, $-\text{C}\equiv\text{N}$, $-\text{CF}_3$, C_1-C_6 alkoxy, -O-phenyl, and $-\text{NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

(XXI) $-(\text{CH}_2)_{0-1}-\text{CHR}_{C-6}-(\text{CH}_2)_{0-1}-R_{B\text{-aryl}}$ where $R_{B\text{-aryl}}$ is as defined above and R_{C-6} is $-(\text{CH}_2)_{0-6}-\text{OH}$,

(XXII) $-(\text{CH}_2)_{0-1}-\text{CHR}_{B-6}-(\text{CH}_2)_{0-1}-R_{B\text{-heteroaryl}}$ where $R_{B\text{-heteroaryl}}$ and R_{C-6} is as defined above,

(XXIII) $-\text{CH}(-\text{R}_{\text{B-aryl}}$ or $\text{R}_{\text{B-heteroaryl}})-\text{CO}-\text{O}(\text{C}_1-\text{C}_4 \text{ alkyl})$ where $\text{R}_{\text{B-aryl}}$

and $\text{R}_{\text{B-heteroaryl}}$ are as defined above,

(XXIV) $-\text{CH}(-\text{CH}_2-\text{OH})-\text{CH}(-\text{OH})-\text{micro-NO}_2$,

(XXV) $(\text{C}_1-\text{C}_6 \text{ alkyl})-\text{O}-(\text{C}_1-\text{C}_6 \text{ alkyl})-\text{OH}$,

5 (XXVII) $-\text{CH}_2-\text{NH}-\text{CH}_2-\text{CH}(-\text{O}-\text{CH}_2-\text{CH}_3)_2$,

(XXVIII) $-\text{H}$, or

(XXIX) $-(\text{CH}_2)_{0-6}-\text{C}(=\text{NR}_{1-a})(\text{NR}_{1-a}\text{R}_{1-b})$ where R_{1-a} and R_{1-b} are as defined above; and

where PROTECTING GROUP is selected from the group consisting of *t*-

10 butoxycarbonyl, benzyloxycarbonyl, formyl, trityl, acetyl, trichloroacetyl, dichloroacetyl, chloroacetyl, trifluoroacetyl, difluoroacetyl, fluoroacetyl, 4-phenylbenzyloxycarbonyl, 2-methylbenzyloxycarbonyl, 4-ethoxybenzyloxycarbonyl, 4-fluorobenzyloxycarbonyl, 4-chlorobenzyloxycarbonyl, 3-chlorobenzyloxycarbonyl, 2-chlorobenzyloxycarbonyl, 2,4-
15 dichlorobenzyloxycarbonyl, 4-bromobenzyloxycarbonyl, 3-bromobenzyloxycarbonyl, 4-nitrobenzyloxycarbonyl, 4-cyanobenzyloxycarbonyl, 2-(4-xenyl)isopropoxycarbonyl, 1,1-diphenyleth-1-yloxycarbonyl, 1,1-diphenylprop-1-yloxycarbonyl, 2-phenylprop-2-yloxycarbonyl, 2-(*p*-toluyl)prop-2-yloxycarbonyl, cyclopentanyloxycarbonyl, 1-methylcyclopentanyloxycarbonyl,
20 cyclohexanyloxycarbonyl, 1-methylcyclohexanyloxycarbonyl, 2-methylcyclohexanyloxycarbonyl, 2-(4-toluylsulfonyl)ethoxycarbonyl, 2-(methylsulfonyl)ethoxycarbonyl, 2-(triphenylphosphino)ethoxycarbonyl, fluorenylmethoxycarbonyl, 2-(trimethylsilyl)ethoxycarbonyl, allyloxycarbonyl, 1-(trimethylsilylmethyl)prop-1-enyloxycarbonyl, 5-benzisoxalylmethoxycarbonyl, 4-
25 acetoxybenzyloxycarbonyl, 2,2,2-trichloroethoxycarbonyl, 2-ethynyl-2-propoxycarbonyl, cyclopropylmethoxycarbonyl, 4-(decyloxy)benzyloxycarbonyl, isobornyloxycarbonyl and 1-piperidyloxycarbonyl, 9-fluorenylmethyl carbonate, $-\text{CH}-\text{CH}=\text{CH}_2$ and phenyl- $\text{C}(=\text{N})-\text{H}$.

30 32. A protected compound according to claim 31

where R_1 is:

$-(\text{CH}_2)_{0-1}-(\text{R}_{1-\text{aryl}})$, or

$-(\text{CH}_2)_{n1}-(\text{R}_{1-\text{heteroaryl}})$;

where R_N is:

$R_{N-1}-X_N-$, where X_N is selected from the group consisting of:

$-\text{CO}-$, and

$-\text{SO}_2-$,

where R_{N-1} is selected from the group consisting of:

5

$-\text{R}_{N\text{-aryl}}$, and

$-\text{R}_{N\text{-heteroaryl}}$, or

$-\text{CO}-\text{CH}(-(\text{CH}_2)_{0-2}-\text{O}-\text{R}_{N-10})-(\text{CH}_2)_{0-2}-\text{R}_{N\text{-aryl}}/\text{R}_{N\text{-heteroaryl}}$;

where R_A is:

$-\text{C}_1-\text{C}_8$ alkyl,

10

$-(\text{CH}_2)_{0-3}-(\text{C}_3-\text{C}_7)$ cycloalkyl,

$-(\text{CR}_{A-x}\text{R}_{A-y})_{0-4}-\text{R}_{A\text{-aryl}}$,

$-(\text{CR}_{A-x}\text{R}_{A-y})_{0-4}-\text{R}_{A\text{-heteroaryl}}$,

$-(\text{CR}_{A-x}\text{R}_{A-y})_{0-4}-\text{R}_{A\text{-heterocycle}}$,

$-\text{cyclopentyl}$ or $-\text{cyclohexyl}$ ring fused to $\text{R}_{A\text{-aryl}}$ or $\text{R}_{A\text{-heteroaryl}}$ or $\text{R}_{A\text{-}}$

15

heterocycle;

where R_B is:

$-\text{C}_1-\text{C}_8$ alkyl,

$-(\text{CH}_2)_{0-3}-(\text{C}_3-\text{C}_7)$ cycloalkyl,

$-(\text{CR}_{A-x}\text{R}_{A-y})_{0-4}-\text{R}_{A\text{-aryl}}$,

20

$-(\text{CR}_{A-x}\text{R}_{A-y})_{0-4}-\text{R}_{A\text{-heteroaryl}}$,

$-(\text{CR}_{A-x}\text{R}_{A-y})_{0-4}-\text{R}_{A\text{-heterocycle}}$,

$-\text{cyclopentyl}$ or $-\text{cyclohexyl}$ ring fused to $\text{R}_{A\text{-aryl}}$ or $\text{R}_{A\text{-heteroaryl}}$ or $\text{R}_{A\text{-}}$

heterocycle.

25 33. A protected compound according to claim 31

where R_1 is:

$-(\text{CH}_2)-(\text{R}_{1\text{-aryl}})$, or

$-(\text{CH}_2)-(\text{R}_{1\text{-heteroaryl}})$;

where R_2 is $-\text{H}$;

30

where R_3 is $-\text{H}$;

where R_N is:

$R_{N-1}-X_N-$ where X_N is:

$-\text{CO}-$,

where R_{N-1} is selected from the group consisting of:

$-R_{N\text{-aryl}}$, and

$-R_{N\text{-heteroaryl}}$;

where R_A is:

$-C_1\text{-}C_8$ alkyl,

5 $-(CH_2)_{0-3}\text{-(}C_3\text{-}C_7\text{) cycloalkyl}$,

$-(CR_{A-x}R_{A-y})_{0-4}\text{-}R_{A\text{-aryl}}$,

$-(CR_{A-x}R_{A-y})_{0-4}\text{-}R_{A\text{-heteroaryl}}$,

$-(CR_{A-x}R_{A-y})_{0-4}\text{-}R_{A\text{-heterocycle}}$, or

10 $-\text{cyclopentyl}$ or $-\text{cyclohexyl}$ ring fused to $R_{A\text{-aryl}}$ or $R_{A\text{-heteroaryl}}$ or $R_{A\text{-heterocycle}}$;

where R_B is:

$-C_1\text{-}C_8$ alkyl,

$-(CH_2)_{0-3}\text{-(}C_3\text{-}C_7\text{) cycloalkyl}$,

$-(CR_{B-x}R_{B-y})_{0-4}\text{-}R_{B\text{-aryl}}$, or

15 $-(CR_{B-x}R_{B-y})_{0-4}\text{-}R_{B\text{-heteroaryl}}$.

$-(CR_{B-x}R_{B-y})_{0-4}\text{-}R_{B\text{-heterocycle}}$,


$-\text{cyclopentyl}$ or $-\text{cyclohexyl}$ ring fused to $R_{B\text{-aryl}}$ or $R_{B\text{-heteroaryl}}$ or $R_{B\text{-heterocycle}}$.

20 34. A protected compound according to claim 31 where PROTECTING GROUP is *t*-butoxycarbonyl.

35. A protected compound according to claim 31 where PROTECTING GROUP is benzyloxycarbonyl.

25

36. A protected compound of the formula (III)



30

where R_1 is:

(I) $C_1\text{-}C_6$ alkyl, optionally substituted with one, two or three substituents selected from the group consisting of $C_1\text{-}C_3$ alkyl, $C_1\text{-}C_7$ alkyl

(optionally substituted with C₁-C₃ alkyl and C₁-C₃ alkoxy), -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₃ alkoxy, -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are -H or C₁-C₆ alkyl, and -OC=O NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(II) -CH₂-S(O)₀₋₂-(C₁-C₆ alkyl),

5 (III) -CH₂-CH₂-S(O)₀₋₂-(C₁-C₆ alkyl),

(IV) C₂-C₆ alkenyl with one or two double bonds, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, -C≡N, -CF₃, C₁-C₃ alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are -H or C₁-C₆ alkyl,

10 (V) C₂-C₆ alkynyl with one or two triple bonds, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, -C≡N, -CF₃, C₁-C₃ alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are -H or C₁-C₆ alkyl,

(VI) -(CH₂)_{n1}-(R_{1-aryl}) where n₁ is zero or one and where R_{1-aryl} is 15 phenyl, 1-naphthyl, 2-naphthyl and indanyl, indenyl, dihydronaphthalyl, or tetralinyl optionally substituted with one, two, three or four of the following substituents on the aryl ring:

(A) C₁-C₆ alkyl optionally substituted with one, two or three substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, 20 -SH, -C≡N, -CF₃, and C₁-C₃ alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(B) C₂-C₆ alkenyl with one or two double bonds, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, -C≡N, -CF₃, C₁-C₃ alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are 25 -H or C₁-C₆ alkyl,

(C) C₂-C₆ alkynyl with one or two triple bonds, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, -C≡N, -CF₃, C₁-C₃ alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are -H or C₁-C₆ alkyl,

30 (D) -F, Cl, -Br or -I,

(F) -C₁-C₆ alkoxy optionally substituted with one, two or three of -F,

(G) -NR_{N-2}R_{N-3} where R_{N-2} and R_{N-3} are as defined below,

(H) -OH,

(I) -C≡N,

(J) C₃-C₇ cycloalkyl, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, -C≡N, -

5 CF₃, C₁-C₃ alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are -H or C₁-C₆ alkyl,

(K) -CO-(C₁-C₄ alkyl),

(L) -SO₂-NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(M) -CO-NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

or

10 (N) -SO₂-(C₁-C₄ alkyl),

(VII) -(CH₂)_{n1}-(R_{1-heteroaryl}) where n₁ is as defined above and where

R_{1-heteroaryl} is selected from the group consisting of:

pyridinyl,

pyrimidinyl,

15 quinolinyl,

benzothienyl,

indolyl,

indolinyl,

pyridazinyl,

20 pyrazinyl,

isoquinolyl,

quinazolinyl,

quinoxalinyl,

phthalazinyl,

25 imidazolyl,

isoxazolyl,

pyrazolyl,

oxazolyl,

thiazolyl,

30 indolizinyl,

indazolyl,

benzothiazolyl,

benzimidazolyl,

	benzofuranyl,
	furanyl,
	thienyl,
	pyrrolyl,
5	oxadiazolyl,
	thiadiazolyl,
	triazolyl,
	tetrazolyl,
	oxazolopyridinyl,
10	imidazopyridinyl,
	isothiazolyl,
	naphthyridinyl,
	cinnolinyI,
	carbazolyl,
15	beta-carbolinyl,
	isochromanyl,
	chromanyl,
	tetrahydroisoquinolinyl,
	isoindolinyl,
20	isobenzotetrahydrofuranyl,
	isobenzotetrahydrothienyl,
	isobenzothienyl,
	benzoxazolyl,
	pyridopyridinyl,
25	benzotetrahydrofuranyl,
	benzotetrahydrothienyl,
	purinyl,
	benzodioxolyl,
	triazinyl,
30	phenoxazinyl,
	phenothiazinyl,
	pteridinyl,
	benzothiazolyl,
	imidazopyridinyl,

	imidazothiazolyl,
	dihydrobenzisoxazinyl,
	benzisoxazinyl,
	benzoxazinyl,
5	dihydrobenzothiazinyl,
	benzopyranyl,
	benzothiopyranyl,
	coumarinyl,
	isocoumarinyl,
10	chromonyl,
	chromanonyl,
	pyridinyl-N-oxide,
	tetrahydroquinolinyl
	dihydroquinolinyl
15	dihydroquinolinonyl
	dihydroisoquinolinonyl
	dihydrocoumarinyl
	dihydroisocoumarinyl
	isoindolinonyl
20	benzodioxanyl
	benzoxazolinonyl
	pyrrolyl N-oxide,
	pyrimidinyl N-oxide,
	pyridazinyl N-oxide,
25	pyrazinyl N-oxide,
	quinolinyl N-oxide,
	indolyl N-oxide,
	indolinyl N-oxide,
	isoquinolyl N-oxide,
30	quinazolinyl N-oxide,
	quinoxalinyl N-oxide,
	phthalazinyl N-oxide,
	imidazolyl N-oxide,
	isoxazolyl N-oxide,

oxazolyl N-oxide,
 thiazolyl N-oxide,
 indoliziny N-oxide,
 indazolyl N-oxide,
 5 benzothiazolyl N-oxide,
 benzimidazolyl N-oxide,
 pyrrolyl N-oxide,
 oxadiazolyl N-oxide,
 thiadiazolyl N-oxide,
 10 triazolyl N-oxide,
 tetrazolyl N-oxide,
 benzothiopyranyl S-oxide, and
 benzothiopyranyl S,S-dioxide,

where the $R_{1\text{-heteroaryl}}$ group is bonded to $-(CH_2)_{n1}-$ by any ring atom
 15 of the parent $R_{1\text{-heteroaryl}}$ group substituted by hydrogen such that the new bond to the
 $R_{1\text{-heteroaryl}}$ group replaces the hydrogen atom and its bond, where heteroaryl is
 optionally substituted with one, two, three or four of:

(1) C_1 - C_6 alkyl optionally substituted with one, two or
 three substituents selected from the group consisting of C_1 - C_3 alkyl, -F, -Cl, -Br, -I,
 20 -OH, -SH, $-C\equiv N$, $-CF_3$, C_1 - C_3 alkoxy, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as
 defined above,

(2) C_2 - C_6 alkenyl with one or two double bonds,
 optionally substituted with one, two or three substituents selected from the group
 consisting of -F, -Cl, -OH, -SH, $-C\equiv N$, $-CF_3$, C_1 - C_3 alkoxy, and $-NR_{1-a}R_{1-b}$ where R_{1-}
 25 a and R_{1-b} are -H or C_1 - C_6 alkyl,

(3) C_2 - C_6 alkynyl with one or two triple bonds,
 optionally substituted with one, two or three substituents selected from the group
 consisting of -F, -Cl, -OH, -SH, $-C\equiv N$, $-CF_3$, C_1 - C_3 alkoxy, and $-NR_{1-a}R_{1-b}$ where R_{1-}
 a and R_{1-b} are -H or C_1 - C_6 alkyl,

30 (4) -F, Cl, -Br or -I,

(6) $-C_1$ - C_6 alkoxy optionally substituted with one, two,
 or three of -F,

(7) $-NR_{N-2}R_{N-3}$ where R_{N-2} and R_{N-3} are as defined
below,

(8) $-OH$,

(9) $-C\equiv N$,

5 (10) C_3-C_7 cycloalkyl, optionally substituted with one,
two or three substituents selected from the group consisting of $-F$, $-Cl$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, C_1-C_3 alkoxy, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are $-H$ or C_1-C_6 alkyl,

(11) $-CO-(C_1-C_4 \text{ alkyl})$,

(12) $-SO_2-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined

10 above,

(13) $-CO-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined

above, or

(14) $-SO_2-(C_1-C_4 \text{ alkyl})$, with the proviso that when n_1
is zero $R_{1-\text{heteroaryl}}$ is not bonded to the carbon chain by nitrogen; or

15 (VIII) $-(CH_2)_{n_1}-(R_{1-\text{heterocycle}})$ where n_1 is as defined above and $R_{1-\text{heterocycle}}$ is selected from the group consisting of:

morpholinyl,

thiomorpholinyl,

thiomorpholinyl S-oxide,

20 thiomorpholinyl S,S-dioxide,

piperazinyl,

homopiperazinyl,

pyrrolidinyl,

pyrrolinyl,

25 tetrahydropyranyl,

piperidinyl,

tetrahydrofuranyl,

tetrahydrothienyl,

homopiperidinyl,

30 homomorpholinyl,

homothiomorpholinyl,

homothiomorpholinyl S,S-dioxide,

oxazolidinonyl,

dihydropyrazolyl,
 dihydropyrrolyl,
 dihydropyrazinyl,
 dihydropyridinyl,
 5 dihydropyrimidinyl,
 dihydrofuryl,
 dihydropyranyl,
 tetrahydrothienyl S-oxide,
 tetrahydrothienyl S,S-dioxide, and
 10 homothiomorpholinyl S-oxide,

where the $R_{1\text{-heterocycle}}$ group is bonded by any atom of the parent $R_{1\text{-heterocycle}}$ group substituted by hydrogen such that the new bond to the $R_{1\text{-heterocycle}}$ group replaces the hydrogen atom and its bond, where heterocycle is optionally substituted with one, two, three or four:

15 (1) $C_1\text{-}C_6$ alkyl optionally substituted with one, two or three substituents selected from the group consisting of $C_1\text{-}C_3$ alkyl, -F, -Cl, -Br, -I, -OH, -SH, $-C\equiv N$, $-CF_3$, $C_1\text{-}C_3$ alkoxy, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

20 (2) $C_2\text{-}C_6$ alkenyl with one or two double bonds, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, $-C\equiv N$, $-CF_3$, $C_1\text{-}C_3$ alkoxy, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are -H or $C_1\text{-}C_6$ alkyl,

25 (3) $C_2\text{-}C_6$ alkynyl with one or two triple bonds, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, $-C\equiv N$, $-CF_3$, $C_1\text{-}C_3$ alkoxy, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are -H or $C_1\text{-}C_6$ alkyl,

30 (4) -F, Cl, -Br or -I,
 (5) $C_1\text{-}C_6$ alkoxy,
 (6) $-C_1\text{-}C_6$ alkoxy optionally substituted with one, two, or three -F,
 (7) $-NR_{N-2}R_{N-3}$ where R_{N-2} and R_{N-3} are as defined below,

(8) -OH,

- (9) $-\text{C}\equiv\text{N}$,
- (10) $\text{C}_3\text{-C}_7$ cycloalkyl, optionally substituted with one, two or three substituents selected from the group consisting of $-\text{F}$, $-\text{Cl}$, $-\text{OH}$, $-\text{SH}$, $-\text{C}\equiv\text{N}$, $-\text{CF}_3$, $\text{C}_1\text{-C}_3$ alkoxy, and $-\text{NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are $-\text{H}$ or $\text{C}_1\text{-C}_6$ alkyl,
- 5 (11) $-\text{CO}(\text{C}_1\text{-C}_4 \text{ alkyl})$,
- (12) $-\text{SO}_2\text{-NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,
- (13) $-\text{CO-NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,
- 10 (14) $-\text{SO}_2(\text{C}_1\text{-C}_4 \text{ alkyl})$, or
- (15) $=\text{O}$, with the proviso that when n_1 is zero $\text{R}_{1\text{-heterocycle}}$ is not bonded to the carbon chain by nitrogen;

where R_2 is:

- 15 (I) $-\text{H}$,
- (II) $\text{C}_1\text{-C}_6$ alkyl, optionally substituted with one, two or three substituents selected from the group consisting of $\text{C}_1\text{-C}_3$ alkyl, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, $-\text{OH}$, $-\text{SH}$, $-\text{C}\equiv\text{N}$, $-\text{CF}_3$, $\text{C}_1\text{-C}_3$ alkoxy, and $-\text{NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,
- 20 (III) $-(\text{CH}_2)_{0-4}\text{-R}_{2-1}$ where R_{2-1} is $\text{R}_{1\text{-aryl}}$ or $\text{R}_{1\text{-heteroaryl}}$ where $\text{R}_{1\text{-aryl}}$ and $\text{R}_{1\text{-heteroaryl}}$ are as defined above;
- (IV) $\text{C}_2\text{-C}_6$ alkenyl with one or two double bonds, optionally substituted with one, two or three substituents selected from the group consisting of $-\text{F}$, $-\text{Cl}$, $-\text{OH}$, $-\text{SH}$, $-\text{C}\equiv\text{N}$, $-\text{CF}_3$, $\text{C}_1\text{-C}_3$ alkoxy, and $-\text{NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are
- 25 $-\text{H}$ or $\text{C}_1\text{-C}_6$ alkyl,
- (V) $\text{C}_2\text{-C}_6$ alkynyl with one or two triple bonds, optionally substituted with one, two or three substituents selected from the group consisting of $-\text{F}$, $-\text{Cl}$, $-\text{OH}$, $-\text{SH}$, $-\text{C}\equiv\text{N}$, $-\text{CF}_3$, $\text{C}_1\text{-C}_3$ alkoxy, and $-\text{NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are $-\text{H}$ or $\text{C}_1\text{-C}_6$ alkyl, or
- 30 (VI) $-(\text{CH}_2)_{0-4}\text{-C}_3\text{-C}_7$ cycloalkyl, optionally substituted with one, two or three substituents selected from the group consisting of $-\text{F}$, $-\text{Cl}$, $-\text{OH}$, $-\text{SH}$, $-\text{C}\equiv\text{N}$, $-\text{CF}_3$, $\text{C}_1\text{-C}_3$ alkoxy, and $-\text{NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are $-\text{H}$ or $\text{C}_1\text{-C}_6$ alkyl;

where R_3 is:

(I)-H,

(II) C_1-C_6 alkyl, optionally substituted with one, two or three substituents selected from the group consisting of C_1-C_3 alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C \equiv N, -CF $_3$, C_1-C_3 alkoxy, and -NR $_{1-a}$ R $_{1-b}$ where R $_{1-a}$ and R $_{1-b}$ are as defined above,

(III) -(CH $_2$) $_{0-4}$ -R $_{2-1}$ where R $_{2-1}$ is R $_{1-aryl}$ or R $_{1-heteroaryl}$ where R $_{1-aryl}$ and R $_{1-heteroaryl}$ are as defined above

(IV) C_2-C_6 alkenyl with one or two double bonds,

(V) C_2-C_6 alkynyl with one or two triple bonds; or

(VI) -(CH $_2$) $_{0-4}$ - C_3-C_7 cycloalkyl, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, -C \equiv N, -CF $_3$, C_1-C_3 alkoxy, and -NR $_{1-a}$ R $_{1-b}$ where R $_{1-a}$ and R $_{1-b}$ are -H or C_1-C_6 alkyl,

and where R $_2$ and R $_3$ are taken together with the carbon to which they are attached to form a carbocycle of three, four, five, six, or seven carbon atoms, optionally where one carbon atom is replaced by a heteroatom selected from the group consisting of -O-, -S-, -SO $_2$ -, and -NR $_{N-2}$ -, where R $_{N-2}$ is selected from the group consisting of:

(a) -H,

(b) C_1-C_6 alkyl optionally substituted with one substituent selected from the group consisting of:

(i) -OH, and

(ii) -NH $_2$,

(c) C_1-C_6 alkyl optionally substituted with one, two, or three -F, -Cl, -Br, or -I,

(d) C_3-C_7 cycloalkyl,

(e) -(C_1-C_2 alkyl)-(C_3-C_7 cycloalkyl),

(f) -(C_1-C_6 alkyl)-O-(C_1-C_3 alkyl),

(g) C_2-C_6 alkenyl with one or two double bonds,

(h) C_2-C_6 alkynyl with one or two triple bonds,

(i) -C₁-C₆ alkyl chain with one double bond

and one triple bond,

(j) -R_{1-aryl} where R_{1-aryl} is as defined above, and

(k) -R_{1-heteroaryl} where R_{1-heteroaryl} is as defined

5 above;

where R_N is:

(I) R_{N-1}-X_N- where X_N is selected from the group consisting of:

(A) -CO-,

10

(B) -SO₂-,

(C) -(CR'R'')₁₋₆ where R' and R'' are the same or different and are -H and C₁-C₄ alkyl,

(D) -CO-(CR'R'')₁₋₆-X_{N-1} where X_{N-1} is selected from the group consisting of -O-, -S- and -NR'- and where R' and R'' are as defined above,

15 and

(E) a single bond;

where R_{N-1} is selected from the group consisting of:

(A) R_{N-aryl} where R_{N-aryl} is phenyl, 1-naphthyl, 2-naphthyl, tetralinyl, indanyl, 6,7,8,9-tetrahydro-5H-benzo[a]cycloheptenyl, or dihydronaphthyl
20 optionally substituted with one, two or three of the following substituents which can be the same or different and are:

(1) C₁-C₆ alkyl, optionally substituted with one, two or three substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₃ alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as
25 defined above,

(2) -OH,

(3) -NO₂,

(4) -F, -Cl, -Br, -I,

(5) -CO-OH,

30

(6) -C≡N,

(7) -(CH₂)₀₋₄-CO-NR_{N-2}R_{N-3} where R_{N-2} and R_{N-3} are the same or different and are selected from the group consisting of:

(a) -H,

(b) -C₁-C₆ alkyl optionally substituted with one substituent selected from the group consisting of:

(i) -OH, and

(ii) -NH₂,

5 (c) -C₁-C₆ alkyl optionally substituted with one, two, or three -F, -Cl, -Br, or -I,

(d) -C₃-C₇ cycloalkyl,

(e) -(C₁-C₂ alkyl)-(C₃-C₇ cycloalkyl),

(f) -(C₁-C₆ alkyl)-O-(C₁-C₃ alkyl),

10 (g) -C₂-C₆ alkenyl with one or two double bonds,

(h) -C₂-C₆ alkynyl with one or two triple

bonds,

(i) -C₁-C₆ alkyl chain with one double bond

15 and one triple bond,

(j) -R_{1-aryl} where R_{1-aryl} is as defined above, and

(k) -R_{1-heteroaryl} where R_{1-heteroaryl} is as defined above,

(8) -(CH₂)₀₋₄-CO-(C₁-C₁₂ alkyl),

20 (9) -(CH₂)₀₋₄-CO-(C₂-C₁₂ alkenyl with one, two or three double bonds),

(10) -(CH₂)₀₋₄-CO-(C₂-C₁₂ alkynyl with one, two or

three triple bonds),

(11) -(CH₂)₀₋₄-CO-(C₃-C₇ cycloalkyl),

25 (12) -(CH₂)₀₋₄-CO-R_{1-aryl} where R_{1-aryl} is as defined above,

(13) -(CH₂)₀₋₄-CO-R_{1-heteroaryl} where R_{1-heteroaryl} is as

defined above,

(14) -(CH₂)₀₋₄-CO-R_{1-heterocycle} where R_{1-heterocycle} is as

30 defined above,

(15) -(CH₂)₀₋₄-CO-R_{N-4} where R_{N-4} is selected from

the group consisting of morpholinyl, thiomorpholinyl, piperazinyl, piperidinyl, homomorpholinyl, homothiomorpholinyl, homothiomorpholinyl S-oxide, homothiomorpholinyl S,S-dioxide, pyrrolinyl and pyrrolidinyl where each group is

optionally substituted with one, two, three, or four of C₁-C₆ alkyl,

(16) $-(CH_2)_{0-4}-CO-O-R_{N-5}$ where R_{N-5} is

selected from the group consisting of:

(a) C₁-C₆ alkyl,

5 (b) $-(CH_2)_{0-2}-(R_{1-aryl})$ where R_{1-aryl} is as defined above,

(c) C₂-C₆ alkenyl containing one or two double bonds,

10 (d) C₂-C₆ alkynyl containing one or two triple bonds,

(e) C₃-C₇ cycloalkyl, and

(f) $-(CH_2)_{0-2}-(R_{1-heteroaryl})$ where R_{1-heteroaryl} is as defined above,

15 (17) $-(CH_2)_{0-4}-SO_2-NR_{N-2}R_{N-3}$ where R_{N-2} and R_{N-3} are as defined above,

(18) $-(CH_2)_{0-4}-SO-(C_1-C_8 \text{ alkyl})$,

(19) $-(CH_2)_{0-4}-SO_2-(C_1-C_{12} \text{ alkyl})$,

(20) $-(CH_2)_{0-4}-SO_2-(C_3-C_7 \text{ cycloalkyl})$,

20 (21) $-(CH_2)_{0-4}-N(H \text{ or } R_{N-5})-CO-O-R_{N-5}$ where R_{N-5} can be the same or different and is as defined above,

(22) $-(CH_2)_{0-4}-N(H \text{ or } R_{N-5})-CO-N(R_{N-5})_2$, where R_{N-5} can be the same or different and is as defined above,

(23) $-(CH_2)_{0-4}-N-CS-N(R_{N-5})_2$, where R_{N-5} can be the same or different and is as defined above,

25 (24) $-(CH_2)_{0-4}-N(-H \text{ or } R_{N-5})-CO-R_{N-2}$ where R_{N-5} and R_{N-2} can be the same or different and are as defined above,

(25) $-(CH_2)_{0-4}-NR_{N-2}R_{N-3}$ where R_{N-2} and R_{N-3} can be the same or different and are as defined above,

(26) $-(CH_2)_{0-4}-R_{N-4}$ where R_{N-4} is as defined above,

30 (27) $-(CH_2)_{0-4}-O-CO-(C_1-C_6 \text{ alkyl})$,

(28) $-(CH_2)_{0-4}-O-P(O)-(OR_{N-aryl-1})_2$ where R_{N-aryl-1} is – H or C₁-C₄ alkyl,

(29) $-(CH_2)_{0-4}-O-CO-N(R_{N-5})_2$ where R_{N-5} is as defined above,

- (30) $-(\text{CH}_2)_{0-4}-\text{O}-\text{CS}-\text{N}(\text{R}_{\text{N}-5})_2$ where $\text{R}_{\text{N}-5}$ is as defined above,
- (31) $-(\text{CH}_2)_{0-4}-\text{O}-(\text{R}_{\text{N}-5})_2$ where $\text{R}_{\text{N}-5}$ is as defined above,
- 5 (32) $-(\text{CH}_2)_{0-4}-\text{O}-(\text{R}_{\text{N}-5})_2-\text{COOH}$ where $\text{R}_{\text{N}-5}$ is as defined above,
- (33) $-(\text{CH}_2)_{0-4}-\text{S}-(\text{R}_{\text{N}-5})_2$ where $\text{R}_{\text{N}-5}$ is as defined above,
- (34) $-(\text{CH}_2)_{0-4}-\text{O}-(\text{C}_1-\text{C}_6 \text{ alkyl optionally substituted with one, two, three, four, or five of } -\text{F}),$
- 10 (35) $\text{C}_3-\text{C}_7 \text{ cycloalkyl},$
- (36) $\text{C}_2-\text{C}_6 \text{ alkenyl with one or two double bonds optionally substituted with } \text{C}_1-\text{C}_3 \text{ alkyl, } -\text{F, } -\text{Cl, } -\text{Br, } -\text{I, } -\text{OH, } -\text{SH, } -\text{C}\equiv\text{N, } -\text{CF}_3, \text{ C}_1-\text{C}_3 \text{ alkoxy, or } -\text{NR}_{1-\text{a}}\text{R}_{1-\text{b}} \text{ where } \text{R}_{1-\text{a}} \text{ and } \text{R}_{1-\text{b}} \text{ are as defined above,}$
- 15 (37) $\text{C}_2-\text{C}_6 \text{ alkynyl with one or two triple bonds optionally substituted with } \text{C}_1-\text{C}_3 \text{ alkyl, } -\text{F, } -\text{Cl, } -\text{Br, } -\text{I, } -\text{OH, } -\text{SH, } -\text{C}\equiv\text{N, } -\text{CF}_3, \text{ C}_1-\text{C}_3 \text{ alkoxy, or } -\text{NR}_{1-\text{a}}\text{R}_{1-\text{b}} \text{ where } \text{R}_{1-\text{a}} \text{ and } \text{R}_{1-\text{b}} \text{ are as defined above,}$
- (38) $-(\text{CH}_2)_{0-4}-\text{N}(-\text{H or } \text{R}_{\text{N}-5})-\text{SO}_2-\text{R}_{\text{N}-2}$ where $\text{R}_{\text{N}-5}$ and $\text{R}_{\text{N}-2}$ can be the same or different and are as described above, or
- 20 (39) $-(\text{CH}_2)_{0-4}-\text{C}_3-\text{C}_7 \text{ cycloalkyl},$
- (B) $-\text{R}_{\text{N-heteroaryl}}$ where $\text{R}_{\text{N-heteroaryl}}$ is selected from the group consisting of:
- pyridinyl,
- pyrimidinyl,
- 25 quinolinyl,
- benzothienyl,
- indolyl,
- indolinyl,
- pyridazinyl,
- 30 pyrazinyl,
- isoindolyl,
- isoquinolyl,
- quinazolinyl,

	quinoxaliny,
	phthalaziny,
	imidazolyl,
	isoxazolyl,
5	pyrazolyl,
	oxazolyl,
	thiazolyl,
	indoliziny,
	indazolyl,
10	benzothiazolyl,
	benzimidazolyl,
	benzofuranyl,
	furanyl,
	thienyl,
15	pyrrolyl,
	oxadiazolyl,
	thiadiazolyl,
	triazolyl,
	tetrazolyl,
20	oxazolopyridiny,
	imidazopyridiny,
	isothiazolyl,
	naphthyridiny,
	cinnoliny,
25	carbazolyl,
	beta-carboliny,
	isochromanyl,
	chromanyl,
	tetrahydroisoquinoliny,
30	isoindoliny,
	isobenzotetrahydrofuranyl,
	isobenzotetrahydrothienyl,
	isobenzothienyl,
	benzoxazolyl,

pyridopyridinyl,
benzotetrahydrofuranyl,
benzotetrahydrothienyl,
purinyl,
5 benzodioxolyl,
triazinyl,
phenoxazinyl,
phenothiazinyl,
pteridinyl,
10 benzothiazolyl,
imidazopyridinyl,
imidazothiazolyl,
dihydrobenzisoaxazinyl,
benzisoaxazinyl,
15 benzoxazinyl,
dihydrobenzisothiazinyl,
benzopyranyl,
benzothiopyranyl,
coumarinyl,
20 isocoumarinyl,
chromonyl,
chromanonyl,
pyridinyl-N-oxide,
tetrahydroquinolinyl,
25 dihydroquinolinyl,
dihydroquinolinonyl,
dihydroisoquinolinonyl,
dihydrocoumarinyl,
dihydroisocoumarinyl,
30 isoindolinonyl,
benzodioxanyl,
benzoxazolinonyl,
pyrrolyl N-oxide,
pyrimidinyl N-oxide,

- pyridazinyl N-oxide,
 pyrazinyl N-oxide,
 quinolinyl N-oxide,
 indolyl N-oxide,
 5 indoliny N-oxide,
 isoquinolyl N-oxide,
 quinazolinyl N-oxide,
 quinoxaliny N-oxide,
 phthalazinyl N-oxide,
 10 imidazolyl N-oxide,
 isoxazolyl N-oxide,
 oxazolyl N-oxide,
 thiazolyl N-oxide,
 indoliziny N-oxide,
 15 indazolyl N-oxide,
 benzothiazolyl N-oxide,
 benzimidazolyl N-oxide,
 pyrrolyl N-oxide,
 oxadiazolyl N-oxide,
 20 thiadiazolyl N-oxide,
 triazolyl N-oxide,
 tetrazolyl N-oxide,
 benzothiopyranyl S-oxide, and
 benzothiopyranyl S,S-dioxide
- 25 where the $R_{N\text{-heteroaryl}}$ group is bonded by any atom of the
 parent $R_{N\text{-heteroaryl}}$ group substituted by hydrogen such that the new bond to the $R_{N\text{-heteroaryl}}$ group replaces the hydrogen atom and its bond, where heteroaryl is
 optionally substituted with one, two, three, or four of:
- (1) $C_1\text{-}C_6$ alkyl, optionally substituted with one, two or
 30 three substituents selected from the group consisting of $C_1\text{-}C_3$ alkyl, -F, -Cl, -Br, -I,
 -OH, -SH, $-C\equiv N$, $-CF_3$, $C_1\text{-}C_3$ alkoxy, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as
 defined above,
- (2) -OH,

- (3) $-\text{NO}_2$,
 (4) $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, or $-\text{I}$,
 (5) $-\text{CO}-\text{OH}$,
 (6) $-\text{C}\equiv\text{N}$,
 5 (7) $-(\text{CH}_2)_{0-4}-\text{CO}-\text{NR}_{\text{N}-2}\text{R}_{\text{N}-3}$ where $\text{R}_{\text{N}-2}$ and $\text{R}_{\text{N}-3}$ are the same or different and are selected from the group consisting of:
 (a) $-\text{H}$,
 (b) $-\text{C}_1-\text{C}_6$ alkyl optionally substituted with one substituent selected from the group consisting of:
 10 (i) $-\text{OH}$, and
 (ii) $-\text{NH}_2$,
 (c) $-\text{C}_1-\text{C}_6$ alkyl optionally substituted with one, two, or three $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$,
 (d) $-\text{C}_3-\text{C}_7$ cycloalkyl,
 15 (e) $-(\text{C}_1-\text{C}_2 \text{ alkyl})-(\text{C}_3-\text{C}_7 \text{ cycloalkyl})$,
 (f) $-(\text{C}_1-\text{C}_6 \text{ alkyl})-\text{O}-(\text{C}_1-\text{C}_3 \text{ alkyl})$,
 (g) $-\text{C}_2-\text{C}_6$ alkenyl with one or two double bonds,
 (h) $-\text{C}_2-\text{C}_6$ alkynyl with one or two triple bonds,
 20 (i) $-\text{C}_1-\text{C}_6$ alkyl chain with one double bond and one triple bond,
 (j) $-\text{R}_{1-\text{aryl}}$ where $\text{R}_{1-\text{aryl}}$ is as defined above,
 (k) $-\text{R}_{1-\text{heteroaryl}}$ where $\text{R}_{1-\text{heteroaryl}}$ is as defined above,
 25 above,
 (8) $-(\text{CH}_2)_{0-4}-\text{CO}-(\text{C}_1-\text{C}_{12} \text{ alkyl})$,
 (9) $-(\text{CH}_2)_{0-4}-\text{CO}-(\text{C}_2-\text{C}_{12} \text{ alkenyl with one, two or three double bonds})$,
 (10) $-(\text{CH}_2)_{0-4}-\text{CO}-(\text{C}_2-\text{C}_{12} \text{ alkynyl with one, two or three triple bonds})$,
 30 (11) $-(\text{CH}_2)_{0-4}-\text{CO}-(\text{C}_3-\text{C}_7 \text{ cycloalkyl})$,
 (12) $-(\text{CH}_2)_{0-4}-\text{CO}-\text{R}_{1-\text{aryl}}$ where $\text{R}_{1-\text{aryl}}$ is as defined above,

(13) $-(\text{CH}_2)_{0-4}-\text{CO}-\text{R}_{1-\text{heteroaryl}}$ where $\text{R}_{1-\text{heteroaryl}}$ is as defined above,

(14) $-(\text{CH}_2)_{0-4}-\text{CO}-\text{R}_{1-\text{heterocycle}}$ where $\text{R}_{1-\text{heterocycle}}$ is as defined above,

5 (15) $-(\text{CH}_2)_{0-4}-\text{CO}-\text{R}_{\text{N-4}}$ where $\text{R}_{\text{N-4}}$ is selected from the group consisting of morpholinyl, thiomorpholinyl, piperazinyl, piperidinyl, homomorpholinyl, homothiomorpholinyl, homothiomorpholinyl S-oxide, homothiomorpholinyl S,S-dioxide, pyrrolinyl and pyrrolidinyl where each group is optionally substituted with one, two, three, or four of C_1-C_6 alkyl,

10 (16) $-(\text{CH}_2)_{0-4}-\text{CO}-\text{O}-\text{R}_{\text{N-5}}$ where $\text{R}_{\text{N-5}}$ is selected from the group consisting of:

(a) C_1-C_6 alkyl,

(b) $-(\text{CH}_2)_{0-2}-(\text{R}_{1-\text{aryl}})$ where $\text{R}_{1-\text{aryl}}$ is as defined above,

15 (c) C_2-C_6 alkenyl containing one or two double bonds,

(d) C_2-C_6 alkynyl containing one or two triple bonds,

(e) C_3-C_7 cycloalkyl, and

20 (f) $-(\text{CH}_2)_{0-2}-(\text{R}_{1-\text{heteroaryl}})$ where $\text{R}_{1-\text{heteroaryl}}$ is as defined above,

(17) $-(\text{CH}_2)_{0-4}-\text{SO}_2-\text{NR}_{\text{N-2}}\text{R}_{\text{N-3}}$ where $\text{R}_{\text{N-2}}$ and $\text{R}_{\text{N-3}}$ are as defined above,

(18) $-(\text{CH}_2)_{0-4}-\text{SO}-(\text{C}_1-\text{C}_8 \text{ alkyl})$,

25 (19) $-(\text{CH}_2)_{0-4}-\text{SO}_2-(\text{C}_1-\text{C}_{12} \text{ alkyl})$,

(20) $-(\text{CH}_2)_{0-4}-\text{SO}_2-(\text{C}_3-\text{C}_7 \text{ cycloalkyl})$,

(21) $-(\text{CH}_2)_{0-4}-\text{N}(\text{H or } \text{R}_{\text{N-5}})-\text{CO}-\text{O}-\text{R}_{\text{N-5}}$ where $\text{R}_{\text{N-5}}$ can be the same or different and is as defined above,

30 (22) $-(\text{CH}_2)_{0-4}-\text{N}(\text{H or } \text{R}_{\text{N-5}})-\text{CO}-\text{N}(\text{R}_{\text{N-5}})_2$, where $\text{R}_{\text{N-5}}$ can be the same or different and is as defined above,

(23) $-(\text{CH}_2)_{0-4}-\text{N}-\text{CS}-\text{N}(\text{R}_{\text{N-5}})_2$, where $\text{R}_{\text{N-5}}$ can be the same or different and is as defined above,

(24) $-(\text{CH}_2)_{0-4}-\text{N}(-\text{H or } \text{R}_{\text{N-5}})-\text{CO}-\text{R}_{\text{N-2}}$ where $\text{R}_{\text{N-5}}$ and $\text{R}_{\text{N-2}}$ can be the same or different and are as defined above,

(25) $-(CH_2)_{0-4}-NR_{N-2}R_{N-3}$ where R_{N-2} and R_{N-3} can be the same or different and are as defined above,

(26) $-(CH_2)_{0-4}-R_{N-4}$ where R_{N-4} is as defined above,

(27) $-(CH_2)_{0-4}-O-CO-(C_1-C_6 \text{ alkyl})$,

5 (28) $-(CH_2)_{0-4}-O-P(O)-(OR_{N-aryl-1})_2$ where $R_{N-aryl-1}$ is – H or C_1-C_4 alkyl,

(29) $-(CH_2)_{0-4}-O-CO-N(R_{N-5})_2$ where R_{N-5} is as defined above,

10 (30) $-(CH_2)_{0-4}-O-CS-N(R_{N-5})_2$ where R_{N-5} is as defined above,

(31) $-(CH_2)_{0-4}-O-(R_{N-5})_2$ where R_{N-5} is as defined above,

(32) $-(CH_2)_{0-4}-O-(R_{N-5})_2-COOH$ where R_{N-5} is as defined above,

15 (33) $-(CH_2)_{0-4}-S-(R_{N-5})_2$ where R_{N-5} is as defined above,

(34) $-(CH_2)_{0-4}-O-(C_1-C_6 \text{ alkyl optionally substituted with one, two, three, four, or five of } -F)$,

(35) C_3-C_7 cycloalkyl,

20 (36) C_2-C_6 alkenyl with one or two double bonds optionally substituted with C_1-C_3 alkyl, $-F$, $-Cl$, $-Br$, $-I$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, C_1-C_3 alkoxy, or $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

(37) C_2-C_6 alkynyl with one or two triple bonds optionally substituted with C_1-C_3 alkyl, $-F$, $-Cl$, $-Br$, $-I$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, C_1-C_3 alkoxy, or $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

25 (38) $-(CH_2)_{0-4}-N(-H \text{ or } R_{N-5})-SO_2-R_{N-2}$ where R_{N-5} and R_{N-2} can be the same or different and are as defined above, or

(39) $-(CH_2)_{0-4}-C_3-C_7$ cycloalkyl,

30 (C) $R_{N-aryl}-W-R_{N-aryl}$, where R_{N-aryl} can be the same or different,

(D) $R_{N-aryl}-W-R_{N-heteroaryl}$,

(E) $R_{N-aryl}-W-R_{N-1-heterocycle}$, wherein $R_{N-1-heterocycle}$ is the same as $R_{1-heterocycle}$, and $R_{1-heterocycle}$ is as defined above

(F) $R_{N\text{-heteroaryl}}\text{-}W\text{-}R_{N\text{-aryl}}$,

(G) $R_{N\text{-heteroaryl}}\text{-}W\text{-}R_{N\text{-heteroaryl}}$,

(H) $R_{N\text{-heteroaryl}}\text{-}W\text{-}R_{N\text{-1-heterocycle}}$,

(I) $R_{N\text{-heterocycle}}\text{-}W\text{-}R_{N\text{-aryl}}$, wherein $R_{N\text{-heterocycle}}$ is the same as
 5 $R_{1\text{-heterocycle}}$, and $R_{1\text{-heterocycle}}$ is as defined above, and $R_{N\text{-aryl}}$ is as defined above,

(J) $R_{N\text{-heterocycle}}\text{-}W\text{-}R_{N\text{-heteroaryl}}$, and

(K) $R_{N\text{-heterocycle}}\text{-}W\text{-}R_{N\text{-1-heterocycle}}$,

where W is

(9) $\text{-(CH}_2\text{)}_{0-4}\text{-}$,

10 (10) -O- ,

(11) $\text{-S(O)}_{0-2}\text{-}$,

(12) $\text{-N(R}_{N-5}\text{)-}$ where R_{N-5} is as defined

above, or

(5) -CO- ;

15 (II) $\text{-CO-(C}_1\text{-C}_{10}\text{ alkyl)}$ where alkyl is optionally substituted with one
 three substituents selected from the group consisting of:

(A) -OH ,

(B) $\text{-C}_1\text{-C}_6\text{ alkoxy}$,

(C) $\text{-C}_1\text{-C}_6\text{ thioalkoxy}$,

20 (D) -CO-O-R_{N-8} where R_{N-8} is -H , $\text{C}_1\text{-C}_6\text{ alkyl}$ or -phenyl ,

(E) $\text{-CO-NR}_{N-2}\text{R}_{N-3}$ where R_{N-2} and R_{N-3} are the same or

different and are as defined above,

(F) -CO-R_{N-4} where R_{N-4} is as defined above,

(G) $\text{-SO}_2\text{-(C}_1\text{-C}_8\text{ alkyl)}$,

25 (H) $\text{-SO}_2\text{-NR}_{N-2}\text{R}_{N-3}$ where R_{N-2} and R_{N-3} are the same or
 different and are as defined above,

(I) $\text{-NH-CO-(C}_1\text{-C}_6\text{ alkyl)}$,

(J) -NH-CO-O-R_{N-8} where R_{N-8} is as defined above,

(K) $\text{-NR}_{N-2}\text{R}_{N-3}$ where R_{N-2} and R_{N-3} are the same or different

30 and are as defined above,

(L) -R_{N-4} where R_{N-4} is as defined above,

(M) $\text{-O-CO-(C}_1\text{-C}_6\text{ alkyl)}$,

(N) $\text{-O-CO-NR}_{N-8}\text{R}_{N-8}$ where R_{N-8} are the same or different

and are as defined above,

(O) -O-(C₁-C₅ alkyl)-COOH,

(P) -O-(C₁-C₆ alkyl optionally substituted with one, two, or three of -F, -Cl, -Br, or -I),

(Q) -NH-SO₂-(C₁-C₆ alkyl), and

5 (R) -F, or -Cl,

(III) -CO-(C₁-C₆ alkyl)-O-(C₁-C₆ alkyl) where alkyl is optionally substituted with one, two, or three substituents selected from the group consisting of:

(A) -OH,

10 (B) -C₁-C₆ alkoxy,

(C) -C₁-C₆ thioalkoxy,

(D) -CO-O-R_{N-8} where R_{N-8} is -H, C₁-C₆ alkyl or -phenyl,

(E) -CO-NR_{N-2}R_{N-3} where R_{N-2} and R_{N-3} are the same or different and are as defined above,

15 (F) -CO-R_{N-4} where R_{N-4} is as defined above,

(G) -SO₂-(C₁-C₈ alkyl),

(H) -SO₂-NR_{N-2}R_{N-3} where R_{N-2} and R_{N-3} are the same or different and are as defined above,

(I) -NH-CO-(C₁-C₆ alkyl),

20 (J) -NH-CO-O-R_{N-8} where R_{N-8} is as defined above,

(K) -NR_{N-2}R_{N-3} where R_{N-2} and R_{N-3} are the same or different and are as defined above,

(L) -R_{N-4} where R_{N-4} is as defined above,

(M) -O-CO-(C₁-C₆ alkyl),

25 (N) -O-CO-NR_{N-8}R_{N-8} where R_{N-8} are the same or different and are as defined above,

(O) -O-(C₁-C₅ alkyl)-COOH,

(P) -O-(C₁-C₆ alkyl optionally substituted with one, two, or three of -F, -Cl, -Br, or -I),

30 (Q) -NH-SO₂-(C₁-C₆ alkyl), and

(R) -F, or -Cl,

(IV) $-\text{CO}-(\text{C}_1\text{-C}_6 \text{ alkyl})-\text{S}-(\text{C}_1\text{-C}_6 \text{ alkyl})$ where alkyl is optionally substituted with one, two, or three of substituents selected from the group consisting of:

- 5 (A) $-\text{OH}$,
 (B) $-\text{C}_1\text{-C}_6 \text{ alkoxy}$,
 (C) $-\text{C}_1\text{-C}_6 \text{ thioalkoxy}$,
 (D) $-\text{CO}-\text{O}-\text{R}_{\text{N-8}}$ where $\text{R}_{\text{N-8}}$ is as defined above,
 (E) $-\text{CO}-\text{NR}_{\text{N-2}}\text{R}_{\text{N-3}}$ where $\text{R}_{\text{N-2}}$ and $\text{R}_{\text{N-3}}$ are the same or different and are as defined above,
 10 (F) $-\text{CO}-\text{R}_{\text{N-4}}$ where $\text{R}_{\text{N-4}}$ is as defined above,
 (G) $-\text{SO}_2-(\text{C}_1\text{-C}_8 \text{ alkyl})$,
 (H) $-\text{SO}_2-\text{NR}_{\text{N-2}}\text{R}_{\text{N-3}}$ where $\text{R}_{\text{N-2}}$ and $\text{R}_{\text{N-3}}$ are the same or different and are as defined above,
 (I) $-\text{NH}-\text{CO}-(\text{C}_1\text{-C}_6 \text{ alkyl})$,
 15 (J) $-\text{NH}-\text{CO}-\text{O}-\text{R}_{\text{N-8}}$ where $\text{R}_{\text{N-8}}$ is as defined above,
 (K) $-\text{NR}_{\text{N-2}}\text{R}_{\text{N-3}}$ where $\text{R}_{\text{N-2}}$ and $\text{R}_{\text{N-3}}$ are the same or different and are as defined above,
 (L) $-\text{R}_{\text{N-4}}$ where $\text{R}_{\text{N-4}}$ is as defined above,
 (M) $-\text{O}-\text{CO}-(\text{C}_1\text{-C}_6 \text{ alkyl})$,
 20 (N) $-\text{O}-\text{CO}-\text{NR}_{\text{N-8}}\text{R}_{\text{N-8}}$ where $\text{R}_{\text{N-8}}$ are the same or different and are as defined above,
 (O) $-\text{O}-(\text{C}_1\text{-C}_5 \text{ alkyl})-\text{COOH}$,
 (P) $-\text{O}-(\text{C}_1\text{-C}_6 \text{ alkyl})$ optionally substituted with one, two, or three of $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, or $-\text{I}$,
 25 (Q) $-\text{NH}-\text{SO}_2-(\text{C}_1\text{-C}_6 \text{ alkyl})$, and
 (R) $-\text{F}$, or $-\text{Cl}$,
 (V) $-\text{CO}-\text{CH}(-(\text{CH}_2)_{0-2}-\text{O}-\text{R}_{\text{N-10}})-(\text{CH}_2)_{0-2}-\text{R}_{\text{N-aryl}}/\text{R}_{\text{N-heteroaryl}}$ where $\text{R}_{\text{N-aryl}}$ and $\text{R}_{\text{N-heteroaryl}}$ are as defined above, where $\text{R}_{\text{N-10}}$ is selected from the group consisting of:
 30 (A) $-\text{H}$,
 (B) $\text{C}_1\text{-C}_6 \text{ alkyl}$,
 (C) $\text{C}_3\text{-C}_7 \text{ cycloalkyl}$,
 (D) $\text{C}_2\text{-C}_6 \text{ alkenyl}$ with one double bond,

- (E) C₂-C₆ alkynyl with one triple bond,
 (F) R_{1-aryl} where R_{1-aryl} is as defined above, and
 (G) R_{N-heteroaryl} where R_{N-heteroaryl} is as defined above, or
 (VI) -CO-(C₃-C₈ cycloalkyl) where alkyl is optionally substituted
 5 with one or two substituents selected from the group consisting of:
 (A) -(CH₂)₀₋₄-OH,
 (B) -(CH₂)₀₋₄-C₁-C₆ alkoxy,
 (C) -(CH₂)₀₋₄-C₁-C₆ thioalkoxy,
 (D) -(CH₂)₀₋₄-CO-O-R_{N-8} where R_{N-8} is -H, C₁-C₆ alkyl or
 10 phenyl,
 (E) -(CH₂)₀₋₄-CO-NR_{N-2}R_{N-3} where R_{N-2} and R_{N-3} are the
 same or different and are as defined above,
 (F) -(CH₂)₀₋₄-CO-R_{N-4} where R_{N-4} is as defined above,
 (G) -(CH₂)₀₋₄-SO₂-(C₁-C₈ alkyl),
 15 (H) -(CH₂)₀₋₄-SO₂-NR_{N-2}R_{N-3} where R_{N-2} and R_{N-3} are the
 same or different and are as defined above,
 (I) -(CH₂)₀₋₄-NH-CO-(C₁-C₆ alkyl),
 (J) -NH-CO-O-R_{N-8} where R_{N-8} is as defined above,
 (K) -(CH₂)₀₋₄-NR_{N-2}R_{N-3} where R_{N-2} and R_{N-3} are the same or
 20 different and are as defined above,
 (L) -(CH₂)₀₋₄-R_{N-4} where R_{N-4} is as defined above,
 (M) -O-CO-(C₁-C₆ alkyl),
 (N) -O-CO-NR_{N-8}R_{N-8} where R_{N-8} are the same or different
 and are as defined above,
 25 (O) -O-(C₁-C₅ alkyl)-COOH,
 (P) -O-(C₁-C₆ alkyl optionally substituted with one, two, or
 three of -F, -Cl, -Br, or -I),
 (Q) -NH-SO₂-(C₁-C₆ alkyl), and
 (R) -F, or -Cl;

30

where R_A is:

(I)-C₁-C₁₀ alkyl optionally substituted with one, two or three substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above, -OC=O NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above, -S(=O)₀₋₂ R_{1-a} where R_{1-a} is as defined above, -NR_{1-a}C=O NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above, -C=O NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above, and -S(=O)₂ NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(II) -(CH₂)₀₋₃-(C₃-C₈) cycloalkyl where cycloalkyl can be optionally substituted with one, two or three substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, -CO-OH, -CO-O-(C₁-C₄ alkyl), and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(III) -(CR_{A-x}R_{A-y})₀₋₄-R_{A-aryl} where R_{A-x} and R_{A-y} are

- (A) -H,
- (B) C₁-C₄ alkyl optionally substituted with one or two -OH,
- (C) C₁-C₄ alkoxy optionally substituted with one, two, or three of -F,
- (D) -(CH₂)₀₋₄-C₃-C₇ cycloalkyl,
- (E) C₂-C₆ alkenyl containing one or two double bonds,
- (F) C₂-C₆ alkynyl containing one or two triple bonds, or
- (G) phenyl,

and where R_{A-x} and R_{A-y} are taken together with the carbon to which they are attached to form a carbocycle of three, four, five, six or seven carbon atoms, optionally where one carbon atom is replaced by a heteroatom selected from the group consisting of -O-, -S-, -SO₂-, and -NR_{N-2}- and R_{A-aryl} is the same as R_{N-aryl},

(IV) -(CR_{A-x}R_{A-y})₀₋₄-R_{A-heteroaryl} where R_{A-heteroaryl} is the same as R_{N-heteroaryl} and R_{A-x} and R_{A-y} are as defined above,

(V) -(CR_{A-x}R_{A-y})₀₋₄-R_{A-aryl}-R_{A-aryl} where R_{A-aryl}, R_{A-x} and R_{A-y} are as defined above,

(VI) -(CR_{A-x}R_{A-y})₀₋₄-R_{A-aryl}-R_{A-heteroaryl} where R_{A-aryl}, R_{A-heteroaryl}, R_{A-x} and R_{A-y} are as defined above,

(VII) -(CR_{A-x}R_{A-y})₀₋₄-R_{A-heteroaryl}-R_{A-aryl} where R_{A-heteroaryl}, R_{A-aryl}, R_{A-x} and R_{A-y} are as defined above,

(VIII) $-(\text{CR}_{\text{A-x}}\text{R}_{\text{A-y}})_{0-4}-\text{R}_{\text{A-heteroaryl}}-\text{R}_{\text{A-heteroaryl}}$ where $\text{R}_{\text{A-heteroaryl}}$, $\text{R}_{\text{A-x}}$ and $\text{R}_{\text{A-y}}$ are as defined above,

(IX) $-(\text{CR}_{\text{A-x}}\text{R}_{\text{A-y}})_{0-4}-\text{R}_{\text{A-aryl}}-\text{R}_{\text{A-heterocycle}}$ where $\text{R}_{\text{A-heterocycle}}$ is defined as $\text{R}_{1\text{-heterocycle}}$, and where $\text{R}_{\text{A-aryl}}$, $\text{R}_{\text{A-x}}$ and $\text{R}_{\text{A-y}}$ are as defined above,

5 (X) $-(\text{CR}_{\text{A-x}}\text{R}_{\text{A-y}})_{0-4}-\text{R}_{\text{A-heteroaryl}}-\text{R}_{\text{A-heterocycle}}$ where $\text{R}_{\text{A-heteroaryl}}$, $\text{R}_{\text{A-heterocycle}}$, $\text{R}_{\text{A-x}}$ and $\text{R}_{\text{A-y}}$ are as defined above,

(XI) $-(\text{CR}_{\text{A-x}}\text{R}_{\text{A-y}})_{0-4}-\text{R}_{\text{A-heterocycle}}-\text{R}_{\text{A-aryl}}$ where $\text{R}_{\text{A-heterocycle}}$, $\text{R}_{\text{A-aryl}}$, $\text{R}_{\text{A-x}}$ and $\text{R}_{\text{A-y}}$ are as defined above,

10 (XII) $-(\text{CR}_{\text{A-x}}\text{R}_{\text{A-y}})_{0-4}-\text{R}_{\text{A-heterocycle}}-\text{R}_{\text{A-heteroaryl}}$ where $\text{R}_{\text{A-heterocycle}}$, $\text{R}_{\text{A-heteroaryl}}$, $\text{R}_{\text{A-x}}$ and $\text{R}_{\text{A-y}}$ are as defined above,

(XIII) $-(\text{CR}_{\text{A-x}}\text{R}_{\text{A-y}})_{0-4}-\text{R}_{\text{A-heterocycle}}-\text{R}_{\text{A-heterocycle}}$ where $\text{R}_{\text{A-heterocycle}}$, $\text{R}_{\text{A-x}}$ and $\text{R}_{\text{A-y}}$ are as defined above,

(XIV) $-(\text{CR}_{\text{A-x}}\text{R}_{\text{A-y}})_{0-4}-\text{R}_{\text{A-heterocycle}}$ where $\text{R}_{\text{A-heterocycle}}$, $\text{R}_{\text{A-x}}$ and $\text{R}_{\text{A-y}}$ are as defined above,

15 (XV) $-[\text{C}(\text{R}_{\text{A-1}})(\text{R}_{\text{A-2}})]_{1-3}-\text{CO}-\text{N}-(\text{R}_{\text{A-3}})_2$ where $\text{R}_{\text{A-1}}$ and $\text{R}_{\text{A-2}}$ are the same or different and are selected from the group consisting of:

(A) -H,

(B) $\text{C}_1\text{-C}_6$ alkyl, optionally substituted with one, two or three substituents selected from the group consisting of $\text{C}_1\text{-C}_3$ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C \equiv N, -CF $_3$, $\text{C}_1\text{-C}_6$ alkoxy, -O-phenyl, and -NR $_{1-a}$ R $_{1-b}$ where R $_{1-a}$ and R $_{1-b}$ are as defined above,

20

(C) $\text{C}_2\text{-C}_6$ alkenyl with one or two double bonds, optionally substituted with one, two or three substituents selected from the group consisting of $\text{C}_1\text{-C}_3$ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C \equiv N, -CF $_3$, $\text{C}_1\text{-C}_6$ alkoxy, -O-phenyl, and -NR $_{1-a}$ R $_{1-b}$ where R $_{1-a}$ and R $_{1-b}$ are as defined above,

25

(D) $\text{C}_2\text{-C}_6$ alkynyl with one or two triple bonds, optionally substituted with one, two or three substituents selected from the group consisting of $\text{C}_1\text{-C}_3$ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C \equiv N, -CF $_3$, $\text{C}_1\text{-C}_6$ alkoxy, -O-phenyl, and -NR $_{1-a}$ R $_{1-b}$ where R $_{1-a}$ and R $_{1-b}$ are as defined above,

30 (E) $-(\text{CH}_2)_{1-2}-\text{S}(\text{O})_{0-2}-(\text{C}_1\text{-C}_6 \text{ alkyl})$,

(F) $-(\text{CH}_2)_{0-4}-\text{C}_3\text{-C}_7$ cycloalkyl, optionally substituted with one, two or three substituents selected from the group consisting of $\text{C}_1\text{-C}_3$ alkyl, -F,

-Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(G) -(C₁-C₄ alkyl)-R_{A'-aryl} where R_{A'-aryl} is as defined for R_{1-a}

aryl,

5 (H) -(C₁-C₄ alkyl)-R_{A-heteroaryl} where R_{A-heteroaryl} is as defined above,

(I) -(C₁-C₄ alkyl)-R_{A-heterocycle} where R_{A-heterocycle} is as defined above,

(J) -R_{A-heteroaryl} where R_{A-heteroaryl} is as defined above,

10 (K) -R_{A-heterocycle} where R_{A-heterocycle} is as defined above,

(M) -(CH₂)₁₋₄-R_{A-4}-(CH₂)₀₋₄-R_{A'-aryl} where R_{A-4} is -O-, -S- or -NR_{A-5}- where R_{A-5} is C₁-C₆ alkyl, and where R_{A'-aryl} is defined above,

(N) -(CH₂)₁₋₄-R_{A-4}-(CH₂)₀₋₄-R_{A-heteroaryl} where R_{A-4} and R_{A-heteroaryl} are as defined above, and

15 (O) -R_{A'-aryl} where R_{A'-aryl} is as defined above,

and where R_{A-3} is the same or different and is:

(A) -H,

(B) -C₁-C₆ alkyl optionally substituted with one, two or three substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(C) C₂-C₆ alkenyl with one or two double bonds, optionally substituted with one, two or three substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(D) C₂-C₆ alkynyl with one or two triple bonds, optionally substituted with one, two or three substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

30 (E) -(CH₂)₀₋₄-C₃-C₇ cycloalkyl, optionally substituted with one, two or three substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(F) $-R_{A'-aryl}$ where $R_{A'-aryl}$ is as defined above,

(G) $-R_{A-heteroaryl}$ where $R_{A-heteroaryl}$ is as defined above,

(H) $-R_{A-heterocycle}$ where $R_{A-heterocycle}$ is as defined above,

(I) $-(C_1-C_4 \text{ alkyl})-R_{A'-aryl}$ where $R_{A'-aryl}$ is as defined
5 above,

(J) $-(C_1-C_4 \text{ alkyl})-R_{A-heteroaryl}$ where $R_{A-heteroaryl}$ is as defined
above,

(K) $-(C_1-C_4 \text{ alkyl})-R_{A-heterocycle}$ where $R_{A-heterocycle}$ is as defined
above, or

(XVI) $-\text{CH}(R_{A-aryl})_2$ where R_{A-aryl} are the same or different and are as
10 defined above,

(XVII) $-\text{CH}(R_{A-heteroaryl})_2$ where $R_{A-heteroaryl}$ are the same or different
and are as defined above,

(XVIII) $-\text{CH}(R_{A-aryl})(R_{A-heteroaryl})$ where R_{A-aryl} and $R_{A-heteroaryl}$ are as
15 defined above,

(XIX) -cyclopentyl, -cyclohexyl, or -cycloheptyl ring fused to R_{A-aryl} ,
 $R_{A-heteroaryl}$, $R_{A-heterocycle}$ where R_{A-aryl} or $R_{A-heteroaryl}$ or $R_{A-heterocycle}$ are as defined above
where one carbon of cyclopentyl, cyclohexyl, or -cycloheptyl is optionally replaced
with NH, NR_{N-5} , O, or $\text{S}(=\text{O})_{0-2}$, and where cyclopentyl, cyclohexyl, or -cycloheptyl
20 can be optionally substituted with one or two $-C_1-C_3$ alkyl, -F, -OH, -SH, $-C\equiv N$, -
 CF_3 , C_1-C_6 alkoxy, =O, or $-\text{NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

(XX) C_2-C_{10} alkenyl containing one or two double bonds optionally
substituted with one, two or three substituents selected from the group consisting of
 C_1-C_3 alkyl, -F, -Cl, -Br, -I, -OH, -SH, $-C\equiv N$, $-\text{CF}_3$, C_1-C_6 alkoxy, -O-phenyl, and -
25 $\text{NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

(XXI) C_2-C_{10} alkynyl containing one or two triple bonds optionally
substituted with one, two or three substituents selected from the group consisting of
 C_1-C_3 alkyl, -F, -Cl, -Br, -I, -OH, -SH, $-C\equiv N$, $-\text{CF}_3$, C_1-C_6 alkoxy, -O-phenyl, $-\text{NR}_{1-a}\text{R}_{1-b}$
30 R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(XXI) $-(\text{CH}_2)_{0-1}-\text{CHR}_{A-6}-(\text{CH}_2)_{0-1}-R_{A-aryl}$ where R_{A-aryl} is as defined
above and R_{A-6} is $-(\text{CH}_2)_{0-6}-\text{OH}$,

(XXII) $-(\text{CH}_2)_{0-1}-\text{CHR}_{A-6}-(\text{CH}_2)_{0-1}-R_{A-heteroaryl}$ where $R_{A-heteroaryl}$ and
 R_{A-6} is as defined above,

(XXIII) $-\text{CH}(-\text{R}_{\text{A-aryl}} \text{ or } \text{R}_{\text{A-heteroaryl}})-\text{CO}-\text{O}(\text{C}_1-\text{C}_4 \text{ alkyl})$ where $\text{R}_{\text{A-aryl}}$

and $\text{R}_{\text{A-heteroaryl}}$ are as defined above,

(XXIV) $-\text{CH}(-\text{CH}_2-\text{OH})-\text{CH}(-\text{OH})-\text{micro-NO}_2$,

(XXV) $(\text{C}_1-\text{C}_6 \text{ alkyl})-\text{O}-(\text{C}_1-\text{C}_6 \text{ alkyl})-\text{OH}$,

5 (XXVII) $-\text{CH}_2-\text{NH}-\text{CH}_2-\text{CH}(-\text{O}-\text{CH}_2-\text{CH}_3)_2$,

(XXVIII) $-\text{H}$,

(XXIX) $-(\text{CH}_2)_{0-6}-\text{C}(=\text{NR}_{1-a})(\text{NR}_{1-a}\text{R}_{1-b})$ where R_{1-a} and R_{1-b} are as defined above; or

(XXX)

10 $-\text{C}=\text{OC}(\text{HR}_6)\text{NHR}_7$, where R_6 and R_7 are as defined below,

$-\text{C}=\text{OR}_7$, where R_7 is as defined below,

$-\text{C}=\text{OOR}_7$, where R_7 is as defined below, or

$-\text{SOOR}_7$ where R_7 is as defined below,

wherein R_6 is:

15 hydrogen,

$\text{C}_1 - \text{C}_3$ alkyl,

phenyl,

thioalkoxyalkyl,

alkyl substituted aryl,

20 cycloalkyl,

cycloalkylalkyl,

hydroxyalkyl,

alkoxyalkyl,

aryloxyalkyl,

25 haloalkyl,

carboxyalkyl,

alkoxycarbonylalkyl,

aminoalkyl,

(N-protected)aminoalkyl,

30 alkylaminoalkyl,

((N-protected)(alkyl)amino)alkyl,

dialkylaminoalkyl,

guanidinoalkyl,

lower alkenyl,

heterocyclic,
(heterocyclic)alkyl),
arylthioalkyl,
arylsulfonylalkyl,
5 (heterocyclic)thioalkyl,
(heterocyclic)sulfonylalkyl,
(heterocyclic)oxyalkyl,
arylalkoxyalkyl,
arylthioalkoxyalkyl,
10 arylalkylsulfonylalkyl,
(heterocyclic))alkoxyalkyl,
(heterocyclic)thioalkoxyalkyl,
(heterocyclic)alkylsulfonylalkyl,
cycloalkyloxyalkyl,
15 cycloalkylthioalkyl,
cycloalkylsulfonylalkyl,
cycloalkylalkoxyalkyl,
cycloalkylthioalkoxyalkyl,
cycloalkylalkylsulfonylalkyl,
20 aminocarbonyl,
alkylaminocarbonyl,
dialkylaminocarbonyl,
aroalkyl,
(heterocyclic)carbonylalkyl,
25 polyhydroxyalkyl,
aminocarbonylalkyl,
alkylaminocarbonylalkyl,
dialkylaminocarbonylalkyl,
aryloxyalkyl, or
30 alkylsulfonylalkyl,

wherein heterocyclic is pyridyl, thiazolyl, isothiazolyl, oxazolyl, isoxazolyl, furanyl, thienyl, tetrahydrofuranyl, tetrahydrothienyl and tetrahydro[2H]pyranyl and wherein the heterocycle is unsubstituted or substituted with one to three substituents independently selected from hydroxy, halo, amino,

alkylamino, dialkylamino, alkoxy, polyalkoxy, haloalkyl, cycloalkyl, cycloalkylalkyl, aryl, arylalkyl, COOH, -SO₃H, lower alkenyl or lower alkyl;

wherein R₇ is:

- 5 C₁ - C₃ alkyl,
phenyl,
thioalkoxyalkyl,
(aryl)alkyl,
cycloalkyl,
cycloalkylalkyl,
10 hydroxyalkyl,
alkoxyalkyl,
aryloxyalkyl,
haloalkyl,
carboxyalkyl,
15 alkoxycarbonylalkyl,
aminoalkyl,
(N-protected)aminocalkyl,
alkylaminoalkyl,
((N-protected)(alkyl)amino)alkyl,
20 dialkylaminoalkyl,
guanidinoalkyl,
lower alkenyl,
heterocyclic,
(heterocyclic)alkyl),
25 arylthioalkyl,
arylsulfonylalkyl,
(heterocyclic)thioalkyl,
(heterocyclic)sulfonylalkyl,
(heterocyclic)oxyalkyl,
30 arylalkoxyalkyl,
arylthioalkoxyalkyl,
arylalkylsulfonylalkyl,
(heterocyclic))alkoxyalkyl,
(heterocyclic)thioalkoxyalkyl,

(heterocyclic)alkylsulfonylalkyl,
 cycloalkyloxyalkyl,
 cycloalkylthioalkyl,
 cycloalkylsulfonylalkyl,
 5 cycloalkylalkoxyalkyl,
 cycloalkylthioalkoxyalkyl,
 cycloalkylalkylsulfonylalkyl,
 aminocarbonyl,
 alkylaminocarbonyl,
 10 dialkylaminocarbonyl,
 aroylalkyl,
 (heterocyclic)carbonylalkyl,
 polyhydroxyalkyl,
 aminocarbonylalkyl,
 15 alkylaminocarbonylalkyl,
 dialkylaminocarbonylalkyl,
 aryloxyalkyl, or
 alkylsulfonylalkyl,
 wherein heterocyclic is pyridyl, thiazolyl, isothiazolyl,
 20 oxazolyl, isoxazolyl, furanyl, thienyl, tetrahydrofuranyl, tetrahydrothienyl and
 tetrahydro[2H]pyranyl and wherein the heterocycle is unsubstituted or substituted
 with one to three substituents independently selected from hydroxy, halo, amino,
 alkylamino, dialkylamino, alkoxy, polyalkoxy, haloalkyl, cycloalkyl,
 cycloalkylalkyl, aryl, arylalkyl, COOH, -SO₃H, lower alkenyl or lower alkyl;

25

where R_B is:

(I)-C₁-C₁₀ alkyl optionally substituted with one, two or three
 substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH,
 -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as
 30 defined above, -OC=O NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above, -S(=O)₀₋₂
 R_{1-a} where R_{1-a} is as defined above, -NR_{1-a}C=O NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as
 defined above, -C=O NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above, and -
 S(=O)₂ NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(II) $-(CH_2)_{0-3}-(C_3-C_8)$ cycloalkyl where cycloalkyl can be optionally substituted with one, two or three substituents selected from the group consisting of C_1-C_3 alkyl, $-F$, $-Cl$, $-Br$, $-I$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, C_1-C_6 alkoxy, $-O$ -phenyl, $-CO-OH$, $-CO-O-(C_1-C_4 \text{ alkyl})$, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

5 (III) $-(CR_{B-x}R_{B-y})_{0-4}-R_{B-aryl}$ where R_{B-x} and R_{B-y} are

(A) $-H$,

(B) C_1-C_4 alkyl optionally substituted with one or two $-OH$,

(C) C_1-C_4 alkoxy optionally substituted with one, two, or
three of $-F$,

10 (D) $-(CH_2)_{0-4}-C_3-C_7$ cycloalkyl,

(E) C_2-C_6 alkenyl containing one or two double bonds,

(F) C_2-C_6 alkynyl containing one or two triple bonds, or

(G) phenyl,

and where R_{B-x} and R_{B-y} are taken together with the carbon to

15 which they are attached to form a carbocycle of three, four, five, six or seven carbon atoms, optionally where one carbon atom is replaced by a heteroatom selected from the group consisting of $-O-$, $-S-$, $-SO_2-$, and $-NR_{N-2}$ where R_{N-2} is as defined above, and R_{B-aryl} is the same as R_{N-aryl} and is defined above

(IV) $-(CR_{B-x}R_{B-y})_{0-4}-R_{B-heteroaryl}$ where $R_{B-heteroaryl}$ is the same as R_{N-}

20 heteroaryl, R_{B-x} , and R_{B-y} are as defined above,

(V) $-(CR_{B-x}R_{B-y})_{0-4}-R_{B-aryl}-R_{B-aryl}$ where R_{B-aryl} , R_{B-x} , and R_{B-y} are as defined above,

(VI) $-(CR_{B-x}R_{B-y})_{0-4}-R_{B-aryl}-R_{B-heteroaryl}$ where R_{B-aryl} , $R_{B-heteroaryl}$, R_{B-x} and R_{B-y} are as defined above,

25 (VII) $-(CR_{B-x}R_{B-y})_{0-4}-R_{B-heteroaryl}-R_{B-aryl}$ where $R_{B-heteroaryl}$, R_{B-aryl} , R_{B-x} and R_{B-y} are as defined above,

(VIII) $-(CR_{B-x}R_{B-y})_{0-4}-R_{B-heteroaryl}-R_{B-heteroaryl}$ where $R_{B-heteroaryl}$, R_{B-x} and R_{B-y} are as defined above,

(IX) $-(CR_{B-x}R_{B-y})_{0-4}-R_{B-aryl}-R_{B-heterocycle}$ where $R_{B-heterocycle}$ is defined as
30 $R_{1-heterocycle}$, and where R_{B-aryl} , R_{B-x} and R_{B-y} are as defined above,

(X) $-(CR_{B-x}R_{B-y})_{0-4}-R_{B-heteroaryl}-R_{B-heterocycle}$ where $R_{B-heteroaryl}$, $R_{B-heterocycle}$, R_{B-x} and R_{B-y} are as defined above,

(XI) $-(\text{CR}_{\text{B-x}}\text{R}_{\text{B-y}})_{0-4}-\text{R}_{\text{B-heterocycle}}-\text{R}_{\text{B-aryl}}$ where $\text{R}_{\text{B-heterocycle}}$, $\text{R}_{\text{B-aryl}}$, $\text{R}_{\text{B-x}}$ and $\text{R}_{\text{B-y}}$ are as defined above,

(XII) $-(\text{CR}_{\text{B-x}}\text{R}_{\text{B-y}})_{0-4}-\text{R}_{\text{B-heterocycle}}-\text{R}_{\text{B-heteroaryl}}$ where $\text{R}_{\text{B-heterocycle}}$, $\text{R}_{\text{B-heteroaryl}}$, $\text{R}_{\text{B-x}}$ and $\text{R}_{\text{B-y}}$ are as defined above,

5 (XIII) $-(\text{CR}_{\text{B-x}}\text{R}_{\text{B-y}})_{0-4}-\text{R}_{\text{B-heterocycle}}-\text{R}_{\text{B-heterocycle}}$ where $\text{R}_{\text{B-heterocycle}}$, $\text{R}_{\text{B-x}}$ and $\text{R}_{\text{B-y}}$ are as defined above,

(XIV) $-(\text{CR}_{\text{B-x}}\text{R}_{\text{B-y}})_{0-4}-\text{R}_{\text{B-heterocycle}}$ where $\text{R}_{\text{B-heterocycle}}$, $\text{R}_{\text{B-x}}$ and $\text{R}_{\text{B-y}}$ are as defined above,

(XV) $-\text{[C(R}_{\text{B-1}})(\text{R}_{\text{B-2}})]_{1-3}-\text{CO-N(R}_{\text{B-3}})_2$ where $\text{R}_{\text{B-1}}$ and $\text{R}_{\text{B-2}}$ are the
10 same or different and are selected from the group consisting of:

(A) -H,

(B) $\text{C}_1\text{-C}_6$ alkyl, optionally substituted with one, two or three substituents selected from the group consisting of $\text{C}_1\text{-C}_3$ alkyl, -F, -Cl, -Br, -I, -OH, -SH, $\text{-C}\equiv\text{N}$, -CF_3 , $\text{C}_1\text{-C}_6$ alkoxy, -O-phenyl, and $\text{-NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are as
15 defined above,

(C) $\text{C}_2\text{-C}_6$ alkenyl with one or two double bonds, optionally substituted with one, two or three substituents selected from the group consisting of $\text{C}_1\text{-C}_3$ alkyl, -F, -Cl, -Br, -I, -OH, -SH, $\text{-C}\equiv\text{N}$, -CF_3 , $\text{C}_1\text{-C}_6$ alkoxy, -O-phenyl, and $\text{-NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

20 (D) $\text{C}_2\text{-C}_6$ alkynyl with one or two triple bonds, optionally substituted with one, two or three substituents selected from the group consisting of $\text{C}_1\text{-C}_3$ alkyl, -F, -Cl, -Br, -I, -OH, -SH, $\text{-C}\equiv\text{N}$, -CF_3 , $\text{C}_1\text{-C}_6$ alkoxy, -O-phenyl, and $\text{-NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

(E) $-(\text{CH}_2)_{1-2}-\text{S(O)}_{0-2}-(\text{C}_1\text{-C}_6 \text{ alkyl})$,

25 (F) $-(\text{CH}_2)_{0-4}-\text{C}_3\text{-C}_7$ cycloalkyl, optionally substituted with one, two or three substituents selected from the group consisting of $\text{C}_1\text{-C}_3$ alkyl, -F, -Cl, -Br, -I, -OH, -SH, $\text{-C}\equiv\text{N}$, -CF_3 , $\text{C}_1\text{-C}_6$ alkoxy, -O-phenyl, and $\text{-NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

(G) $-(\text{C}_1\text{-C}_4 \text{ alkyl})-\text{R}_{\text{B'-aryl}}$ where $\text{R}_{\text{B'-aryl}}$ is as defined above for
30 $\text{R}_{1\text{-aryl}}$,

(H) $-(\text{C}_1\text{-C}_4 \text{ alkyl})-\text{R}_{\text{B-heteroaryl}}$ where $\text{R}_{\text{B-heteroaryl}}$ is as defined above,

(I) $-(C_1-C_4 \text{ alkyl})-R_{B\text{-heterocycle}}$ where $R_{B\text{-heterocycle}}$ is as defined above,

(J) $-R_{B\text{-heteroaryl}}$ where $R_{B\text{-heteroaryl}}$ is as defined above,

(K) $-R_{B\text{-heterocycle}}$ where $R_{B\text{-heterocycle}}$ is as defined above,

5 (M) $-(CH_2)_{1-4}-R_{B-4}-(CH_2)_{0-4}-R_{B'\text{-aryl}}$ where R_{B-4} is $-O-$, $-S-$ or $-NR_{B-5}-$ where R_{B-5} is C_1-C_6 alkyl, and where $R_{B'\text{-aryl}}$ is defined above,

(N) $-(CH_2)_{1-4}-R_{B-4}-(CH_2)_{0-4}-R_{B\text{-heteroaryl}}$ where R_{B-4} and $R_{B\text{-heteroaryl}}$ are as defined above, and

(O) $-R_{B'\text{-aryl}}$ where $R_{B'\text{-aryl}}$ is as defined above,

10 and where R_{B-3} is the same or different and is:

(A) $-H$,

(B) C_1-C_6 alkyl optionally substituted with one, two or three substituents selected from the group consisting of C_1-C_3 alkyl, $-F$, $-Cl$, $-Br$, $-I$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, C_1-C_6 alkoxy, $-O$ -phenyl, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

(C) C_2-C_6 alkenyl with one or two double bonds, optionally substituted with one, two or three substituents selected from the group consisting of C_1-C_3 alkyl, $-F$, $-Cl$, $-Br$, $-I$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, C_1-C_6 alkoxy, $-O$ -phenyl, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

20 (D) C_2-C_6 alkynyl with one or two triple bonds, optionally substituted with one, two or three substituents selected from the group consisting of C_1-C_3 alkyl, $-F$, $-Cl$, $-Br$, $-I$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, C_1-C_6 alkoxy, $-O$ -phenyl, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

(E) $-(CH_2)_{0-4}-C_3-C_7$ cycloalkyl, optionally substituted with one, two or three substituents selected from the group consisting of C_1-C_3 alkyl, $-F$, $-Cl$, $-Br$, $-I$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, C_1-C_6 alkoxy, $-O$ -phenyl, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

(F) $-R_{B'\text{-aryl}}$ where $R_{B'\text{-aryl}}$ is as defined above,

(G) $-R_{B\text{-heteroaryl}}$ where $R_{B\text{-heteroaryl}}$ is as defined above,

30 (H) $-R_{B\text{-heterocycle}}$ where $R_{B\text{-heterocycle}}$ is as defined above,

(I) $-(C_1-C_4 \text{ alkyl})-R_{B'\text{-aryl}}$ where $R_{B'\text{-aryl}}$ is as defined above,

(J) $-(C_1-C_4 \text{ alkyl})-R_{B\text{-heteroaryl}}$ where $R_{B\text{-heteroaryl}}$ is as defined above,

(K) $-(C_1-C_4 \text{ alkyl})-R_{B\text{-heterocycle}}$ where $R_{B\text{-heterocycle}}$ is as defined above, or

5 (XVI) $-\text{CH}(R_{B\text{-aryl}})_2$ where $R_{B\text{-aryl}}$ are the same or different and are as defined above,

(XVII) $-\text{CH}(R_{B\text{-heteroaryl}})_2$ where $R_{B\text{-heteroaryl}}$ are the same or different and are as defined above,

10 (XVIII) $-\text{CH}(R_{B\text{-aryl}})(R_{B\text{-heteroaryl}})$ where $R_{B\text{-aryl}}$ and $R_{B\text{-heteroaryl}}$ are as defined above,

(XIX) -cyclopentyl, -cyclohexyl, or -cycloheptyl ring fused to $R_{B\text{-aryl}}$ or $R_{B\text{-heteroaryl}}$ or $R_{B\text{-heterocycle}}$ where $R_{B\text{-aryl}}$ or $R_{B\text{-heteroaryl}}$ or $R_{B\text{-heterocycle}}$ are as defined above where one carbon of cyclopentyl, cyclohexyl, or -cycloheptyl is optionally replaced with NH, NR_{N-5} , O, or $\text{S}(=\text{O})_{0-2}$, and where cyclopentyl, cyclohexyl, or -cycloheptyl can be optionally substituted with one or two $-C_1-C_3$ alkyl, -F, -OH, -SH, $-\text{C}\equiv\text{N}$, $-\text{CF}_3$, C_1-C_6 alkoxy, =O, or $-\text{NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

20 (XX) C_2-C_{10} alkenyl containing one or two double bonds optionally substituted with one, two or three substituents selected from the group consisting of C_1-C_3 alkyl, -F, -Cl, -Br, -I, -OH, -SH, $-\text{C}\equiv\text{N}$, $-\text{CF}_3$, C_1-C_6 alkoxy, -O-phenyl, and $-\text{NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

(XXI) C_2-C_{10} alkynyl containing one or two triple bonds optionally substituted with one, two or three substituents selected from the group consisting of C_1-C_3 alkyl, -F, -Cl, -Br, -I, -OH, -SH, $-\text{C}\equiv\text{N}$, $-\text{CF}_3$, C_1-C_6 alkoxy, -O-phenyl, and $-\text{NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

(XXI) $-(\text{CH}_2)_{0-1}-\text{CHR}_{C-6}-(\text{CH}_2)_{0-1}-R_{B\text{-aryl}}$ where $R_{B\text{-aryl}}$ is as defined above and R_{C-6} is $-(\text{CH}_2)_{0-6}-\text{OH}$,

(XXII) $-(\text{CH}_2)_{0-1}-\text{CHR}_{B-6}-(\text{CH}_2)_{0-1}-R_{B\text{-heteroaryl}}$ where $R_{B\text{-heteroaryl}}$ and R_{C-6} is as defined above,

30 (XXIII) $-\text{CH}(-R_{B\text{-aryl}} \text{ or } R_{B\text{-heteroaryl}})-\text{CO}-\text{O}(C_1-C_4 \text{ alkyl})$ where $R_{B\text{-aryl}}$ and $R_{B\text{-heteroaryl}}$ are as defined above,

(XXIV) $-\text{CH}(-\text{CH}_2-\text{OH})-\text{CH}(-\text{OH})-\text{micro-NO}_2$,

(XXV) $(C_1-C_6 \text{ alkyl})-\text{O}-(C_1-C_6 \text{ alkyl})-\text{OH}$,

(XXVII) $-\text{CH}_2-\text{NH}-\text{CH}_2-\text{CH}(-\text{O}-\text{CH}_2-\text{CH}_3)_2$,

(XXVIII) $-\text{H}$, or

(XXIX) $-(\text{CH}_2)_{0-6}-\text{C}(=\text{NR}_{1-a})(\text{NR}_{1-a}\text{R}_{1-b})$ where R_{1-a} and R_{1-b} are as defined above; and

5

where PROTECTING GROUP is selected from the group consisting of *t*-butoxycarbonyl, benzyloxycarbonyl, formyl, trityl, acetyl, trichloroacetyl, dichloroacetyl, chloroacetyl, trifluoroacetyl, difluoroacetyl, fluoroacetyl, 4-phenylbenzyloxycarbonyl, 2-methylbenzyloxycarbonyl, 4-ethoxybenzyloxycarbonyl, 4-fluorobenzyloxycarbonyl, 4-chlorobenzyloxycarbonyl, 10 3-chlorobenzyloxycarbonyl, 2-chlorobenzyloxycarbonyl, 2,4-dichlorobenzyloxycarbonyl, 4-bromobenzyloxycarbonyl, 3-bromobenzyloxycarbonyl, 4-nitrobenzyloxycarbonyl, 4-cyanobenzyloxycarbonyl, 2-(4-xenyl)isopropoxycarbonyl, 1,1-diphenyleth-1-yloxycarbonyl, 1,1-diphenylprop-1-yloxycarbonyl, 2-phenylprop-2-yloxycarbonyl, 2-(*p*-toluyl)prop-2-yloxycarbonyl, 15 cyclopentanyloxycarbonyl, 1-methylcyclopentanyloxycarbonyl, cyclohexanyloxycarbonyl, 1-methylcyclohexanyloxycarbonyl, 2-methylcyclohexanyloxycarbonyl, 2-(4-toluylsulfonyl)ethoxycarbonyl, 2-(methylsulfonyl)ethoxycarbonyl, 2-(triphenylphosphino)ethoxycarbonyl, 20 fluorenylmethoxycarbonyl, 2-(trimethylsilyl)ethoxycarbonyl, allyloxycarbonyl, 1-(trimethylsilylmethyl)prop-1-enyloxycarbonyl, 5-benzisoxalylmethoxycarbonyl, 4-acetoxybenzyloxycarbonyl, 2,2,2-trichloroethoxycarbonyl, 2-ethynyl-2-propoxycarbonyl, cyclopropylmethoxycarbonyl, 4-(decyloxy)benzyloxycarbonyl, isobornyloxycarbonyl and 1-piperidyloxycarbonyl, 9-fluorenylmethyl carbonate, -25 $\text{CH}-\text{CH}=\text{CH}_2$ and phenyl- $\text{C}(=\text{N})-\text{H}$.

37. A protected compound according to claim 36

where R_1 is:

30 $-(\text{CH}_2)_{0-1}-(\text{R}_{1-\text{aryl}})$, or
 $-(\text{CH}_2)_{n1}-(\text{R}_{1-\text{heteroaryl}})$;

where R_N is:

$\text{R}_{N-1}-\text{X}_N$, where X_N is selected from the group consisting of:

$-\text{CO}-$, and

$-\text{SO}_2-$,

where R_{N-1} is selected from the group consisting of:

$-R_{N-aryl}$, and

$-R_{N-heteroaryl}$, or

$-\text{CO}-\text{CH}(-(\text{CH}_2)_{0-2}-\text{O}-R_{N-10})-(\text{CH}_2)_{0-2}-R_{N-aryl}/R_{N-heteroaryl}$;

5 where R_A is:

$-\text{C}_1-\text{C}_8$ alkyl,

$-(\text{CH}_2)_{0-3}-(\text{C}_3-\text{C}_7)$ cycloalkyl,

$-(\text{CR}_{A-x}\text{R}_{A-y})_{0-4}-R_{A-aryl}$,

$-(\text{CR}_{A-x}\text{R}_{A-y})_{0-4}-R_{A-heteroaryl}$,

10 $-(\text{CR}_{A-x}\text{R}_{A-y})_{0-4}-R_{A-heterocycle}$,

$-\text{cyclopentyl}$ or $-\text{cyclohexyl}$ ring fused to R_{A-aryl} or $R_{A-heteroaryl}$ or R_{A-}

heterocycle; and

where R_B is:

$-\text{C}_1-\text{C}_8$ alkyl,

15 $-(\text{CH}_2)_{0-3}-(\text{C}_3-\text{C}_7)$ cycloalkyl,

$-(\text{CR}_{A-x}\text{R}_{A-y})_{0-4}-R_{A-aryl}$,

$-(\text{CR}_{A-x}\text{R}_{A-y})_{0-4}-R_{A-heteroaryl}$,

$-(\text{CR}_{A-x}\text{R}_{A-y})_{0-4}-R_{A-heterocycle}$,

$-\text{cyclopentyl}$ or $-\text{cyclohexyl}$ ring fused to R_{A-aryl} or $R_{A-heteroaryl}$ or $R_{A-heterocycle}$.

20

38. A protected compound according to claim 37

where R_1 is:

$-(\text{CH}_2)-(\text{R}_{1-aryl})$, or

$-(\text{CH}_2)-(\text{R}_{1-heteroaryl})$;

25 where R_2 is $-\text{H}$;

where R_3 is $-\text{H}$;

where R_N is:

$R_{N-1}-X_N$ where X_N is:

$-\text{CO}-$,

30 where R_{N-1} is selected from the group consisting of:

$-R_{N-aryl}$, and

$-R_{N-heteroaryl}$;

where R_A is:

$-\text{C}_1-\text{C}_8$ alkyl,

- 5 $-(CH_2)_{0-3}-(C_3-C_7)$ cycloalkyl,
 $-(CR_{A-x}R_{A-y})_{0-4}-R_{A-aryl}$,
 $-(CR_{A-x}R_{A-y})_{0-4}-R_{A-heteroaryl}$,
 $-(CR_{A-x}R_{A-y})_{0-4}-R_{A-heterocycle}$,
 -cyclopentyl or -cyclohexyl ring fused to R_{A-aryl} or $R_{A-heteroaryl}$ or $R_{A-heterocycle}$;

where R_B is:


- 10 $-C_1-C_8$ alkyl,
 $-(CH_2)_{0-3}-(C_3-C_7)$ cycloalkyl,
 $-(CR_{B-x}R_{B-y})_{0-4}-R_{B-aryl}$,
 $-(CR_{B-x}R_{B-y})_{0-4}-R_{B-heteroaryl}$,
 $-(CR_{B-x}R_{B-y})_{0-4}-R_{B-heterocycle}$,
 -cyclopentyl or -cyclohexyl ring fused to R_{B-aryl} or $R_{B-heteroaryl}$ or $R_{B-heterocycle}$.

15

39. A protected compound according to claim 36 where PROTECTING GROUP is *t*-butoxycarbonyl.

20 40. A protected compound according to claim 36 where PROTECTING GROUP is benzyloxycarbonyl.

41. A protected compound of the formula (IV)



25

where R_1 is:

(I) C_1-C_6 alkyl, optionally substituted with one, two or three
 substituents selected from the group consisting of C_1-C_3 alkyl, C_1-C_7 alkyl
 (optionally substituted with C_1-C_3 alkyl and C_1-C_3 alkoxy), -F, -Cl, -Br, -I, -OH, -

SH, $-C\equiv N$, $-CF_3$, C_1-C_3 alkoxy, $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are $-H$ or C_1-C_6 alkyl, and $-OC=O NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

(II) $-CH_2-S(O)_{0-2}-(C_1-C_6 \text{ alkyl})$,

(III) $-CH_2-CH_2-S(O)_{0-2}-(C_1-C_6 \text{ alkyl})$,

5 (IV) C_2-C_6 alkenyl with one or two double bonds, optionally substituted with one, two or three substituents selected from the group consisting of $-F$, $-Cl$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, C_1-C_3 alkoxy, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are $-H$ or C_1-C_6 alkyl,

(V) C_2-C_6 alkynyl with one or two triple bonds, optionally substituted
10 with one, two or three substituents selected from the group consisting of $-F$, $-Cl$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, C_1-C_3 alkoxy, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are $-H$ or C_1-C_6 alkyl,

(VI) $-(CH_2)_{n1}-(R_{1-aryl})$ where n_1 is zero or one and where R_{1-aryl} is phenyl, 1-naphthyl, 2-naphthyl and indanyl, indenyl, dihydronaphthalyl, or tetralinyl
15 optionally substituted with one, two, three or four of the following substituents on the aryl ring:

(A) C_1-C_6 alkyl optionally substituted with one, two or three substituents selected from the group consisting of C_1-C_3 alkyl, $-F$, $-Cl$, $-Br$, $-I$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, and C_1-C_3 alkoxy, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as
20 defined above,

(B) C_2-C_6 alkenyl with one or two double bonds, optionally substituted with one, two or three substituents selected from the group consisting of $-F$, $-Cl$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, C_1-C_3 alkoxy, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are $-H$ or C_1-C_6 alkyl,

25 (C) C_2-C_6 alkynyl with one or two triple bonds, optionally substituted with one, two or three substituents selected from the group consisting of $-F$, $-Cl$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, C_1-C_3 alkoxy, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are $-H$ or C_1-C_6 alkyl,

(D) $-F$, Cl , $-Br$ or $-I$,

30 (F) C_1-C_6 alkoxy optionally substituted with one, two or three of $-F$,

(G) $-NR_{N-2}R_{N-3}$ where R_{N-2} and R_{N-3} are as defined below,

(H) $-OH$,

(I) $-C\equiv N$,

(J) C_3-C_7 cycloalkyl, optionally substituted with one, two or three substituents selected from the group consisting of $-F$, $-Cl$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, C_1-C_3 alkoxy, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are $-H$ or C_1-C_6 alkyl,

5 (K) $-\text{CO}-(C_1-C_4 \text{ alkyl})$,

(L) $-\text{SO}_2-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

(M) $-\text{CO}-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

or

(N) $-\text{SO}_2-(C_1-C_4 \text{ alkyl})$,

10 (VII) $-(\text{CH}_2)_{n_1}-(R_{1-\text{heteroaryl}})$ where n_1 is as defined above and where $R_{1-\text{heteroaryl}}$ is selected from the group consisting of:

pyridinyl,

pyrimidinyl,

quinolinyl,

15 benzothienyl,

indolyl,

indolinyl,

pyridazinyl,

pyrazinyl,

20 isoquinolyl,

quinazolinyl,

quinoxalinyl,

phthalazinyl,

imidazolyl,

25 isoxazolyl,

pyrazolyl,

oxazolyl,

thiazolyl,

indolizinyl,

30 indazolyl,

benzothiazolyl,

benzimidazolyl,

benzofuranyl,

	furanyl,
	thienyl,
	pyrrolyl,
	oxadiazolyl,
5	thiadiazolyl,
	triazolyl,
	tetrazolyl,
	oxazolopyridinyl,
	imidazopyridinyl,
10	isothiazolyl,
	naphthyridinyl,
	cinnolinyl,
	carbazolyl,
	beta-carbolinyl,
15	isochromanyl,
	chromanyl,
	tetrahydroisoquinolinyl,
	isoindolinyl,
	isobenzotetrahydrofuranyl,
20	isobenzotetrahydrothienyl,
	isobenzothienyl,
	benzoxazolyl,
	pyridopyridinyl,
	benzotetrahydrofuranyl,
25	benzotetrahydrothienyl,
	purinyl,
	benzodioxolyl,
	triazinyl,
	phenoxazinyl,
30	phenothiazinyl,
	pteridinyl,
	benzothiazolyl,
	imidazopyridinyl,
	imidazothiazolyl,

	dihydrobenzisoxaziny1,
	benzisoxaziny1,
	benzoxaziny1,
	dihydrobenzisothiaziny1,
5	benzopyrany1,
	benzothiopyrany1,
	coumariny1,
	isocoumariny1,
	chromony1,
10	chromanony1,
	pyridiny1-N-oxide,
	tetrahydroquinoliny1
	dihydroquinoliny1
	dihydroquinolinony1
15	dihydroisoquinolinony1
	dihydrocoumariny1
	dihydroisocoumariny1
	isoindolinony1
	benzodioxany1
20	benzoxazolinony1
	pyrroly1 N-oxide,
	pyrimidiny1 N-oxide,
	pyridaziny1 N-oxide,
	pyraziny1 N-oxide,
25	quinoliny1 N-oxide,
	indoly1 N-oxide,
	indoliny1 N-oxide,
	isoquinoly1 N-oxide,
	quinazolinyl N-oxide,
30	quinoxaliny1 N-oxide,
	phthalaziny1 N-oxide,
	imidazolyl N-oxide,
	isoxazolyl N-oxide,
	oxazolyl N-oxide,

thiazolyl N-oxide,
indoliziny N-oxide,
indazolyl N-oxide,
benzothiazolyl N-oxide,
5 benzimidazolyl N-oxide,
pyrrolyl N-oxide,
oxadiazolyl N-oxide,
thiadiazolyl N-oxide,
triazolyl N-oxide,
10 tetrazolyl N-oxide,
benzothiopyranyl S-oxide, and
benzothiopyranyl S,S-dioxide,

where the $R_{1\text{-heteroaryl}}$ group is bonded to $-(CH_2)_{n1}-$ by any ring atom of the parent $R_{1\text{-heteroaryl}}$ group substituted by hydrogen such that the new bond to the
15 $R_{1\text{-heteroaryl}}$ group replaces the hydrogen atom and its bond, where heteroaryl is optionally substituted with one, two, three or four of:

(1) C_1-C_6 alkyl optionally substituted with one, two or three substituents selected from the group consisting of C_1-C_3 alkyl, -F, -Cl, -Br, -I, -OH, -SH, $-C\equiv N$, $-CF_3$, C_1-C_3 alkoxy, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as
20 defined above,

(2) C_2-C_6 alkenyl with one or two double bonds, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, $-C\equiv N$, $-CF_3$, C_1-C_3 alkoxy, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are -H or C_1-C_6 alkyl,
25

(3) C_2-C_6 alkynyl with one or two triple bonds, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, $-C\equiv N$, $-CF_3$, C_1-C_3 alkoxy, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are -H or C_1-C_6 alkyl,
25

(4) -F, Cl, -Br or -I,
30 (6) $-C_1-C_6$ alkoxy optionally substituted with one, two, or three of -F,

(7) $-NR_{N-2}R_{N-3}$ where R_{N-2} and R_{N-3} are as defined below,

(8) -OH,

(9) -C≡N,

(10) C₃-C₇ cycloalkyl, optionally substituted with one,
two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, -
5 C≡N, -CF₃, C₁-C₃ alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are -H or C₁-C₆ alkyl,

(11) -CO-(C₁-C₄ alkyl),

(12) -SO₂-NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined
above,

(13) -CO-NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined
10 above, or

(14) -SO₂-(C₁-C₄ alkyl), with the proviso that when n₁
is zero R_{1-heteroaryl} is not bonded to the carbon chain by nitrogen; or

(VIII) -(CH₂)_{n1}-(R_{1-heterocycle}) where n₁ is as defined above and R<sub>1-
heterocycle</sub> is selected from the group consisting of:

15 morpholinyl,
thiomorpholinyl,
thiomorpholinyl S-oxide,
thiomorpholinyl S,S-dioxide,
piperazinyl,
20 homopiperazinyl,
pyrrolidinyl,
pyrrolinyl,
tetrahydropyranyl,
piperidinyl,
25 tetrahydrofuranyl,
tetrahydrothienyl,
homopiperidinyl,
homomorpholinyl,
homothiomorpholinyl,
30 homothiomorpholinyl S,S-dioxide,
oxazolidinonyl,
dihydropyrazolyl,
dihydropyrrolyl,

245

dihydropyrazinyl,
 dihydropyridinyl,
 dihydropyrimidinyl,
 dihydrofuryl,
 5 dihydropyranyl,
 tetrahydrothienyl S-oxide,
 tetrahydrothienyl S,S-dioxide, and
 homothiomorpholinyl S-oxide,

where the $R_{1\text{-heterocycle}}$ group is bonded by any atom of the parent $R_{1\text{-heterocycle}}$ group substituted by hydrogen such that the new bond to the $R_{1\text{-heterocycle}}$ group replaces the hydrogen atom and its bond, where heterocycle is optionally substituted with one, two, three or four:

(1) $C_1\text{-}C_6$ alkyl optionally substituted with one, two or three substituents selected from the group consisting of $C_1\text{-}C_3$ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C \equiv N, -CF₃, $C_1\text{-}C_3$ alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(2) $C_2\text{-}C_6$ alkenyl with one or two double bonds, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, -C \equiv N, -CF₃, $C_1\text{-}C_3$ alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are -H or $C_1\text{-}C_6$ alkyl,

(3) $C_2\text{-}C_6$ alkynyl with one or two triple bonds, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, -C \equiv N, -CF₃, $C_1\text{-}C_3$ alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are -H or $C_1\text{-}C_6$ alkyl,

(4) -F, Cl, -Br or -I,
 (5) $C_1\text{-}C_6$ alkoxy,
 (6) $C_1\text{-}C_6$ alkoxy optionally substituted with one, two, or three -F,

(7) -NR_{N-2}R_{N-3} where R_{N-2} and R_{N-3} are as defined below,

(8) -OH,
 (9) -C \equiv N,

- (10) C₃-C₇ cycloalkyl, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, -C≡N, -CF₃, C₁-C₃ alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are -H or C₁-C₆ alkyl,
- (11) -CO-(C₁-C₄ alkyl),
- 5 (12) -SO₂-NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,
- (13) -CO-NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,
- (14) -SO₂-(C₁-C₄ alkyl), or
- 10 (15) =O, with the proviso that when n₁ is zero R_{1-heterocycle} is not bonded to the carbon chain by nitrogen;

where R₂ is:

- (I)-H,
- 15 (II) C₁-C₆ alkyl, optionally substituted with one, two or three substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₃ alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,
- (III) -(CH₂)₀₋₄-R₂₋₁ where R₂₋₁ is R_{1-aryl} or R_{1-heteroaryl} where R_{1-aryl} and R_{1-heteroaryl} are as defined above;
- 20 (IV) C₂-C₆ alkenyl with one or two double bonds, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, -C≡N, -CF₃, C₁-C₃ alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are -H or C₁-C₆ alkyl,
- 25 (V) C₂-C₆ alkynyl with one or two triple bonds, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, -C≡N, -CF₃, C₁-C₃ alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are -H or C₁-C₆ alkyl, or
- (VI) -(CH₂)₀₋₄-C₃-C₇ cycloalkyl, optionally substituted with one, two
- 30 or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, -C≡N, -CF₃, C₁-C₃ alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are -H or C₁-C₆ alkyl;

where R₃ is:

(I)-H,

(II) C₁-C₆ alkyl, optionally substituted with one, two or three substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₃ alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined

5 above,

(III) -(CH₂)₀₋₄-R₂₋₁ where R₂₋₁ is R_{1-aryl} or R_{1-heteroaryl} where R_{1-aryl} and R_{1-heteroaryl} are as defined above

(IV) C₂-C₆ alkenyl with one or two double bonds,

(V) C₂-C₆ alkynyl with one or two triple bonds; or

10 (VI) -(CH₂)₀₋₄- C₃-C₇ cycloalkyl, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, -C≡N, -CF₃, C₁-C₃ alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are -H or C₁-C₆ alkyl,

and where R₂ and R₃ are taken together with the carbon to which they are attached to form a carbocycle of three, four, five, six, or seven carbon atoms,

15 optionally where one carbon atom is replaced by a heteroatom selected from the group consisting of -O-, -S-, -SO₂-, and -NR_{N-2}-, where R_{N-2} is selected from the group consisting of:

(a) -H,

(b) -C₁-C₆ alkyl optionally substituted with one

20 substituent selected from the group consisting of:

(i) -OH, and

(ii) -NH₂,

(c) -C₁-C₆ alkyl optionally substituted with

one, two, or three -F, -Cl, -Br, or -I,

25 (d) -C₃-C₇ cycloalkyl,

(e) -(C₁-C₂ alkyl)-(C₃-C₇ cycloalkyl),

(f) -(C₁-C₆ alkyl)-O-(C₁-C₃ alkyl),

(g) -C₂-C₆ alkenyl with one or two double

bonds,

30 (h) -C₂-C₆ alkynyl with one or two triple

bonds,

(i) -C₁-C₆ alkyl chain with one double bond

and one triple bond,

(j) $-R_{1-aryl}$ where R_{1-aryl} is as defined above, and

(k) $-R_{1-heteroaryl}$ where $R_{1-heteroaryl}$ is as defined

above;

5 where R_N is:

(I) $R_{N-1}-X_N-$ where X_N is selected from the group consisting of:

(A) $-\text{CO}-$,

(B) $-\text{SO}_2-$,

(C) $-(\text{CR}'\text{R}'')_{1-6}$ where R' and R'' are the same or different and

10 are $-\text{H}$ and $\text{C}_1\text{-C}_4$ alkyl,

(D) $-\text{CO}-(\text{CR}'\text{R}'')_{1-6}-X_{N-1}$ where X_{N-1} is selected from the group consisting of $-\text{O}-$, $-\text{S}-$ and $-\text{NR}'-$ and where R' and R'' are as defined above, and

(E) a single bond;

15 where R_{N-1} is selected from the group consisting of:

(A) R_{N-aryl} where R_{N-aryl} is phenyl, 1-naphthyl, 2-naphthyl, tetralinyl, indanyl, 6,7,8,9-tetrahydro-5H-benzo[a]cycloheptenyl, or dihydronaphthyl optionally substituted with one, two or three of the following substituents which can be the same or different and are:

20 (1) $\text{C}_1\text{-C}_6$ alkyl, optionally substituted with one, two or three substituents selected from the group consisting of $\text{C}_1\text{-C}_3$ alkyl, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, $-\text{OH}$, $-\text{SH}$, $-\text{C}\equiv\text{N}$, $-\text{CF}_3$, $\text{C}_1\text{-C}_3$ alkoxy, and $-\text{NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

(2) $-\text{OH}$,

25 (3) $-\text{NO}_2$,

(4) $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$,

(5) $-\text{CO}-\text{OH}$,

(6) $-\text{C}\equiv\text{N}$,

(7) $-(\text{CH}_2)_{0-4}-\text{CO}-\text{NR}_{N-2}\text{R}_{N-3}$ where R_{N-2} and R_{N-3} are

30 the same or different and are selected from the group consisting of:

(a) $-\text{H}$,

(b) $\text{C}_1\text{-C}_6$ alkyl optionally substituted with one substituent selected from the group consisting of:

- (i) -OH, and
(ii) -NH₂,
(c) -C₁-C₆ alkyl optionally substituted with one, two, or three -F, -Cl, -Br, or -I,
5 (d) -C₃-C₇ cycloalkyl,
(e) -(C₁-C₂ alkyl)-(C₃-C₇ cycloalkyl),
(f) -(C₁-C₆ alkyl)-O-(C₁-C₃ alkyl),
(g) -C₂-C₆ alkenyl with one or two double bonds,
10 (h) -C₂-C₆ alkynyl with one or two triple bonds,
(i) -C₁-C₆ alkyl chain with one double bond and one triple bond,
(j) -R_{1-aryl} where R_{1-aryl} is as defined above, and
15 (k) -R_{1-heteroaryl} where R_{1-heteroaryl} is as defined above,
(8) -(CH₂)₀₋₄-CO-(C₁-C₁₂ alkyl),
(9) -(CH₂)₀₋₄-CO-(C₂-C₁₂ alkenyl with one, two or three double bonds),
20 (10) -(CH₂)₀₋₄-CO-(C₂-C₁₂ alkynyl with one, two or three triple bonds),
(11) -(CH₂)₀₋₄-CO-(C₃-C₇ cycloalkyl),
(12) -(CH₂)₀₋₄-CO-R_{1-aryl} where R_{1-aryl} is as defined above,
25 (13) -(CH₂)₀₋₄-CO-R_{1-heteroaryl} where R_{1-heteroaryl} is as defined above,
(14) -(CH₂)₀₋₄-CO-R_{1-heterocycle} where R_{1-heterocycle} is as defined above,
(15) -(CH₂)₀₋₄-CO-R_{N-4} where R_{N-4} is selected from
30 the group consisting of morpholinyl, thiomorpholinyl, piperazinyl, piperidinyl, homomorpholinyl, homothiomorpholinyl, homothiomorpholinyl S-oxide, homothiomorpholinyl S,S-dioxide, pyrrolinyl and pyrrolidinyl where each group is optionally substituted with one, two, three, or four of C₁-C₆ alkyl,

(16) $-(\text{CH}_2)_{0-4}-\text{CO}-\text{O}-\text{R}_{\text{N}-5}$ where $\text{R}_{\text{N}-5}$ is

selected from the group consisting of:

(a) C_1-C_6 alkyl,

(b) $-(\text{CH}_2)_{0-2}-(\text{R}_{1-\text{aryl}})$ where $\text{R}_{1-\text{aryl}}$ is as defined

5 above,

(c) C_2-C_6 alkenyl containing one or two double

bonds,

(d) C_2-C_6 alkynyl containing one or two triple

bonds,

10

(e) C_3-C_7 cycloalkyl, and

(f) $-(\text{CH}_2)_{0-2}-(\text{R}_{1-\text{heteroaryl}})$ where $\text{R}_{1-\text{heteroaryl}}$ is as

defined above,

(17) $-(\text{CH}_2)_{0-4}-\text{SO}_2-\text{NR}_{\text{N}-2}\text{R}_{\text{N}-3}$ where $\text{R}_{\text{N}-2}$ and $\text{R}_{\text{N}-3}$ are

as defined above,

15

(18) $-(\text{CH}_2)_{0-4}-\text{SO}-(\text{C}_1-\text{C}_8 \text{ alkyl})$,

(19) $-(\text{CH}_2)_{0-4}-\text{SO}_2-(\text{C}_1-\text{C}_{12} \text{ alkyl})$,

(20) $-(\text{CH}_2)_{0-4}-\text{SO}_2-(\text{C}_3-\text{C}_7 \text{ cycloalkyl})$,

(21) $-(\text{CH}_2)_{0-4}-\text{N}(\text{H or } \text{R}_{\text{N}-5})-\text{CO}-\text{O}-\text{R}_{\text{N}-5}$ where $\text{R}_{\text{N}-5}$

can be the same or different and is as defined above,

20

(22) $-(\text{CH}_2)_{0-4}-\text{N}(\text{H or } \text{R}_{\text{N}-5})-\text{CO}-\text{N}(\text{R}_{\text{N}-5})_2$, where $\text{R}_{\text{N}-5}$

can be the same or different and is as defined above,

(23) $-(\text{CH}_2)_{0-4}-\text{N}-\text{CS}-\text{N}(\text{R}_{\text{N}-5})_2$, where $\text{R}_{\text{N}-5}$ can be the

same or different and is as defined above,

(24) $-(\text{CH}_2)_{0-4}-\text{N}(-\text{H or } \text{R}_{\text{N}-5})-\text{CO}-\text{R}_{\text{N}-2}$ where $\text{R}_{\text{N}-5}$ and

25

$\text{R}_{\text{N}-2}$ can be the same or different and are as defined above,

(25) $-(\text{CH}_2)_{0-4}-\text{NR}_{\text{N}-2}\text{R}_{\text{N}-3}$ where $\text{R}_{\text{N}-2}$ and $\text{R}_{\text{N}-3}$ can be

the same or different and are as defined above,

(26) $-(\text{CH}_2)_{0-4}-\text{R}_{\text{N}-4}$ where $\text{R}_{\text{N}-4}$ is as defined above,

(27) $-(\text{CH}_2)_{0-4}-\text{O}-\text{CO}-(\text{C}_1-\text{C}_6 \text{ alkyl})$,

30

(28) $-(\text{CH}_2)_{0-4}-\text{O}-\text{P}(\text{O})-(\text{OR}_{\text{N-aryl-1}})_2$ where $\text{R}_{\text{N-aryl-1}}$ is -

H or C_1-C_4 alkyl,

(29) $-(\text{CH}_2)_{0-4}-\text{O}-\text{CO}-\text{N}(\text{R}_{\text{N}-5})_2$ where $\text{R}_{\text{N}-5}$ is as

defined above,

- above,
- (30) $-(\text{CH}_2)_{0-4}-\text{O}-\text{CS}-\text{N}(\text{R}_{\text{N}-5})_2$ where $\text{R}_{\text{N}-5}$ is as defined
- above,
- (31) $-(\text{CH}_2)_{0-4}-\text{O}-(\text{R}_{\text{N}-5})_2$ where $\text{R}_{\text{N}-5}$ is as defined
- above,
- 5 (32) $-(\text{CH}_2)_{0-4}-\text{O}-(\text{R}_{\text{N}-5})_2-\text{COOH}$ where $\text{R}_{\text{N}-5}$ is as defined above,
- (33) $-(\text{CH}_2)_{0-4}-\text{S}-(\text{R}_{\text{N}-5})_2$ where $\text{R}_{\text{N}-5}$ is as defined
- above,
- (34) $-(\text{CH}_2)_{0-4}-\text{O}-(\text{C}_1-\text{C}_6 \text{ alkyl optionally substituted$
- 10 with one, two, three, four, or five of $-\text{F}$),
- (35) C_3-C_7 cycloalkyl,
- (36) C_2-C_6 alkenyl with one or two double bonds optionally substituted with C_1-C_3 alkyl, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, $-\text{OH}$, $-\text{SH}$, $-\text{C}\equiv\text{N}$, $-\text{CF}_3$, C_1-C_3 alkoxy, or $-\text{NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,
- 15 (37) C_2-C_6 alkynyl with one or two triple bonds optionally substituted with C_1-C_3 alkyl, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, $-\text{OH}$, $-\text{SH}$, $-\text{C}\equiv\text{N}$, $-\text{CF}_3$, C_1-C_3 alkoxy, or $-\text{NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,
- (38) $-(\text{CH}_2)_{0-4}-\text{N}(-\text{H} \text{ or } \text{R}_{\text{N}-5})-\text{SO}_2-\text{R}_{\text{N}-2}$ where $\text{R}_{\text{N}-5}$ and $\text{R}_{\text{N}-2}$ can be the same or different and are as described above, or
- 20 (39) $-(\text{CH}_2)_{0-4}-\text{C}_3-\text{C}_7$ cycloalkyl,
- (B) $-\text{R}_{\text{N-heteroaryl}}$ where $\text{R}_{\text{N-heteroaryl}}$ is selected from the group consisting of:
- pyridinyl,
- pyrimidinyl,
- 25 quinolinyl,
- benzothienyl,
- indolyl,
- indolinyl,
- pyridazinyl,
- 30 pyrazinyl,
- isoindolyl,
- isoquinolyl,
- quinazolinyl,

	quinoxaliny1,
	phthalaziny1,
	imidazolyl,
	isoxazolyl,
5	pyrazolyl,
	oxazolyl,
	thiazolyl,
	indoliziny1,
	indazolyl,
10	benzothiazolyl,
	benzimidazolyl,
	benzofurany1,
	furany1,
	thienyl,
15	pyrroly1,
	oxadiazolyl,
	thiadiazolyl,
	triazolyl,
	tetrazolyl,
20	oxazolopyridiny1,
	imidazopyridiny1,
	isothiazolyl,
	naphthyridiny1,
	cinnoliny1,
25	carbazolyl,
	beta-carboliny1,
	isochromany1,
	chromany1,
	tetrahydroisoquinoliny1,
30	isoindoliny1,
	isobenzotetrahydrofurany1,
	isobenzotetrahydrothienyl,
	isobenzothiény1,
	benzoxazolyl,

	pyridopyridinyl,
	benzotetrahydrofuranyl,
	benzotetrahydrothienyl,
	purinyl,
5	benzodioxolyl,
	triazinyl,
	phenoxazinyl,
	phenothiazinyl,
	pteridinyl,
10	benzothiazolyl,
	imidazopyridinyl,
	imidazothiazolyl,
	dihydrobenzisoxazinyl,
	benzisoxazinyl,
15	benzoxazinyl,
	dihydrobenzisothoniazinyl,
	benzopyranyl,
	benzothiopyranyl,
	coumarinyl,
20	isocoumarinyl,
	chromonyl,
	chromanonyl,
	pyridinyl-N-oxide,
	tetrahydroquinolinyl,
25	dihydroquinolinyl,
	dihydroquinolinonyl,
	dihydroisoquinolinonyl,
	dihydrocoumarinyl,
	dihydroisocoumarinyl,
30	isoindolinonyl,
	benzodioxanyl,
	benzoxazolinonyl,
	pyrrolyl N-oxide,
	pyrimidinyl N-oxide,

5 pyridazinyl N-oxide,
 pyrazinyl N-oxide,
 quinolinyl N-oxide,
 indolyl N-oxide,
 indoliny N-oxide,
 isoquinolyl N-oxide,
 quinazoliny N-oxide,
 quinoxaliny N-oxide,
 10 phthalazinyl N-oxide,
 imidazolyl N-oxide,
 isoxazolyl N-oxide,
 oxazolyl N-oxide,
 thiazolyl N-oxide,
 indoliziny N-oxide,
 15 indazolyl N-oxide,
 benzothiazolyl N-oxide,
 benzimidazolyl N-oxide,
 pyrrolyl N-oxide,
 oxadiazolyl N-oxide,
 20 thiadiazolyl N-oxide,
 triazolyl N-oxide,
 tetrazolyl N-oxide,
 benzothiopyranyl S-oxide, and
 benzothiopyranyl S,S-dioxide

25 where the $R_{N\text{-heteroaryl}}$ group is bonded by any atom of the
 parent $R_{N\text{-heteroaryl}}$ group substituted by hydrogen such that the new bond to the $R_{N\text{-heteroaryl}}$
 group replaces the hydrogen atom and its bond, where heteroaryl is
 optionally substituted with one, two, three, or four of:

30 (1) $C_1\text{-}C_6$ alkyl, optionally substituted with one, two or
 three substituents selected from the group consisting of $C_1\text{-}C_3$ alkyl, -F, -Cl, -Br, -I,
 -OH, -SH, $-C\equiv N$, $-CF_3$, $C_1\text{-}C_3$ alkoxy, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as
 defined above,

(2) -OH,

- (3) $-\text{NO}_2$,
 (4) $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, or $-\text{I}$,
 (5) $-\text{CO}-\text{OH}$,
 (6) $-\text{C}\equiv\text{N}$,
 5 (7) $-(\text{CH}_2)_{0-4}-\text{CO}-\text{NR}_{\text{N}-2}\text{R}_{\text{N}-3}$ where $\text{R}_{\text{N}-2}$ and $\text{R}_{\text{N}-3}$ are the same or different and are selected from the group consisting of:
 (a) $-\text{H}$,
 (b) $-\text{C}_1-\text{C}_6$ alkyl optionally substituted with one substituent selected from the group consisting of:
 10 (i) $-\text{OH}$, and
 (ii) $-\text{NH}_2$,
 (c) $-\text{C}_1-\text{C}_6$ alkyl optionally substituted with one, two, or three $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$,
 (d) $-\text{C}_3-\text{C}_7$ cycloalkyl,
 15 (e) $-(\text{C}_1-\text{C}_2 \text{ alkyl})-(\text{C}_3-\text{C}_7 \text{ cycloalkyl})$,
 (f) $-(\text{C}_1-\text{C}_6 \text{ alkyl})-\text{O}-(\text{C}_1-\text{C}_3 \text{ alkyl})$,
 (g) $-\text{C}_2-\text{C}_6$ alkenyl with one or two double bonds,
 (h) $-\text{C}_2-\text{C}_6$ alkynyl with one or two triple bonds,
 20 (i) $-\text{C}_1-\text{C}_6$ alkyl chain with one double bond and one triple bond,
 (j) $-\text{R}_{1-\text{aryl}}$ where $\text{R}_{1-\text{aryl}}$ is as defined above,
 (k) $-\text{R}_{1-\text{heteroaryl}}$ where $\text{R}_{1-\text{heteroaryl}}$ is as defined above,
 25 above,
 (8) $-(\text{CH}_2)_{0-4}-\text{CO}-(\text{C}_1-\text{C}_{12} \text{ alkyl})$,
 (9) $-(\text{CH}_2)_{0-4}-\text{CO}-(\text{C}_2-\text{C}_{12} \text{ alkenyl with one, two or three double bonds})$,
 (10) $-(\text{CH}_2)_{0-4}-\text{CO}-(\text{C}_2-\text{C}_{12} \text{ alkynyl with one, two or three triple bonds})$,
 30 (11) $-(\text{CH}_2)_{0-4}-\text{CO}-(\text{C}_3-\text{C}_7 \text{ cycloalkyl})$,
 (12) $-(\text{CH}_2)_{0-4}-\text{CO}-\text{R}_{1-\text{aryl}}$ where $\text{R}_{1-\text{aryl}}$ is as defined above,

(13) $-(\text{CH}_2)_{0-4}-\text{CO}-\text{R}_{1-\text{heteroaryl}}$ where $\text{R}_{1-\text{heteroaryl}}$ is as defined above,

(14) $-(\text{CH}_2)_{0-4}-\text{CO}-\text{R}_{1-\text{heterocycle}}$ where $\text{R}_{1-\text{heterocycle}}$ is as defined above,

5 (15) $-(\text{CH}_2)_{0-4}-\text{CO}-\text{R}_{\text{N-4}}$ where $\text{R}_{\text{N-4}}$ is selected from the group consisting of morpholinyl, thiomorpholinyl, piperazinyl, piperidinyl, homomorpholinyl, homothiomorpholinyl, homothiomorpholinyl S-oxide, homothiomorpholinyl S,S-dioxide, pyrrolinyl and pyrrolidinyl where each group is optionally substituted with one, two, three, or four of C_1-C_6 alkyl,

10 (16) $-(\text{CH}_2)_{0-4}-\text{CO}-\text{O}-\text{R}_{\text{N-5}}$ where $\text{R}_{\text{N-5}}$ is selected from the group consisting of:

(a) C_1-C_6 alkyl,

(b) $-(\text{CH}_2)_{0-2}-(\text{R}_{1-\text{aryl}})$ where $\text{R}_{1-\text{aryl}}$ is as defined above,

15 (c) C_2-C_6 alkenyl containing one or two double bonds,

(d) C_2-C_6 alkynyl containing one or two triple bonds,

(e) C_3-C_7 cycloalkyl, and

20 (f) $-(\text{CH}_2)_{0-2}-(\text{R}_{1-\text{heteroaryl}})$ where $\text{R}_{1-\text{heteroaryl}}$ is as defined above,

(17) $-(\text{CH}_2)_{0-4}-\text{SO}_2-\text{NR}_{\text{N-2}}\text{R}_{\text{N-3}}$ where $\text{R}_{\text{N-2}}$ and $\text{R}_{\text{N-3}}$ are as defined above,

(18) $-(\text{CH}_2)_{0-4}-\text{SO}-(\text{C}_1-\text{C}_8 \text{ alkyl})$,

25 (19) $-(\text{CH}_2)_{0-4}-\text{SO}_2-(\text{C}_1-\text{C}_{12} \text{ alkyl})$,

(20) $-(\text{CH}_2)_{0-4}-\text{SO}_2-(\text{C}_3-\text{C}_7 \text{ cycloalkyl})$,

(21) $-(\text{CH}_2)_{0-4}-\text{N}(\text{H or } \text{R}_{\text{N-5}})-\text{CO}-\text{O}-\text{R}_{\text{N-5}}$ where $\text{R}_{\text{N-5}}$ can be the same or different and is as defined above,

30 (22) $-(\text{CH}_2)_{0-4}-\text{N}(\text{H or } \text{R}_{\text{N-5}})-\text{CO}-\text{N}(\text{R}_{\text{N-5}})_2$, where $\text{R}_{\text{N-5}}$ can be the same or different and is as defined above,

(23) $-(\text{CH}_2)_{0-4}-\text{N}-\text{CS}-\text{N}(\text{R}_{\text{N-5}})_2$, where $\text{R}_{\text{N-5}}$ can be the same or different and is as defined above,

(24) $-(\text{CH}_2)_{0-4}-\text{N}(\text{H or } \text{R}_{\text{N-5}})-\text{CO}-\text{R}_{\text{N-2}}$ where $\text{R}_{\text{N-5}}$ and $\text{R}_{\text{N-2}}$ can be the same or different and are as defined above,

- (25) $-(\text{CH}_2)_{0-4}-\text{NR}_{\text{N}-2}\text{R}_{\text{N}-3}$ where $\text{R}_{\text{N}-2}$ and $\text{R}_{\text{N}-3}$ can be the same or different and are as defined above,
- (26) $-(\text{CH}_2)_{0-4}-\text{R}_{\text{N}-4}$ where $\text{R}_{\text{N}-4}$ is as defined above,
- (27) $-(\text{CH}_2)_{0-4}-\text{O}-\text{CO}-(\text{C}_1-\text{C}_6 \text{ alkyl})$,
- 5 (28) $-(\text{CH}_2)_{0-4}-\text{O}-\text{P}(\text{O})-(\text{OR}_{\text{N-aryl-1}})_2$ where $\text{R}_{\text{N-aryl-1}}$ is – H or C_1-C_4 alkyl,
- (29) $-(\text{CH}_2)_{0-4}-\text{O}-\text{CO}-\text{N}(\text{R}_{\text{N}-5})_2$ where $\text{R}_{\text{N}-5}$ is as defined above,
- (30) $-(\text{CH}_2)_{0-4}-\text{O}-\text{CS}-\text{N}(\text{R}_{\text{N}-5})_2$ where $\text{R}_{\text{N}-5}$ is as defined
- 10 above,
- (31) $-(\text{CH}_2)_{0-4}-\text{O}-(\text{R}_{\text{N}-5})_2$ where $\text{R}_{\text{N}-5}$ is as defined above,
- (32) $-(\text{CH}_2)_{0-4}-\text{O}-(\text{R}_{\text{N}-5})_2-\text{COOH}$ where $\text{R}_{\text{N}-5}$ is as defined above,
- 15 (33) $-(\text{CH}_2)_{0-4}-\text{S}-(\text{R}_{\text{N}-5})_2$ where $\text{R}_{\text{N}-5}$ is as defined above,
- (34) $-(\text{CH}_2)_{0-4}-\text{O}-(\text{C}_1-\text{C}_6 \text{ alkyl optionally substituted with one, two, three, four, or five of } -\text{F})$,
- (35) $\text{C}_3-\text{C}_7 \text{ cycloalkyl}$,
- 20 (36) $\text{C}_2-\text{C}_6 \text{ alkenyl}$ with one or two double bonds optionally substituted with $\text{C}_1-\text{C}_3 \text{ alkyl}$, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, $-\text{OH}$, $-\text{SH}$, $-\text{C}\equiv\text{N}$, $-\text{CF}_3$, $\text{C}_1-\text{C}_3 \text{ alkoxy}$, or $-\text{NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,
- (37) $\text{C}_2-\text{C}_6 \text{ alkynyl}$ with one or two triple bonds optionally substituted with $\text{C}_1-\text{C}_3 \text{ alkyl}$, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, $-\text{OH}$, $-\text{SH}$, $-\text{C}\equiv\text{N}$, $-\text{CF}_3$, $\text{C}_1-\text{C}_3 \text{ alkoxy}$, or $-\text{NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,
- 25 (38) $-(\text{CH}_2)_{0-4}-\text{N}(-\text{H or } \text{R}_{\text{N}-5})-\text{SO}_2-\text{R}_{\text{N}-2}$ where $\text{R}_{\text{N}-5}$ and $\text{R}_{\text{N}-2}$ can be the same or different and are as defined above, or
- (39) $-(\text{CH}_2)_{0-4}-\text{C}_3-\text{C}_7 \text{ cycloalkyl}$,
- (C) $\text{R}_{\text{N-aryl}}-\text{W}-\text{R}_{\text{N-aryl}}$, where $\text{R}_{\text{N-aryl}}$ can be the same or
- 30 different,
- (D) $\text{R}_{\text{N-aryl}}-\text{W}-\text{R}_{\text{N-heteroaryl}}$,
- (E) $\text{R}_{\text{N-aryl}}-\text{W}-\text{R}_{\text{N-1-heterocycle}}$, wherein $\text{R}_{\text{N-1-heterocycle}}$ is the same as $\text{R}_{1-\text{heterocycle}}$, and $\text{R}_{1-\text{heterocycle}}$ is as defined above

(F) $R_{N\text{-heteroaryl}}\text{-W-}R_{N\text{-aryl}}$,

(G) $R_{N\text{-heteroaryl}}\text{-W-}R_{N\text{-heteroaryl}}$,

(H) $R_{N\text{-heteroaryl}}\text{-W-}R_{N\text{-1-heterocycle}}$,

(I) $R_{N\text{-heterocycle}}\text{-W-}R_{N\text{-aryl}}$, wherein $R_{N\text{-heterocycle}}$ is the same as

5 $R_{1\text{-heterocycle}}$, and $R_{1\text{-heterocycle}}$ is as defined above, and $R_{N\text{-aryl}}$ is as defined above,

(J) $R_{N\text{-heterocycle}}\text{-W-}R_{N\text{-heteroaryl}}$, and

(K) $R_{N\text{-heterocycle}}\text{-W-}R_{N\text{-1-heterocycle}}$,

where W is

(13) $\text{-(CH}_2\text{)}_{0-4}\text{-}$,

10 (14) -O- ,

(15) $\text{-S(O)}_{0-2}\text{-}$,

(16) $\text{-N(R}_{N-5}\text{)-}$ where R_{N-5} is as defined

above, or

(5) -CO- ;

15 (II) $\text{-CO-(C}_1\text{-C}_{10}\text{ alkyl)}$ where alkyl is optionally substituted with one three substituents selected from the group consisting of:

(A) -OH ,

(B) $\text{-C}_1\text{-C}_6\text{ alkoxy}$,

(C) $\text{-C}_1\text{-C}_6\text{ thioalkoxy}$,

20 (D) $\text{-CO-O-}R_{N-8}$ where R_{N-8} is -H , $\text{C}_1\text{-C}_6\text{ alkyl}$ or -phenyl ,

(E) $\text{-CO-NR}_{N-2}R_{N-3}$ where R_{N-2} and R_{N-3} are the same or

different and are as defined above,

(F) $\text{-CO-}R_{N-4}$ where R_{N-4} is as defined above,

(G) $\text{-SO}_2\text{-(C}_1\text{-C}_8\text{ alkyl)}$,

25 (H) $\text{-SO}_2\text{-NR}_{N-2}R_{N-3}$ where R_{N-2} and R_{N-3} are the same or different and are as defined above,

(I) $\text{-NH-CO-(C}_1\text{-C}_6\text{ alkyl)}$,

(J) $\text{-NH-CO-O-}R_{N-8}$ where R_{N-8} is as defined above,

(K) $\text{-NR}_{N-2}R_{N-3}$ where R_{N-2} and R_{N-3} are the same or different

30 and are as defined above,

(L) -R_{N-4} where R_{N-4} is as defined above,

(M) $\text{-O-CO-(C}_1\text{-C}_6\text{ alkyl)}$,

(N) $\text{-O-CO-NR}_{N-8}R_{N-8}$ where R_{N-8} are the same or different

and are as defined above,

- (O) -O-(C₁-C₅ alkyl)-COOH,
(P) -O-(C₁-C₆ alkyl optionally substituted with one, two, or three of -F, -Cl, -Br, or -I),
(Q) -NH-SO₂-(C₁-C₆ alkyl), and
5 (R) -F, or -Cl,
(III) -CO-(C₁-C₆ alkyl)-O-(C₁-C₆ alkyl) where alkyl is optionally substituted with one, two, or three substituents selected from the group consisting of:
(A) -OH,
10 (B) -C₁-C₆ alkoxy,
(C) -C₁-C₆ thioalkoxy,
(D) -CO-O-R_{N-8} where R_{N-8} is -H, C₁-C₆ alkyl or -phenyl,
(E) -CO-NR_{N-2}R_{N-3} where R_{N-2} and R_{N-3} are the same or different and are as defined above,
15 (F) -CO-R_{N-4} where R_{N-4} is as defined above,
(G) -SO₂-(C₁-C₈ alkyl),
(H) -SO₂-NR_{N-2}R_{N-3} where R_{N-2} and R_{N-3} are the same or different and are as defined above,
(I) -NH-CO-(C₁-C₆ alkyl),
20 (J) -NH-CO-O-R_{N-8} where R_{N-8} is as defined above,
(K) -NR_{N-2}R_{N-3} where R_{N-2} and R_{N-3} are the same or different and are as defined above,
(L) -R_{N-4} where R_{N-4} is as defined above,
(M) -O-CO-(C₁-C₆ alkyl),
25 (N) -O-CO-NR_{N-8}R_{N-8} where R_{N-8} are the same or different and are as defined above,
(O) -O-(C₁-C₅ alkyl)-COOH,
(P) -O-(C₁-C₆ alkyl optionally substituted with one, two, or three of -F, -Cl, -Br, or -I),
30 (Q) -NH-SO₂-(C₁-C₆ alkyl), and
(R) -F, or -Cl,

(IV) $-\text{CO}-(\text{C}_1-\text{C}_6 \text{ alkyl})-\text{S}-(\text{C}_1-\text{C}_6 \text{ alkyl})$ where alkyl is optionally substituted with one, two, or three of substituents selected from the group consisting of:

- 5 (A) $-\text{OH}$,
 - (B) $-\text{C}_1-\text{C}_6 \text{ alkoxy}$,
 - (C) $-\text{C}_1-\text{C}_6 \text{ thioalkoxy}$,
 - (D) $-\text{CO}-\text{O}-\text{R}_{\text{N}-8}$ where $\text{R}_{\text{N}-8}$ is as defined above,
 - (E) $-\text{CO}-\text{NR}_{\text{N}-2}\text{R}_{\text{N}-3}$ where $\text{R}_{\text{N}-2}$ and $\text{R}_{\text{N}-3}$ are the same or different and are as defined above,
 - 10 (F) $-\text{CO}-\text{R}_{\text{N}-4}$ where $\text{R}_{\text{N}-4}$ is as defined above,
 - (G) $-\text{SO}_2-(\text{C}_1-\text{C}_8 \text{ alkyl})$,
 - (H) $-\text{SO}_2-\text{NR}_{\text{N}-2}\text{R}_{\text{N}-3}$ where $\text{R}_{\text{N}-2}$ and $\text{R}_{\text{N}-3}$ are the same or different and are as defined above,
 - (I) $-\text{NH}-\text{CO}-(\text{C}_1-\text{C}_6 \text{ alkyl})$,
 - 15 (J) $-\text{NH}-\text{CO}-\text{O}-\text{R}_{\text{N}-8}$ where $\text{R}_{\text{N}-8}$ is as defined above,
 - (K) $-\text{NR}_{\text{N}-2}\text{R}_{\text{N}-3}$ where $\text{R}_{\text{N}-2}$ and $\text{R}_{\text{N}-3}$ are the same or different and are as defined above,
 - (L) $-\text{R}_{\text{N}-4}$ where $\text{R}_{\text{N}-4}$ is as defined above,
 - (M) $-\text{O}-\text{CO}-(\text{C}_1-\text{C}_6 \text{ alkyl})$,
 - 20 (N) $-\text{O}-\text{CO}-\text{NR}_{\text{N}-8}\text{R}_{\text{N}-8}$ where $\text{R}_{\text{N}-8}$ are the same or different and are as defined above,
 - (O) $-\text{O}-(\text{C}_1-\text{C}_5 \text{ alkyl})-\text{COOH}$,
 - (P) $-\text{O}-(\text{C}_1-\text{C}_6 \text{ alkyl})$ optionally substituted with one, two, or three of $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, or $-\text{I}$,
 - 25 (Q) $-\text{NH}-\text{SO}_2-(\text{C}_1-\text{C}_6 \text{ alkyl})$, and
 - (R) $-\text{F}$, or $-\text{Cl}$,
- (V) $-\text{CO}-\text{CH}(-(\text{CH}_2)_{0-2}-\text{O}-\text{R}_{\text{N}-10})-(\text{CH}_2)_{0-2}-\text{R}_{\text{N-aryl}}/\text{R}_{\text{N-heteroaryl}}$ where $\text{R}_{\text{N-aryl}}$ and $\text{R}_{\text{N-heteroaryl}}$ are as defined above, where $\text{R}_{\text{N}-10}$ is selected from the group consisting of:
- 30 (A) $-\text{H}$,
 - (B) $\text{C}_1-\text{C}_6 \text{ alkyl}$,
 - (C) $\text{C}_3-\text{C}_7 \text{ cycloalkyl}$,
 - (D) $\text{C}_2-\text{C}_6 \text{ alkenyl}$ with one double bond,

- (E) C₂-C₆ alkynyl with one triple bond,
 (F) R_{1-aryl} where R_{1-aryl} is as defined above, and
 (G) R_{N-heteroaryl} where R_{N-heteroaryl} is as defined above, or
 (VI) -CO-(C₃-C₈ cycloalkyl) where alkyl is optionally substituted
 5 with one or two substituents selected from the group consisting of:
 (A) -(CH₂)₀₋₄-OH,
 (B) -(CH₂)₀₋₄-C₁-C₆ alkoxy,
 (C) -(CH₂)₀₋₄-C₁-C₆ thioalkoxy,
 (D) -(CH₂)₀₋₄-CO-O-R_{N-8} where R_{N-8} is -H, C₁-C₆ alkyl or
 10 phenyl,
 (E) -(CH₂)₀₋₄-CO-NR_{N-2}R_{N-3} where R_{N-2} and R_{N-3} are the
 same or different and are as defined above,
 (F) -(CH₂)₀₋₄-CO-R_{N-4} where R_{N-4} is as defined above,
 (G) -(CH₂)₀₋₄-SO₂-(C₁-C₈ alkyl),
 15 (H) -(CH₂)₀₋₄-SO₂-NR_{N-2}R_{N-3} where R_{N-2} and R_{N-3} are the
 same or different and are as defined above,
 (I) -(CH₂)₀₋₄-NH-CO-(C₁-C₆ alkyl),
 (J) -NH-CO-O-R_{N-8} where R_{N-8} is as defined above,
 (K) -(CH₂)₀₋₄-NR_{N-2}R_{N-3} where R_{N-2} and R_{N-3} are the same or
 20 different and are as defined above,
 (L) -(CH₂)₀₋₄-R_{N-4} where R_{N-4} is as defined above,
 (M) -O-CO-(C₁-C₆ alkyl),
 (N) -O-CO-NR_{N-8}R_{N-8} where R_{N-8} are the same or different
 and are as defined above,
 25 (O) -O-(C₁-C₅ alkyl)-COOH,
 (P) -O-(C₁-C₆ alkyl optionally substituted with one, two, or
 three of -F, -Cl, -Br, or -I),
 (Q) -NH-SO₂-(C₁-C₆ alkyl), and
 (R) -F, or -Cl;

30

where R_A is:

(I)-C₁-C₁₀ alkyl optionally substituted with one, two or three substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above, -OC=O NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above, -S(=O)₀₋₂ R_{1-a} where R_{1-a} is as defined above, - NR_{1-a}C=O NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above, -C=O NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above, and -S(=O)₂ NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(II) -(CH₂)₀₋₃-(C₃-C₈) cycloalkyl where cycloalkyl can be optionally substituted with one, two or three substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, -CO-OH, -CO-O-(C₁-C₄ alkyl), and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(III) -(CR_{A-x}R_{A-y})₀₋₄-R_{A-aryl} where R_{A-x} and R_{A-y} are

- (A) -H,
- (B) C₁-C₄ alkyl optionally substituted with one or two -OH,
- (C) C₁-C₄ alkoxy optionally substituted with one, two, or three of -F,
- (D) -(CH₂)₀₋₄-C₃-C₇ cycloalkyl,
- (E) C₂-C₆ alkenyl containing one or two double bonds,
- (F) C₂-C₆ alkynyl containing one or two triple bonds, or
- (G) phenyl,

and where R_{A-x} and R_{A-y} are taken together with the carbon to which they are attached to form a carbocycle of three, four, five, six or seven carbon atoms, optionally where one carbon atom is replaced by a heteroatom selected from the group consisting of -O-, -S-, -SO₂-, and -NR_{N-2}- and R_{A-aryl} is the same as R_{N-aryl},

(IV) -(CR_{A-x}R_{A-y})₀₋₄-R_{A-heteroaryl} where R_{A-heteroaryl} is the same as R_{N-heteroaryl} and R_{A-x} and R_{A-y} are as defined above,

(V) -(CR_{A-x}R_{A-y})₀₋₄-R_{A-aryl}-R_{A-aryl} where R_{A-aryl}, R_{A-x} and R_{A-y} are as defined above,

(VI) -(CR_{A-x}R_{A-y})₀₋₄-R_{A-aryl}-R_{A-heteroaryl} where R_{A-aryl}, R_{A-heteroaryl}, R_{A-x} and R_{A-y} are as defined above,

(VII) -(CR_{A-x}R_{A-y})₀₋₄-R_{A-heteroaryl}-R_{A-aryl} where R_{A-heteroaryl}, R_{A-aryl}, R_{A-x} and R_{A-y} are as defined above,

(VIII) $-(\text{CR}_{\text{A-x}}\text{R}_{\text{A-y}})_{0-4}-\text{R}_{\text{A-heteroaryl}}-\text{R}_{\text{A-heteroaryl}}$ where $\text{R}_{\text{A-heteroaryl}}$, $\text{R}_{\text{A-x}}$ and $\text{R}_{\text{A-y}}$ are as defined above,

(IX) $-(\text{CR}_{\text{A-x}}\text{R}_{\text{A-y}})_{0-4}-\text{R}_{\text{A-aryl}}-\text{R}_{\text{A-heterocycle}}$ where $\text{R}_{\text{A-heterocycle}}$ is defined as $\text{R}_{1\text{-heterocycle}}$, and where $\text{R}_{\text{A-aryl}}$, $\text{R}_{\text{A-x}}$ and $\text{R}_{\text{A-y}}$ are as defined above,

5 (X) $-(\text{CR}_{\text{A-x}}\text{R}_{\text{A-y}})_{0-4}-\text{R}_{\text{A-heteroaryl}}-\text{R}_{\text{A-heterocycle}}$ where $\text{R}_{\text{A-heteroaryl}}$, $\text{R}_{\text{A-heterocycle}}$, $\text{R}_{\text{A-x}}$ and $\text{R}_{\text{A-y}}$ are as defined above,

(XI) $-(\text{CR}_{\text{A-x}}\text{R}_{\text{A-y}})_{0-4}-\text{R}_{\text{A-heterocycle}}-\text{R}_{\text{A-aryl}}$ where $\text{R}_{\text{A-heterocycle}}$, $\text{R}_{\text{A-aryl}}$, $\text{R}_{\text{A-x}}$ and $\text{R}_{\text{A-y}}$ are as defined above,

10 (XII) $-(\text{CR}_{\text{A-x}}\text{R}_{\text{A-y}})_{0-4}-\text{R}_{\text{A-heterocycle}}-\text{R}_{\text{A-heteroaryl}}$ where $\text{R}_{\text{A-heterocycle}}$, $\text{R}_{\text{A-heteroaryl}}$, $\text{R}_{\text{A-x}}$ and $\text{R}_{\text{A-y}}$ are as defined above,

(XIII) $-(\text{CR}_{\text{A-x}}\text{R}_{\text{A-y}})_{0-4}-\text{R}_{\text{A-heterocycle}}-\text{R}_{\text{A-heterocycle}}$ where $\text{R}_{\text{A-heterocycle}}$, $\text{R}_{\text{A-x}}$ and $\text{R}_{\text{A-y}}$ are as defined above,

(XIV) $-(\text{CR}_{\text{A-x}}\text{R}_{\text{A-y}})_{0-4}-\text{R}_{\text{A-heterocycle}}$ where $\text{R}_{\text{A-heterocycle}}$, $\text{R}_{\text{A-x}}$ and $\text{R}_{\text{A-y}}$ are as defined above,

15 (XV) $-[\text{C}(\text{R}_{\text{A-1}})(\text{R}_{\text{A-2}})]_{1-3}-\text{CO}-\text{N}-(\text{R}_{\text{A-3}})_2$ where $\text{R}_{\text{A-1}}$ and $\text{R}_{\text{A-2}}$ are the same or different and are selected from the group consisting of:

(A) -H,

(B) $\text{C}_1\text{-C}_6$ alkyl, optionally substituted with one, two or three substituents selected from the group consisting of $\text{C}_1\text{-C}_3$ alkyl, -F, -Cl, -Br, -I, -OH, -SH, $\text{-C}\equiv\text{N}$, -CF_3 , $\text{C}_1\text{-C}_6$ alkoxy, -O-phenyl, and $\text{-NR}_{1\text{-a}}\text{R}_{1\text{-b}}$ where $\text{R}_{1\text{-a}}$ and $\text{R}_{1\text{-b}}$ are as defined above,

(C) $\text{C}_2\text{-C}_6$ alkenyl with one or two double bonds, optionally substituted with one, two or three substituents selected from the group consisting of $\text{C}_1\text{-C}_3$ alkyl, -F, -Cl, -Br, -I, -OH, -SH, $\text{-C}\equiv\text{N}$, -CF_3 , $\text{C}_1\text{-C}_6$ alkoxy, -O-phenyl, and $\text{-NR}_{1\text{-a}}\text{R}_{1\text{-b}}$ where $\text{R}_{1\text{-a}}$ and $\text{R}_{1\text{-b}}$ are as defined above,

(D) $\text{C}_2\text{-C}_6$ alkynyl with one or two triple bonds, optionally substituted with one, two or three substituents selected from the group consisting of $\text{C}_1\text{-C}_3$ alkyl, -F, -Cl, -Br, -I, -OH, -SH, $\text{-C}\equiv\text{N}$, -CF_3 , $\text{C}_1\text{-C}_6$ alkoxy, -O-phenyl, and $\text{-NR}_{1\text{-a}}\text{R}_{1\text{-b}}$ where $\text{R}_{1\text{-a}}$ and $\text{R}_{1\text{-b}}$ are as defined above,

30 (E) $-(\text{CH}_2)_{1-2}-\text{S}(\text{O})_{0-2}-(\text{C}_1\text{-C}_6 \text{ alkyl})$,

(F) $-(\text{CH}_2)_{0-4}-\text{C}_3\text{-C}_7$ cycloalkyl, optionally substituted with one, two or three substituents selected from the group consisting of $\text{C}_1\text{-C}_3$ alkyl, -F,

-Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(G) -(C₁-C₄ alkyl)-R_{A'-aryl} where R_{A'-aryl} is as defined for R_{1-aryl},

aryl,

5 (H) -(C₁-C₄ alkyl)-R_{A-heteroaryl} where R_{A-heteroaryl} is as defined above,

(I) -(C₁-C₄ alkyl)-R_{A-heterocycle} where R_{A-heterocycle} is as defined above,

(J) -R_{A-heteroaryl} where R_{A-heteroaryl} is as defined above,

10 (K) -R_{A-heterocycle} where R_{A-heterocycle} is as defined above,

(M) -(CH₂)₁₋₄-R_{A-4}-(CH₂)₀₋₄-R_{A'-aryl} where R_{A-4} is -O-, -S- or -NR_{A-5}- where R_{A-5} is C₁-C₆ alkyl, and where R_{A'-aryl} is defined above,

(N) -(CH₂)₁₋₄-R_{A-4}-(CH₂)₀₋₄-R_{A-heteroaryl} where R_{A-4} and R_{A-}

heteroaryl are as defined above, and

15 (O) -R_{A'-aryl} where R_{A'-aryl} is as defined above,

and where R_{A-3} is the same or different and is:

(A) -H,

(B) -C₁-C₆ alkyl optionally substituted with one, two or three substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(C) C₂-C₆ alkenyl with one or two double bonds, optionally substituted with one, two or three substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(D) C₂-C₆ alkynyl with one or two triple bonds, optionally substituted with one, two or three substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

30 (E) -(CH₂)₀₋₄-C₃-C₇ cycloalkyl, optionally substituted with one, two or three substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

- (F) $-R_{A'-aryl}$ where $R_{A'-aryl}$ is as defined above,
- (G) $-R_{A-heteroaryl}$ where $R_{A-heteroaryl}$ is as defined above,
- (H) $-R_{A-heterocycle}$ where $R_{A-heterocycle}$ is as defined above,
- (I) $-(C_1-C_4 \text{ alkyl})-R_{A'-aryl}$ where $R_{A'-aryl}$ is as defined
 5 above,
- (J) $-(C_1-C_4 \text{ alkyl})-R_{A-heteroaryl}$ where $R_{A-heteroaryl}$ is as defined
 above,
- (K) $-(C_1-C_4 \text{ alkyl})-R_{A-heterocycle}$ where $R_{A-heterocycle}$ is as defined
 above, or
- 10 (XVI) $-\text{CH}(R_{A-aryl})_2$ where R_{A-aryl} are the same or different and are as
 defined above,
- (XVII) $-\text{CH}(R_{A-heteroaryl})_2$ where $R_{A-heteroaryl}$ are the same or different
 and are as defined above,
- (XVIII) $-\text{CH}(R_{A-aryl})(R_{A-heteroaryl})$ where R_{A-aryl} and $R_{A-heteroaryl}$ are as
 15 defined above,
- (XIX) -cyclopentyl, -cyclohexyl, or -cycloheptyl ring fused to R_{A-aryl} ,
 $R_{A-heteroaryl}$, $R_{A-heterocycle}$ where R_{A-aryl} or $R_{A-heteroaryl}$ or $R_{A-heterocycle}$ are as defined above
 where one carbon of cyclopentyl, cyclohexyl, or -cycloheptyl is optionally replaced
 with NH, NR_{N-5} , O, or $\text{S}(=\text{O})_{0-2}$, and where cyclopentyl, cyclohexyl, or -cycloheptyl
 20 can be optionally substituted with one or two $-C_1-C_3$ alkyl, -F, -OH, -SH, $-\text{C}\equiv\text{N}$, -
 CF_3 , C_1-C_6 alkoxy, =O, or $-\text{NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,
- (XX) C_2-C_{10} alkenyl containing one or two double bonds optionally
 substituted with one, two or three substituents selected from the group consisting of
 C_1-C_3 alkyl, -F, -Cl, -Br, -I, -OH, -SH, $-\text{C}\equiv\text{N}$, $-\text{CF}_3$, C_1-C_6 alkoxy, -O-phenyl, and -
 25 $\text{NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,
- (XXI) C_2-C_{10} alkynyl containing one or two triple bonds optionally
 substituted with one, two or three substituents selected from the group consisting of
 C_1-C_3 alkyl, -F, -Cl, -Br, -I, -OH, -SH, $-\text{C}\equiv\text{N}$, $-\text{CF}_3$, C_1-C_6 alkoxy, -O-phenyl, $-\text{NR}_{1-a}\text{R}_{1-b}$
 where R_{1-a} and R_{1-b} are as defined above,
- 30 (XXI) $-(\text{CH}_2)_{0-1}-\text{CHR}_{A-6}-(\text{CH}_2)_{0-1}-R_{A-aryl}$ where R_{A-aryl} is as defined
 above and R_{A-6} is $-(\text{CH}_2)_{0-6}-\text{OH}$,
- (XXII) $-(\text{CH}_2)_{0-1}-\text{CHR}_{A-6}-(\text{CH}_2)_{0-1}-R_{A-heteroaryl}$ where $R_{A-heteroaryl}$ and
 R_{A-6} is as defined above,

(XXIII) $-\text{CH}(-\text{R}_{\text{A-aryl}}$ or $\text{R}_{\text{A-heteroaryl}})-\text{CO}-\text{O}(\text{C}_1-\text{C}_4 \text{ alkyl})$ where $\text{R}_{\text{A-aryl}}$

and $\text{R}_{\text{A-heteroaryl}}$ are as defined above,

(XXIV) $-\text{CH}(-\text{CH}_2-\text{OH})-\text{CH}(-\text{OH})-\text{micro-NO}_2$,

(XXV) $(\text{C}_1-\text{C}_6 \text{ alkyl})-\text{O}-(\text{C}_1-\text{C}_6 \text{ alkyl})-\text{OH}$,

5 (XXVII) $-\text{CH}_2-\text{NH}-\text{CH}_2-\text{CH}(-\text{O}-\text{CH}_2-\text{CH}_3)_2$,

(XXVIII) $-\text{H}$,

(XXIX) $-(\text{CH}_2)_{0-6}-\text{C}(=\text{NR}_{1-a})(\text{NR}_{1-a}\text{R}_{1-b})$ where R_{1-a} and R_{1-b} are as

defined above; or

(XXX)

10 $-\text{C}=\text{OC}(\text{HR}_6)\text{NHR}_7$, where R_6 and R_7 are as defined below,

$-\text{C}=\text{OR}_7$, where R_7 is as defined below,

$-\text{C}=\text{OOR}_7$, where R_7 is as defined below, or

$-\text{SOOR}_7$ where R_7 is as defined below,

wherein R_6 is:

15 hydrogen,

$\text{C}_1 - \text{C}_3$ alkyl,

phenyl,

thioalkoxyalkyl,

alkyl substituted aryl,

20 cycloalkyl,

cycloalkylalkyl,

hydroxyalkyl,

alkoxyalkyl,

aryloxyalkyl,

25 haloalkyl,

carboxyalkyl,

alkoxycarbonylalkyl,

aminoalkyl,

(N-protected)aminoalkyl,

30 alkylaminoalkyl,

((N-protected)(alkyl)amino)alkyl,

dialkylaminoalkyl,

guanidinoalkyl,

lower alkenyl,

- heterocyclic,
(heterocyclic)alkyl),
arylthioalkyl,
arylsulfonylalkyl,
5 (heterocyclic)thioalkyl,
(heterocyclic)sulfonylalkyl,
(heterocyclic)oxyalkyl,
arylalkoxyalkyl,
arylthioalkoxyalkyl,
10 arylalkylsulfonylalkyl,
(heterocyclic))alkoxyalkyl,
(heterocyclic)thioalkoxyalkyl,
(heterocyclic)alkylsulfonylalkyl,
cycloalkyloxyalkyl,
15 cycloalkylthioalkyl,
cycloalkylsulfonylalkyl,
cycloalkylalkoxyalkyl,
cycloalkylthioalkoxyalkyl,
cycloalkylalkylsulfonylalkyl,
20 aminocarbonyl,
alkylaminocarbonyl,
dialkylaminocarbonyl,
aroalkyl,
(heterocyclic)carbonylalkyl,
25 polyhydroxyalkyl,
aminocarbonylalkyl,
alkylaminocarbonylalkyl,
dialkylaminocarbonylalkyl,
aryloxyalkyl, or
30 alkylsulfonylalkyl,

wherein heterocyclic is pyridyl, thiazolyl, isothiazolyl, oxazolyl, isoxazolyl, furanyl, thienyl, tetrahydrofuranyl, tetrahydrothienyl and tetrahydro[2H]pyranyl and wherein the heterocycle is unsubstituted or substituted with one to three substituents independently selected from hydroxy, halo, amino,

alkylamino, dialkylamino, alkoxy, polyalkoxy, haloalkyl, cycloalkyl, cycloalkylalkyl, aryl, arylalkyl, COOH, -SO₃H, lower alkenyl or lower alkyl;

wherein R₇ is:

- 5 C₁ - C₃ alkyl,
phenyl,
thioalkoxyalkyl,
(aryl)alkyl,
cycloalkyl,
cycloalkylalkyl,
10 hydroxyalkyl,
alkoxyalkyl,
aryloxyalkyl,
haloalkyl,
carboxyalkyl,
15 alkoxycarbonylalkyl,
aminoalkyl,
(N-protected)aminocalkyl,
alkylaminoalkyl,
((N-protected)(alkyl)amino)alkyl,
20 dialkylaminoalkyl,
guanidinoalkyl,
lower alkenyl,
heterocyclic,
(heterocyclic)alkyl),
25 arylthioalkyl,
arylsulfonylalkyl,
(heterocyclic)thioalkyl,
(heterocyclic)sulfonylalkyl,
(heterocyclic)oxyalkyl,
30 arylalkoxyalkyl,
arylthioalkoxyalkyl,
arylalkylsulfonylalkyl,
(heterocyclic))alkoxyalkyl,
(heterocyclic)thioalkoxyalkyl,

(heterocyclic)alkylsulfonylalkyl,
 cycloalkyloxyalkyl,
 cycloalkylthioalkyl,
 cycloalkylsulfonylalkyl,
 5 cycloalkylalkoxyalkyl,
 cycloalkylthioalkoxyalkyl,
 cycloalkylalkylsulfonylalkyl,
 aminocarbonyl,
 alkylaminocarbonyl,
 10 dialkylaminocarbonyl,
 aroylalkyl,
 (heterocyclic)carbonylalkyl,
 polyhydroxyalkyl,
 aminocarbonylalkyl,
 15 alkylaminocarbonylalkyl,
 dialkylaminocarbonylalkyl,
 aryloxyalkyl, or
 alkylsulfonylalkyl,
 wherein heterocyclic is pyridyl, thiazolyl, isothiazolyl,
 20 oxazolyl, isoxazolyl, furanyl, thienyl, tetrahydrofuranyl, tetrahydrothienyl and
 tetrahydro[2H]pyranyl and wherein the heterocycle is unsubstituted or substituted
 with one to three substituents independently selected from hydroxy, halo, amino,
 alkylamino, dialkylamino, alkoxy, polyalkoxy, haloalkyl, cycloalkyl,
 cycloalkylalkyl, aryl, arylalkyl, COOH, -SO₃H, lower alkenyl or lower alkyl;

25

where R_B is:

(I)-C₁-C₁₀ alkyl optionally substituted with one, two or three
 substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH,
 -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as
 30 defined above, -OC=O NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above, -S(=O)₀₋₂
 R_{1-a} where R_{1-a} is as defined above, -NR_{1-a}C=O NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as
 defined above, -C=O NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above, and -
 S(=O)₂ NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(II) $-(CH_2)_{0-3}-(C_3-C_8)$ cycloalkyl where cycloalkyl can be optionally substituted with one, two or three substituents selected from the group consisting of C_1-C_3 alkyl, $-F$, $-Cl$, $-Br$, $-I$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, C_1-C_6 alkoxy, $-O$ -phenyl, $-CO-OH$, $-CO-O-(C_1-C_4 \text{ alkyl})$, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

5 (III) $-(CR_{B-x}R_{B-y})_{0-4}-R_{B-aryl}$ where R_{B-x} and R_{B-y} are

(A) $-H$,

(B) C_1-C_4 alkyl optionally substituted with one or two $-OH$,

(C) C_1-C_4 alkoxy optionally substituted with one, two, or three of $-F$,

10 (D) $-(CH_2)_{0-4}-C_3-C_7$ cycloalkyl,

(E) C_2-C_6 alkenyl containing one or two double bonds,

(F) C_2-C_6 alkynyl containing one or two triple bonds, or

(G) phenyl,

and where R_{B-x} and R_{B-y} are taken together with the carbon to which they are attached to form a carbocycle of three, four, five, six or seven carbon atoms, optionally where one carbon atom is replaced by a heteroatom selected from the group consisting of $-O-$, $-S-$, $-SO_2-$, and $-NR_{N-2}$ where R_{N-2} is as defined above, and R_{B-aryl} is the same as R_{N-aryl} and is defined above

(IV) $-(CR_{B-x}R_{B-y})_{0-4}-R_{B-heteroaryl}$ where $R_{B-heteroaryl}$ is the same as $R_{N-heteroaryl}$, R_{B-x} , and R_{B-y} are as defined above,

(V) $-(CR_{B-x}R_{B-y})_{0-4}-R_{B-aryl}-R_{B-aryl}$ where R_{B-aryl} , R_{B-x} , and R_{B-y} are as defined above,

(VI) $-(CR_{B-x}R_{B-y})_{0-4}-R_{B-aryl}-R_{B-heteroaryl}$ where R_{B-aryl} , $R_{B-heteroaryl}$, R_{B-x} and R_{B-y} are as defined above,

25 (VII) $-(CR_{B-x}R_{B-y})_{0-4}-R_{B-heteroaryl}-R_{B-aryl}$ where $R_{B-heteroaryl}$, R_{B-aryl} , R_{B-x} and R_{B-y} are as defined above,

(VIII) $-(CR_{B-x}R_{B-y})_{0-4}-R_{B-heteroaryl}-R_{B-heteroaryl}$ where $R_{B-heteroaryl}$, R_{B-x} and R_{B-y} are as defined above,

(IX) $-(CR_{B-x}R_{B-y})_{0-4}-R_{B-aryl}-R_{B-heterocycle}$ where $R_{B-heterocycle}$ is defined as $R_{1-heterocycle}$, and where R_{B-aryl} , R_{B-x} and R_{B-y} are as defined above,

(X) $-(CR_{B-x}R_{B-y})_{0-4}-R_{B-heteroaryl}-R_{B-heterocycle}$ where $R_{B-heteroaryl}$, $R_{B-heterocycle}$, R_{B-x} and R_{B-y} are as defined above,

(XI) $-(\text{CR}_{\text{B-x}}\text{R}_{\text{B-y}})_{0-4}-\text{R}_{\text{B-heterocycle}}-\text{R}_{\text{B-aryl}}$ where $\text{R}_{\text{B-heterocycle}}$, $\text{R}_{\text{B-aryl}}$, $\text{R}_{\text{B-x}}$ and $\text{R}_{\text{B-y}}$ are as defined above,

(XII) $-(\text{CR}_{\text{B-x}}\text{R}_{\text{B-y}})_{0-4}-\text{R}_{\text{B-heterocycle}}-\text{R}_{\text{B-heteroaryl}}$ where $\text{R}_{\text{B-heterocycle}}$, $\text{R}_{\text{B-heteroaryl}}$, $\text{R}_{\text{B-x}}$ and $\text{R}_{\text{B-y}}$ are as defined above,

5 (XIII) $-(\text{CR}_{\text{B-x}}\text{R}_{\text{B-y}})_{0-4}-\text{R}_{\text{B-heterocycle}}-\text{R}_{\text{B-heterocycle}}$ where $\text{R}_{\text{B-heterocycle}}$, $\text{R}_{\text{B-x}}$ and $\text{R}_{\text{B-y}}$ are as defined above,

(XIV) $-(\text{CR}_{\text{B-x}}\text{R}_{\text{B-y}})_{0-4}-\text{R}_{\text{B-heterocycle}}$ where $\text{R}_{\text{B-heterocycle}}$, $\text{R}_{\text{B-x}}$ and $\text{R}_{\text{B-y}}$ are as defined above,

(XV) $-\text{[C(R}_{\text{B-1}})(\text{R}_{\text{B-2}})]_{1-3}-\text{CO-N}(\text{R}_{\text{B-3}})_2$ where $\text{R}_{\text{B-1}}$ and $\text{R}_{\text{B-2}}$ are the
10 same or different and are selected from the group consisting of:

(A) -H,

(B) $\text{C}_1\text{-C}_6$ alkyl, optionally substituted with one, two or three substituents selected from the group consisting of $\text{C}_1\text{-C}_3$ alkyl, -F, -Cl, -Br, -I, -OH, -SH, $\text{-C}\equiv\text{N}$, -CF_3 , $\text{C}_1\text{-C}_6$ alkoxy, -O-phenyl, and $\text{-NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are as
15 defined above,

(C) $\text{C}_2\text{-C}_6$ alkenyl with one or two double bonds, optionally substituted with one, two or three substituents selected from the group consisting of $\text{C}_1\text{-C}_3$ alkyl, -F, -Cl, -Br, -I, -OH, -SH, $\text{-C}\equiv\text{N}$, -CF_3 , $\text{C}_1\text{-C}_6$ alkoxy, -O-phenyl, and $\text{-NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

20 (D) $\text{C}_2\text{-C}_6$ alkynyl with one or two triple bonds, optionally substituted with one, two or three substituents selected from the group consisting of $\text{C}_1\text{-C}_3$ alkyl, -F, -Cl, -Br, -I, -OH, -SH, $\text{-C}\equiv\text{N}$, -CF_3 , $\text{C}_1\text{-C}_6$ alkoxy, -O-phenyl, and $\text{-NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

(E) $-(\text{CH}_2)_{1-2}-\text{S(O)}_{0-2}-(\text{C}_1\text{-C}_6 \text{ alkyl})$,

25 (F) $-(\text{CH}_2)_{0-4}-\text{C}_3\text{-C}_7$ cycloalkyl, optionally substituted with one, two or three substituents selected from the group consisting of $\text{C}_1\text{-C}_3$ alkyl, -F, -Cl, -Br, -I, -OH, -SH, $\text{-C}\equiv\text{N}$, -CF_3 , $\text{C}_1\text{-C}_6$ alkoxy, -O-phenyl, and $\text{-NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

(G) $-(\text{C}_1\text{-C}_4 \text{ alkyl})-\text{R}_{\text{B'-aryl}}$ where $\text{R}_{\text{B'-aryl}}$ is as defined above for
30 $\text{R}_{1\text{-aryl}}$,

(H) $-(\text{C}_1\text{-C}_4 \text{ alkyl})-\text{R}_{\text{B-heteroaryl}}$ where $\text{R}_{\text{B-heteroaryl}}$ is as defined above,

(I) $-(C_1-C_4 \text{ alkyl})-R_{B\text{-heterocycle}}$ where $R_{B\text{-heterocycle}}$ is as defined above,

(J) $-R_{B\text{-heteroaryl}}$ where $R_{B\text{-heteroaryl}}$ is as defined above,

(K) $-R_{B\text{-heterocycle}}$ where $R_{B\text{-heterocycle}}$ is as defined above,

5 (M) $-(CH_2)_{1-4}-R_{B-4}-(CH_2)_{0-4}-R_{B'\text{-aryl}}$ where R_{B-4} is $-O-$, $-S-$ or $-NR_{B-5}-$ where R_{B-5} is C_1-C_6 alkyl, and where $R_{B'\text{-aryl}}$ is defined above,

(N) $-(CH_2)_{1-4}-R_{B-4}-(CH_2)_{0-4}-R_{B\text{-heteroaryl}}$ where R_{B-4} and $R_{B\text{-heteroaryl}}$ are as defined above, and

(O) $-R_{B'\text{-aryl}}$ where $R_{B'\text{-aryl}}$ is as defined above,

10 and where R_{B-3} is the same or different and is:

(A) $-H$,

(B) $-C_1-C_6$ alkyl optionally substituted with one, two or three substituents selected from the group consisting of C_1-C_3 alkyl, $-F$, $-Cl$, $-Br$, $-I$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, C_1-C_6 alkoxy, $-O$ -phenyl, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

(C) C_2-C_6 alkenyl with one or two double bonds, optionally substituted with one, two or three substituents selected from the group consisting of C_1-C_3 alkyl, $-F$, $-Cl$, $-Br$, $-I$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, C_1-C_6 alkoxy, $-O$ -phenyl, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

20 (D) C_2-C_6 alkynyl with one or two triple bonds, optionally substituted with one, two or three substituents selected from the group consisting of C_1-C_3 alkyl, $-F$, $-Cl$, $-Br$, $-I$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, C_1-C_6 alkoxy, $-O$ -phenyl, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

(E) $-(CH_2)_{0-4}-C_3-C_7$ cycloalkyl, optionally substituted with one, two or three substituents selected from the group consisting of C_1-C_3 alkyl, $-F$, $-Cl$, $-Br$, $-I$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, C_1-C_6 alkoxy, $-O$ -phenyl, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

(F) $-R_{B'\text{-aryl}}$ where $R_{B'\text{-aryl}}$ is as defined above,

(G) $-R_{B\text{-heteroaryl}}$ where $R_{B\text{-heteroaryl}}$ is as defined above,

30 (H) $-R_{B\text{-heterocycle}}$ where $R_{B\text{-heterocycle}}$ is as defined above,

(I) $-(C_1-C_4 \text{ alkyl})-R_{B'\text{-aryl}}$ where $R_{B'\text{-aryl}}$ is as defined above,

(J) $-(C_1-C_4 \text{ alkyl})-R_{B\text{-heteroaryl}}$ where $R_{B\text{-heteroaryl}}$ is as defined above,

(K) $-(C_1-C_4 \text{ alkyl})-R_{B\text{-heterocycle}}$ where $R_{B\text{-heterocycle}}$ is as defined above, or

5 (XVI) $-\text{CH}(R_{B\text{-aryl}})_2$ where $R_{B\text{-aryl}}$ are the same or different and are as defined above,

(XVII) $-\text{CH}(R_{B\text{-heteroaryl}})_2$ where $R_{B\text{-heteroaryl}}$ are the same or different and are as defined above,

10 (XVIII) $-\text{CH}(R_{B\text{-aryl}})(R_{B\text{-heteroaryl}})$ where $R_{B\text{-aryl}}$ and $R_{B\text{-heteroaryl}}$ are as defined above,

(XIX) -cyclopentyl, -cyclohexyl, or -cycloheptyl ring fused to $R_{B\text{-aryl}}$ or $R_{B\text{-heteroaryl}}$ or $R_{B\text{-heterocycle}}$ where $R_{B\text{-aryl}}$ or $R_{B\text{-heteroaryl}}$ or $R_{B\text{-heterocycle}}$ are as defined above where one carbon of cyclopentyl, cyclohexyl, or -cycloheptyl is optionally replaced with NH, NR_{N-5} , O, or $\text{S}(=\text{O})_{0-2}$, and where cyclopentyl, cyclohexyl, or -cycloheptyl can be optionally substituted with one or two $-C_1-C_3$ alkyl, -F, -OH, -SH, $-\text{C}\equiv\text{N}$, $-\text{CF}_3$, C_1-C_6 alkoxy, =O, or $-\text{NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

20 (XX) C_2-C_{10} alkenyl containing one or two double bonds optionally substituted with one, two or three substituents selected from the group consisting of C_1-C_3 alkyl, -F, -Cl, -Br, -I, -OH, -SH, $-\text{C}\equiv\text{N}$, $-\text{CF}_3$, C_1-C_6 alkoxy, -O-phenyl, and $-\text{NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

(XXI) C_2-C_{10} alkynyl containing one or two triple bonds optionally substituted with one, two or three substituents selected from the group consisting of C_1-C_3 alkyl, -F, -Cl, -Br, -I, -OH, -SH, $-\text{C}\equiv\text{N}$, $-\text{CF}_3$, C_1-C_6 alkoxy, -O-phenyl, and $-\text{NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

(XXI) $-(\text{CH}_2)_{0-1}-\text{CHR}_{C-6}-(\text{CH}_2)_{0-1}-R_{B\text{-aryl}}$ where $R_{B\text{-aryl}}$ is as defined above and R_{C-6} is $-(\text{CH}_2)_{0-6}-\text{OH}$,

(XXII) $-(\text{CH}_2)_{0-1}-\text{CHR}_{B-6}-(\text{CH}_2)_{0-1}-R_{B\text{-heteroaryl}}$ where $R_{B\text{-heteroaryl}}$ and R_{C-6} is as defined above,

30 (XXIII) $-\text{CH}(-R_{B\text{-aryl}} \text{ or } R_{B\text{-heteroaryl}})-\text{CO}-\text{O}(C_1-C_4 \text{ alkyl})$ where $R_{B\text{-aryl}}$ and $R_{B\text{-heteroaryl}}$ are as defined above,

(XXIV) $-\text{CH}(-\text{CH}_2-\text{OH})-\text{CH}(-\text{OH})-\text{micro-NO}_2$,

(XXV) $(C_1-C_6 \text{ alkyl})-\text{O}-(C_1-C_6 \text{ alkyl})-\text{OH}$,

(XXVII) $-\text{CH}_2\text{-NH-CH}_2\text{-CH(-O-CH}_2\text{-CH}_3)_2$,

(XXVIII) $-\text{H}$, or

(XXIX) $-(\text{CH}_2)_{0-6}\text{-C(=NR}_{1-a})(\text{NR}_{1-a}\text{R}_{1-b})$ where R_{1-a} and R_{1-b} are as defined above; and

5 where PROTECTING GROUP is selected from the group consisting of *t*-butoxycarbonyl, benzyloxycarbonyl, formyl, trityl, acetyl, trichloroacetyl, dichloroacetyl, chloroacetyl, trifluoroacetyl, difluoroacetyl, fluoroacetyl, 4-phenylbenzyloxycarbonyl, 2-methylbenzyloxycarbonyl, 4-ethoxybenzyloxycarbonyl, 4-fluorobenzyloxycarbonyl, 4-chlorobenzyloxycarbonyl,
10 3-chlorobenzyloxycarbonyl, 2-chlorobenzyloxycarbonyl, 2,4-dichlorobenzyloxycarbonyl, 4-bromobenzyloxycarbonyl, 3-bromobenzyloxycarbonyl, 4-nitrobenzyloxycarbonyl, 4-cyanobenzyloxycarbonyl, 2-(4-xenyl)isopropoxycarbonyl, 1,1-diphenyleth-1-yloxycarbonyl, 1,1-diphenylprop-1-yloxycarbonyl, 2-phenylprop-2-yloxycarbonyl, 2-(*p*-toluyl)prop-2-yloxycarbonyl,
15 cyclopentanyloxycarbonyl, 1-methylcyclopentanyloxycarbonyl, cyclohexanyloxycarbonyl, 1-methylcyclohexanyloxycarbonyl, 2-methylcyclohexanyloxycarbonyl, 2-(4-toluylsulfonyl)ethoxycarbonyl, 2-(methylsulfonyl)ethoxycarbonyl, 2-(triphenylphosphino)ethoxycarbonyl, fluorenylmethoxycarbonyl, 2-(trimethylsilyl)ethoxycarbonyl, allyloxycarbonyl, 1-(trimethylsilylmethyl)prop-1-enyloxycarbonyl, 5-benzisoxalylmethoxycarbonyl, 4-acetoxybenzyloxycarbonyl, 2,2,2-trichloroethoxycarbonyl, 2-ethynyl-2-propoxycarbonyl, cyclopropylmethoxycarbonyl, 4-(decyloxyl)benzyloxycarbonyl, isobornyloxycarbonyl and 1-piperidyloxycarbonyl, 9-fluorenylmethyl carbonate, $-\text{CH-CH=CH}_2$ and phenyl- C(=N-)-H .

25

42. A protected compound according to claim 41 where R_1 is:

$-(\text{CH}_2)_{0-1}\text{-(R}_{1\text{-aryl}})$, or

$-(\text{CH}_2)_{n1}\text{-(R}_{1\text{-heteroaryl}})$;

where R_A is:

30

$-\text{C}_1\text{-C}_8$ alkyl,

$-(\text{CH}_2)_{0-3}\text{-(C}_3\text{-C}_7)$ cycloalkyl,

$-(\text{CR}_{A-x}\text{R}_{A-y})_{0-4}\text{-R}_{A\text{-aryl}}$,

$-(\text{CR}_{A-x}\text{R}_{A-y})_{0-4}\text{-R}_{A\text{-heteroaryl}}$,

$-(\text{CR}_{A-x}\text{R}_{A-y})_{0-4}\text{-R}_{A\text{-heterocycle}}$,

-cyclopentyl or -cyclohexyl ring fused to R_{A-aryl} or $R_{A-heteroaryl}$ or $R_{A-heterocycle}$; and

where R_B is:

- C₁-C₈ alkyl,
- 5 -(CH₂)₀₋₃-(C₃-C₇) cycloalkyl,
- (CR_{A-x}R_{A-y})₀₋₄-R_{A-aryl},
- (CR_{A-x}R_{A-y})₀₋₄-R_{A-heteroaryl},
- (CR_{A-x}R_{A-y})₀₋₄-R_{A-heterocycle},

-cyclopentyl or -cyclohexyl ring fused to R_{A-aryl} or $R_{A-heteroaryl}$ or $R_{A-heterocycle}$.

10

43. A protected compound according to claim 42,

where R_1 is:

- (CH₂)-(R_{1-aryl}), or
- (CH₂)-(R_{1-heteroaryl});

15

where R_2 is -H;

where R_3 is -H;

where R_N is:

R_{N-1} -X_N- where X_N is:

-CO-,

20

where R_{N-1} is selected from the group consisting of:

-R_{N-aryl}, and

-R_{N-heteroaryl};

where R_A is:

- C₁-C₈ alkyl,
- 25 -(CH₂)₀₋₃-(C₃-C₇) cycloalkyl,
- (CR_{A-x}R_{A-y})₀₋₄-R_{A-aryl},
- (CR_{A-x}R_{A-y})₀₋₄-R_{A-heteroaryl},
- (CR_{A-x}R_{A-y})₀₋₄-R_{A-heterocycle},

-cyclopentyl or -cyclohexyl ring fused to R_{A-aryl} or $R_{A-heteroaryl}$ or $R_{A-heterocycle}$;

30

heterocycle;

where R_B is:

- C₁-C₈ alkyl,
- (CH₂)₀₋₃-(C₃-C₇) cycloalkyl,
- (CR_{B-x}R_{B-y})₀₋₄-R_{B-aryl},

$-(\text{CR}_{\text{B-x}}\text{R}_{\text{B-y}})_{0-4}-\text{R}_{\text{B-heteroaryl}},$

$-(\text{CR}_{\text{B-x}}\text{R}_{\text{B-y}})_{0-4}-\text{R}_{\text{B-heterocycle}},$

-cyclopentyl or -cyclohexyl ring fused to $\text{R}_{\text{B-aryl}}$ or $\text{R}_{\text{B-heteroaryl}}$ or $\text{R}_{\text{B-}}$


heterocycle.

5

44. A protected compound according to claim 41 where PROTECTING GROUP is *t*-butoxycarbonyl.

45. A protected compound according to claim 41 where PROTECTING GROUP is
10 benzyloxycarbonyl.

46. A protected compound of the formula (XI)



15

where R_1 is:

(I) $\text{C}_1\text{-C}_6$ alkyl, optionally substituted with one, two or three
substituents selected from the group consisting of $\text{C}_1\text{-C}_3$ alkyl, $\text{C}_1\text{-C}_7$ alkyl
(optionally substituted with $\text{C}_1\text{-C}_3$ alkyl and $\text{C}_1\text{-C}_3$ alkoxy), -F, -Cl, -Br, -I, -OH, -
20 SH, -C \equiv N, -CF₃, $\text{C}_1\text{-C}_3$ alkoxy, -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are -H or $\text{C}_1\text{-C}_6$ alkyl,
and -OC=O NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(II) $-\text{CH}_2\text{-S(O)}_{0-2}\text{-(C}_1\text{-C}_6\text{ alkyl)},$

(III) $-\text{CH}_2\text{-CH}_2\text{-S(O)}_{0-2}\text{-(C}_1\text{-C}_6\text{ alkyl)},$

(IV) $\text{C}_2\text{-C}_6$ alkenyl with one or two double bonds, optionally
25 substituted with one, two or three substituents selected from the group consisting of
-F, -Cl, -OH, -SH, -C \equiv N, -CF₃, $\text{C}_1\text{-C}_3$ alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are
-H or $\text{C}_1\text{-C}_6$ alkyl,

(V) $\text{C}_2\text{-C}_6$ alkynyl with one or two triple bonds, optionally substituted
with one, two or three substituents selected from the group consisting of -F, -Cl, -
30 OH, -SH, -C \equiv N, -CF₃, $\text{C}_1\text{-C}_3$ alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are -H or
 $\text{C}_1\text{-C}_6$ alkyl,

(VI) $-(CH_2)_{n1}-(R_{1-aryl})$ where n_1 is zero or one and where R_{1-aryl} is phenyl, 1-naphthyl, 2-naphthyl and indanyl, indenyl, dihydronaphthalyl, or tetralinyl optionally substituted with one, two, three or four of the following substituents on the aryl ring:

5 (A) C_1-C_6 alkyl optionally substituted with one, two or three substituents selected from the group consisting of C_1-C_3 alkyl, $-F$, $-Cl$, $-Br$, $-I$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, and C_1-C_3 alkoxy, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

(B) C_2-C_6 alkenyl with one or two double bonds, optionally
10 substituted with one, two or three substituents selected from the group consisting of $-F$, $-Cl$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, C_1-C_3 alkoxy, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are $-H$ or C_1-C_6 alkyl,

(C) C_2-C_6 alkynyl with one or two triple bonds, optionally
substituted with one, two or three substituents selected from the group consisting of
15 $-F$, $-Cl$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, C_1-C_3 alkoxy, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are $-H$ or C_1-C_6 alkyl,

(D) $-F$, Cl , $-Br$ or $-I$,

(F) $-C_1-C_6$ alkoxy optionally substituted with one, two or
three of $-F$,

20 (G) $-NR_{N-2}R_{N-3}$ where R_{N-2} and R_{N-3} are as defined below,

(H) $-OH$,

(I) $-C\equiv N$,

(J) C_3-C_7 cycloalkyl, optionally substituted with one, two or
three substituents selected from the group consisting of $-F$, $-Cl$, $-OH$, $-SH$, $-C\equiv N$, $-$
25 CF_3 , C_1-C_3 alkoxy, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are $-H$ or C_1-C_6 alkyl,

(K) $-CO-(C_1-C_4 \text{ alkyl})$,

(L) $-SO_2-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

(M) $-CO-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

or

30 (N) $-SO_2-(C_1-C_4 \text{ alkyl})$,

(VII) $-(CH_2)_{n1}-(R_{1-heteroaryl})$ where n_1 is as defined above and where
 $R_{1-heteroaryl}$ is selected from the group consisting of:

pyridinyl,

	pyrimidinyl,
	quinolinyl,
	benzothienyl,
	indolyl,
5	indolinyl,
	pyridazinyl,
	pyrazinyl,
	isoquinolyl,
	quinazolinyl,
10	quinoxalinyl,
	phthalazinyl,
	imidazolyl,
	isoxazolyl,
	pyrazolyl,
15	oxazolyl,
	thiazolyl,
	indolizinyll,
	indazolyl,
	benzothiazolyl,
20	benzimidazolyl,
	benzofuranyl,
	furanyl,
	thienyl,
	pyrrolyl,
25	oxadiazolyl,
	thiadiazolyl,
	triazolyl,
	tetrazolyl,
	oxazolopyridinyl,
30	imidazopyridinyl,
	isothiazolyl,
	naphthyridinyl,
	cinnolinyl,
	carbazolyl,

	beta-carbolinyl,
	isochroman-yl,
	chroman-yl,
	tetrahydroisoquinolin-yl,
5	isoindolin-yl,
	isobenzotetrahydrofuran-yl,
	isobenzotetrahydrothien-yl,
	isobenzothien-yl,
	benzoxazol-yl,
10	pyridopyridin-yl,
	benzotetrahydrofuran-yl,
	benzotetrahydrothien-yl,
	purin-yl,
	benzodioxol-yl,
15	triazin-yl,
	phenoxazin-yl,
	phenothiazin-yl,
	pteridin-yl,
	benzothiazol-yl,
20	imidazopyridin-yl,
	imidazothiazol-yl,
	dihydrobenzisoxazin-yl,
	benzisoxazin-yl,
	benzoxazin-yl,
25	dihydrobenziso-thiazin-yl,
	benzopyran-yl,
	benzothiopyran-yl,
	coumarin-yl,
	isocoumarin-yl,
30	chromon-yl,
	chromanon-yl,
	pyridin-yl-N-oxide,
	tetrahydroquinolin-yl
	dihydroquinolin-yl

	dihydroquinolinonyl
	dihydroisoquinolinonyl
	dihydrocoumarinyl
	dihydroisocoumarinyl
5	isoindolinonyl
	benzodioxanyl
	benzoxazolinonyl
	pyrrolyl N-oxide,
	pyrimidinyl N-oxide,
10	pyridazinyl N-oxide,
	pyrazinyl N-oxide,
	quinolinyl N-oxide,
	indolyl N-oxide,
	indolinyl N-oxide,
15	isoquinolyl N-oxide,
	quinazolinyl N-oxide,
	quinoxalinyl N-oxide,
	phthalazinyl N-oxide,
	imidazolyl N-oxide,
20	isoxazolyl N-oxide,
	oxazolyl N-oxide,
	thiazolyl N-oxide,
	indoliziny N-oxide,
	indazolyl N-oxide,
25	benzothiazolyl N-oxide,
	benzimidazolyl N-oxide,
	pyrrolyl N-oxide,
	oxadiazolyl N-oxide,
	thiadiazolyl N-oxide,
30	triazolyl N-oxide,
	tetrazolyl N-oxide,
	benzothiopyranyl S-oxide, and
	benzothiopyranyl S,S-dioxide,

where the $R_{1\text{-heteroaryl}}$ group is bonded to $-(CH_2)_{n1}-$ by any ring atom of the parent $R_{1\text{-heteroaryl}}$ group substituted by hydrogen such that the new bond to the $R_{1\text{-heteroaryl}}$ group replaces the hydrogen atom and its bond, where heteroaryl is optionally substituted with one, two, three or four of:

- 5 (1) C_1-C_6 alkyl optionally substituted with one, two or three substituents selected from the group consisting of C_1-C_3 alkyl, $-F$, $-Cl$, $-Br$, $-I$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, C_1-C_3 alkoxy, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,
- (2) C_2-C_6 alkenyl with one or two double bonds,
- 10 optionally substituted with one, two or three substituents selected from the group consisting of $-F$, $-Cl$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, C_1-C_3 alkoxy, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are $-H$ or C_1-C_6 alkyl,
- (3) C_2-C_6 alkynyl with one or two triple bonds,
- 15 optionally substituted with one, two or three substituents selected from the group consisting of $-F$, $-Cl$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, C_1-C_3 alkoxy, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are $-H$ or C_1-C_6 alkyl,
- (4) $-F$, $-Cl$, $-Br$ or $-I$,
- (6) $-C_1-C_6$ alkoxy optionally substituted with one, two, or three of $-F$,
- 20 (7) $-NR_{N-2}R_{N-3}$ where R_{N-2} and R_{N-3} are as defined below,
- (8) $-OH$,
- (9) $-C\equiv N$,
- (10) C_3-C_7 cycloalkyl, optionally substituted with one,
- 25 two or three substituents selected from the group consisting of $-F$, $-Cl$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, C_1-C_3 alkoxy, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are $-H$ or C_1-C_6 alkyl,
- (11) $-CO-(C_1-C_4 \text{ alkyl})$,
- (12) $-SO_2-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,
- 30 (13) $-CO-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined above, or
- (14) $-SO_2-(C_1-C_4 \text{ alkyl})$, with the proviso that when n_1 is zero $R_{1\text{-heteroaryl}}$ is not bonded to the carbon chain by nitrogen; or

(VIII) $-(CH_2)_{n_1}-(R_{1-heterocycle})$ where n_1 is as defined above and $R_{1-heterocycle}$

$R_{1-heterocycle}$ is selected from the group consisting of:

- morpholinyl,
- thiomorpholinyl,
- 5 thiomorpholinyl S-oxide,
- thiomorpholinyl S,S-dioxide,
- piperazinyl,
- homopiperazinyl,
- pyrrolidinyl,
- 10 pyrrolinyl,
- tetrahydropyranyl,
- piperidinyl,
- tetrahydrofuranyl,
- tetrahydrothienyl,
- 15 homopiperidinyl,
- homomorpholinyl,
- homothiomorpholinyl,
- homothiomorpholinyl S,S-dioxide,
- oxazolidinonyl,
- 20 dihydropyrazolyl,
- dihydropyrrolyl,
- dihydropyrazinyl,
- dihydropyridinyl,
- dihydropyrimidinyl,
- 25 dihydrofuryl,
- dihydropyranyl,
- tetrahydrothienyl S-oxide,
- tetrahydrothienyl S,S-dioxide, and
- homothiomorpholinyl S-oxide,

30 where the $R_{1-heterocycle}$ group is bonded by any atom of the parent $R_{1-heterocycle}$

$R_{1-heterocycle}$ group substituted by hydrogen such that the new bond to the $R_{1-heterocycle}$ group replaces the hydrogen atom and its bond, where heterocycle is optionally substituted with one, two, three or four:

- (1) C₁-C₆ alkyl optionally substituted with one, two or three substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₃ alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,
- 5 (2) C₂-C₆ alkenyl with one or two double bonds, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, -C≡N, -CF₃, C₁-C₃ alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are -H or C₁-C₆ alkyl,
- 10 (3) C₂-C₆ alkynyl with one or two triple bonds, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, -C≡N, -CF₃, C₁-C₃ alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are -H or C₁-C₆ alkyl,
- (4) -F, Cl, -Br or -I,
- (5) C₁-C₆ alkoxy,
- 15 (6) -C₁-C₆ alkoxy optionally substituted with one, two, or three -F,
- (7) -NR_{N-2}R_{N-3} where R_{N-2} and R_{N-3} are as defined below,
- (8) -OH,
- 20 (9) -C≡N,
- (10) C₃-C₇ cycloalkyl, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, -C≡N, -CF₃, C₁-C₃ alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are -H or C₁-C₆ alkyl,
- (11) -CO-(C₁-C₄ alkyl),
- 25 (12) -SO₂-NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,
- (13) -CO-NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,
- (14) -SO₂-(C₁-C₄ alkyl), or
- 30 (15) =O, with the proviso that when n₁ is zero R_{1-heterocycle} is not bonded to the carbon chain by nitrogen;

where R₂ is:

(I)-H,

(II) C₁-C₆ alkyl, optionally substituted with one, two or three substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₃ alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(III) -(CH₂)₀₋₄-R₂₋₁ where R₂₋₁ is R_{1-aryl} or R_{1-heteroaryl} where R_{1-aryl} and R_{1-heteroaryl} are as defined above;

(IV) C₂-C₆ alkenyl with one or two double bonds, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, -C≡N, -CF₃, C₁-C₃ alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are -H or C₁-C₆ alkyl,

(V) C₂-C₆ alkynyl with one or two triple bonds, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, -C≡N, -CF₃, C₁-C₃ alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are -H or C₁-C₆ alkyl, or

(VI) -(CH₂)₀₋₄-C₃-C₇ cycloalkyl, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, -C≡N, -CF₃, C₁-C₃ alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are -H or C₁-C₆ alkyl;

where R₃ is:

(I)-H,

(II) C₁-C₆ alkyl, optionally substituted with one, two or three substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₃ alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(III) -(CH₂)₀₋₄-R₂₋₁ where R₂₋₁ is R_{1-aryl} or R_{1-heteroaryl} where R_{1-aryl} and R_{1-heteroaryl} are as defined above

(IV) C₂-C₆ alkenyl with one or two double bonds,

(V) C₂-C₆ alkynyl with one or two triple bonds; or

(VI) -(CH₂)₀₋₄-C₃-C₇ cycloalkyl, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, -C≡N, -CF₃, C₁-C₃ alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are -H or C₁-C₆ alkyl,

and where R_2 and R_3 are taken together with the carbon to which they are attached to form a carbocycle of three, four, five, six, or seven carbon atoms, optionally where one carbon atom is replaced by a heteroatom selected from the group consisting of $-O-$, $-S-$, $-SO_2-$, and $-NR_{N-2}-$, where R_{N-2} is selected from the group consisting of:

- (a) $-H$,
- (b) $-C_1-C_6$ alkyl optionally substituted with one substituent selected from the group consisting of:
 - (i) $-OH$, and
 - (ii) $-NH_2$,
- (c) $-C_1-C_6$ alkyl optionally substituted with one, two, or three $-F$, $-Cl$, $-Br$, or $-I$,
- (d) $-C_3-C_7$ cycloalkyl,
- (e) $-(C_1-C_2 \text{ alkyl})-(C_3-C_7 \text{ cycloalkyl})$,
- (f) $-(C_1-C_6 \text{ alkyl})-O-(C_1-C_3 \text{ alkyl})$,
- (g) $-C_2-C_6$ alkenyl with one or two double bonds,
- (h) $-C_2-C_6$ alkynyl with one or two triple bonds,
- (i) $-C_1-C_6$ alkyl chain with one double bond and one triple bond,
- (j) $-R_{1-aryl}$ where R_{1-aryl} is as defined above, and
- (k) $-R_{1-heteroaryl}$ where $R_{1-heteroaryl}$ is as defined above;

25

where R_N is:

- (I) $R_{N-1}-X_N-$ where X_N is selected from the group consisting of:
 - (A) $-CO-$,
 - (B) $-SO_2-$,
 - (C) $-(CR'R'')_{1-6}$ where R' and R'' are the same or different and are $-H$ and C_1-C_4 alkyl,
 - (D) $-CO-(CR'R'')_{1-6}-X_{N-1}$ where X_{N-1} is selected from the group consisting of $-O-$, $-S-$ and $-NR'-$ and where R' and R'' are as defined above, and

30

(E) a single bond;

where R_{N-1} is selected from the group consisting of:

(A) R_{N-aryl} where R_{N-aryl} is phenyl, 1-naphthyl, 2-naphthyl, tetralinyl, indanyl, 6,7,8,9-tetrahydro-5H-benzo[a]cycloheptenyl, or dihydronaphthyl
 5 optionally substituted with one, two or three of the following substituents which can be the same or different and are:

(1) C_1 - C_6 alkyl, optionally substituted with one, two or three substituents selected from the group consisting of C_1 - C_3 alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C \equiv N, -CF₃, C_1 - C_3 alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as
 10 defined above,

(2) -OH,

(3) -NO₂,

(4) -F, -Cl, -Br, -I,

(5) -CO-OH,

15 (6) -C \equiv N,

(7) -(CH₂)₀₋₄-CO-NR_{N-2}R_{N-3} where R_{N-2} and R_{N-3} are the same or different and are selected from the group consisting of:

(a) -H,

(b) - C_1 - C_6 alkyl optionally substituted with one

20 substituent selected from the group consisting of:

(i) -OH, and

(ii) -NH₂,

(c) - C_1 - C_6 alkyl optionally substituted with

one, two, or three -F, -Cl, -Br, or -I,

25 (d) - C_3 - C_7 cycloalkyl,

(e) -(C_1 - C_2 alkyl)-(C_3 - C_7 cycloalkyl),

(f) -(C_1 - C_6 alkyl)-O-(C_1 - C_3 alkyl),

(g) - C_2 - C_6 alkenyl with one or two double

bonds,

30 (h) - C_2 - C_6 alkynyl with one or two triple

bonds,

(i) - C_1 - C_6 alkyl chain with one double bond

and one triple bond,

(j) $-R_{1-aryl}$ where R_{1-aryl} is as defined above, and

(k) $-R_{1-heteroaryl}$ where $R_{1-heteroaryl}$ is as defined

above,

(8) $-(CH_2)_{0-4}-CO-(C_1-C_{12} \text{ alkyl})$,

5 (9) $-(CH_2)_{0-4}-CO-(C_2-C_{12} \text{ alkenyl with one, two or three double bonds})$,

(10) $-(CH_2)_{0-4}-CO-(C_2-C_{12} \text{ alkynyl with one, two or three triple bonds})$,

(11) $-(CH_2)_{0-4}-CO-(C_3-C_7 \text{ cycloalkyl})$,

10 (12) $-(CH_2)_{0-4}-CO-R_{1-aryl}$ where R_{1-aryl} is as defined above,

(13) $-(CH_2)_{0-4}-CO-R_{1-heteroaryl}$ where $R_{1-heteroaryl}$ is as defined above,

15 (14) $-(CH_2)_{0-4}-CO-R_{1-heterocycle}$ where $R_{1-heterocycle}$ is as defined above,

(15) $-(CH_2)_{0-4}-CO-R_{N-4}$ where R_{N-4} is selected from the group consisting of morpholinyl, thiomorpholinyl, piperazinyl, piperidinyl, homomorpholinyl, homothiomorpholinyl, homothiomorpholinyl S-oxide, homothiomorpholinyl S,S-dioxide, pyrrolinyl and pyrrolidinyl where each group is

20 optionally substituted with one, two, three, or four of C_1-C_6 alkyl,

(16) $-(CH_2)_{0-4}-CO-O-R_{N-5}$ where R_{N-5} is selected from the group consisting of:

(a) C_1-C_6 alkyl,

25 (b) $-(CH_2)_{0-2}-(R_{1-aryl})$ where R_{1-aryl} is as defined above,

(c) C_2-C_6 alkenyl containing one or two double bonds,

(d) C_2-C_6 alkynyl containing one or two triple bonds,

30 (e) C_3-C_7 cycloalkyl, and

(f) $-(CH_2)_{0-2}-(R_{1-heteroaryl})$ where $R_{1-heteroaryl}$ is as defined above,

(17) $-(CH_2)_{0-4}-SO_2-NR_{N-2}R_{N-3}$ where R_{N-2} and R_{N-3} are as defined above,

- (18) $-(CH_2)_{0-4}-SO-(C_1-C_8 \text{ alkyl})$,
- (19) $-(CH_2)_{0-4}-SO_2-(C_1-C_{12} \text{ alkyl})$,
- (20) $-(CH_2)_{0-4}-SO_2-(C_3-C_7 \text{ cycloalkyl})$,
- (21) $-(CH_2)_{0-4}-N(H \text{ or } R_{N-5})-CO-O-R_{N-5}$ where R_{N-5}
 5 can be the same or different and is as defined above,
- (22) $-(CH_2)_{0-4}-N(H \text{ or } R_{N-5})-CO-N(R_{N-5})_2$, where R_{N-5}
 can be the same or different and is as defined above,
- (23) $-(CH_2)_{0-4}-N-CS-N(R_{N-5})_2$, where R_{N-5} can be the
 same or different and is as defined above,
- (24) $-(CH_2)_{0-4}-N(-H \text{ or } R_{N-5})-CO-R_{N-2}$ where R_{N-5} and
 10 R_{N-2} can be the same or different and are as defined above,
- (25) $-(CH_2)_{0-4}-NR_{N-2}R_{N-3}$ where R_{N-2} and R_{N-3} can be
 the same or different and are as defined above,
- (26) $-(CH_2)_{0-4}-R_{N-4}$ where R_{N-4} is as defined above,
- (27) $-(CH_2)_{0-4}-O-CO-(C_1-C_6 \text{ alkyl})$,
- (28) $-(CH_2)_{0-4}-O-P(O)-(OR_{N-aryl-1})_2$ where $R_{N-aryl-1}$ is –
 15 H or C_1-C_4 alkyl,
- (29) $-(CH_2)_{0-4}-O-CO-N(R_{N-5})_2$ where R_{N-5} is as
 defined above,
- (30) $-(CH_2)_{0-4}-O-CS-N(R_{N-5})_2$ where R_{N-5} is as defined
 20 above,
- (31) $-(CH_2)_{0-4}-O-(R_{N-5})_2$ where R_{N-5} is as defined
 above,
- (32) $-(CH_2)_{0-4}-O-(R_{N-5})_2-COOH$ where R_{N-5} is as
 25 defined above,
- (33) $-(CH_2)_{0-4}-S-(R_{N-5})_2$ where R_{N-5} is as defined
 above,
- (34) $-(CH_2)_{0-4}-O-(C_1-C_6 \text{ alkyl optionally substituted}$
 with one, two, three, four, or five of –F),
- (35) $C_3-C_7 \text{ cycloalkyl}$,
- (36) $C_2-C_6 \text{ alkenyl with one or two double bonds}$
 30 optionally substituted with C_1-C_3 alkyl, –F, –Cl, –Br, –I, –OH, –SH, –C≡N, –CF₃, C_1-C_3 alkoxy, or –NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(37) C₂-C₆ alkynyl with one or two triple bonds

optionally substituted with C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₃ alkoxy, or -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(38) -(CH₂)₀₋₄-N(-H or R_{N-5})-SO₂-R_{N-2} where R_{N-5} and
5 R_{N-2} can be the same or different and are as described above, or

(39) -(CH₂)₀₋₄-C₃-C₇ cycloalkyl,

(B) -R_{N-heteroaryl} where R_{N-heteroaryl} is selected from the group
consisting of:

10 pyridinyl,
pyrimidinyl,
quinolinyl,
benzothienyl,
indolyl,
indolinyl,
15 pyridazinyl,
pyrazinyl,
isoindolyl,
isoquinolyl,
quinazolinyl,
20 quinoxalinyl,
phthalazinyl,
imidazolyl,
isoxazolyl,
pyrazolyl,
25 oxazolyl,
thiazolyl,
indolizinyl,
indazolyl,
benzothiazolyl,
30 benzimidazolyl,
benzofuranyl,
furanyl,
thienyl,

	pyrrolyl,
	oxadiazolyl,
	thiadiazolyl,
	triazolyl,
5	tetrazolyl,
	oxazolopyridinyl,
	imidazopyridinyl,
	isothiazolyl,
	naphthyridinyl,
10	cinnolyl,
	carbazolyl,
	beta-carbolinyl,
	isochromanyl,
	chromanyl,
15	tetrahydroisoquinolyl,
	isoindolyl,
	isobenzotetrahydrofuranyl,
	isobenzotetrahydrothienyl,
	isobenzothienyl,
20	benzoxazolyl,
	pyridopyridinyl,
	benzotetrahydrofuranyl,
	benzotetrahydrothienyl,
	purinyl,
25	benzodioxolyl,
	triazinyl,
	phenoxazinyl,
	phenothiazinyl,
	pteridinyl,
30	benzothiazolyl,
	imidazopyridinyl,
	imidazothiazolyl,
	dihydrobenzisoxazinyl,
	benzisoxazinyl,

	benzoxazinyl,
	dihydrobenzisothiazinyl,
	benzopyranyl,
	benzothiopyranyl,
5	coumarinyl,
	isocoumarinyl,
	chromonyl,
	chromanonyl,
	pyridinyl-N-oxide,
10	tetrahydroquinolinyl,
	dihydroquinolinyl,
	dihydroquinolinonyl,
	dihydroisoquinolinonyl,
	dihydrocoumarinyl,
15	dihydroisocoumarinyl,
	isoindolinonyl,
	benzodioxanyl,
	benzoxazolinonyl,
	pyrrolyl N-oxide,
20	pyrimidinyl N-oxide,
	pyridazinyl N-oxide,
	pyrazinyl N-oxide,
	quinolinyl N-oxide,
	indolyl N-oxide,
25	indolinyl N-oxide,
	isoquinolyl N-oxide,
	quinazolinyl N-oxide,
	quinoxalinyl N-oxide,
	phthalazinyl N-oxide,
30	imidazolyl N-oxide,
	isoxazolyl N-oxide,
	oxazolyl N-oxide,
	thiazolyl N-oxide,
	indolizinyl N-oxide,

indazolyl N-oxide,
 benzothiazolyl N-oxide,
 benzimidazolyl N-oxide,
 pyrrolyl N-oxide,
 5 oxadiazolyl N-oxide,
 thiadiazolyl N-oxide,
 triazolyl N-oxide,
 tetrazolyl N-oxide,
 benzothiopyranyl S-oxide, and
 10 benzothiopyranyl S,S-dioxide

where the $R_{N\text{-heteroaryl}}$ group is bonded by any atom of the parent $R_{N\text{-heteroaryl}}$ group substituted by hydrogen such that the new bond to the $R_{N\text{-heteroaryl}}$ group replaces the hydrogen atom and its bond, where heteroaryl is optionally substituted with one, two, three, or four of:

15 (1) $C_1\text{-}C_6$ alkyl, optionally substituted with one, two or three substituents selected from the group consisting of $C_1\text{-}C_3$ alkyl, -F, -Cl, -Br, -I, -OH, -SH, $-C\equiv N$, $-CF_3$, $C_1\text{-}C_3$ alkoxy, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

(2) -OH,
 20 (3) $-NO_2$,
 (4) -F, -Cl, -Br, or -I,
 (5) -CO-OH,
 (6) $-C\equiv N$,
 (7) $-(CH_2)_{0-4}\text{-CO-NR}_{N-2}R_{N-3}$ where R_{N-2} and R_{N-3} are

25 the same or different and are selected from the group consisting of:

(a) -H,
 (b) $-C_1\text{-}C_6$ alkyl optionally substituted with one substituent selected from the group consisting of:
 (i) -OH, and
 30 (ii) $-NH_2$,
 (c) $-C_1\text{-}C_6$ alkyl optionally substituted with one, two, or three -F, -Cl, -Br, -I,
 (d) $-C_3\text{-}C_7$ cycloalkyl,

293

- (e) $-(C_1-C_2 \text{ alkyl})-(C_3-C_7 \text{ cycloalkyl})$,
 (f) $-(C_1-C_6 \text{ alkyl})-O-(C_1-C_3 \text{ alkyl})$,
 (g) $-C_2-C_6 \text{ alkenyl}$ with one or two double bonds,
 5 (h) $-C_2-C_6 \text{ alkynyl}$ with one or two triple bonds,
 and one triple bond,
 (j) $-R_{1\text{-aryl}}$ where $R_{1\text{-aryl}}$ is as defined above,
 10 (k) $-R_{1\text{-heteroaryl}}$ where $R_{1\text{-heteroaryl}}$ is as defined above,
 (8) $-(CH_2)_{0-4}-CO-(C_1-C_{12} \text{ alkyl})$,
 (9) $-(CH_2)_{0-4}-CO-(C_2-C_{12} \text{ alkenyl}$ with one, two or three double bonds),
 15 (10) $-(CH_2)_{0-4}-CO-(C_2-C_{12} \text{ alkynyl}$ with one, two or three triple bonds),
 (11) $-(CH_2)_{0-4}-CO-(C_3-C_7 \text{ cycloalkyl})$,
 (12) $-(CH_2)_{0-4}-CO-R_{1\text{-aryl}}$ where $R_{1\text{-aryl}}$ is as defined above,
 20 (13) $-(CH_2)_{0-4}-CO-R_{1\text{-heteroaryl}}$ where $R_{1\text{-heteroaryl}}$ is as defined above,
 (14) $-(CH_2)_{0-4}-CO-R_{1\text{-heterocycle}}$ where $R_{1\text{-heterocycle}}$ is as defined above,
 (15) $-(CH_2)_{0-4}-CO-R_{N-4}$ where R_{N-4} is selected from
 25 the group consisting of morpholinyl, thiomorpholinyl, piperazinyl, piperidinyl, homomorpholinyl, homothiomorpholinyl, homothiomorpholinyl S-oxide, homothiomorpholinyl S,S-dioxide, pyrrolinyl and pyrrolidinyl where each group is optionally substituted with one, two, three, or four of $C_1-C_6 \text{ alkyl}$,
 (16) $-(CH_2)_{0-4}-CO-O-R_{N-5}$ where R_{N-5} is selected from
 30 the group consisting of:
 (a) $C_1-C_6 \text{ alkyl}$,
 (b) $-(CH_2)_{0-2}-(R_{1\text{-aryl}})$ where $R_{1\text{-aryl}}$ is as defined above,

- bonds,
- (c) C₂-C₆ alkenyl containing one or two double bonds,
- (d) C₂-C₆ alkynyl containing one or two triple bonds,
- 5 (e) C₃-C₇ cycloalkyl, and
- (f) -(CH₂)₀₋₂-(R_{1-heteroaryl}) where R_{1-heteroaryl} is as defined above,
- (17) -(CH₂)₀₋₄-SO₂-NR_{N-2}R_{N-3} where R_{N-2} and R_{N-3} are as defined above,
- 10 (18) -(CH₂)₀₋₄-SO-(C₁-C₈ alkyl),
- (19) -(CH₂)₀₋₄-SO₂-(C₁-C₁₂ alkyl),
- (20) -(CH₂)₀₋₄-SO₂-(C₃-C₇ cycloalkyl),
- (21) -(CH₂)₀₋₄-N(H or R_{N-5})-CO-O-R_{N-5} where R_{N-5} can be the same or different and is as defined above,
- 15 (22) -(CH₂)₀₋₄-N(H or R_{N-5})-CO-N(R_{N-5})₂, where R_{N-5} can be the same or different and is as defined above,
- (23) -(CH₂)₀₋₄-N-CS-N(R_{N-5})₂, where R_{N-5} can be the same or different and is as defined above,
- (24) -(CH₂)₀₋₄-N(-H or R_{N-5})-CO-R_{N-2} where R_{N-5} and R_{N-2} can be the same or different and are as defined above,
- 20 (25) -(CH₂)₀₋₄-NR_{N-2}R_{N-3} where R_{N-2} and R_{N-3} can be the same or different and are as defined above,
- (26) -(CH₂)₀₋₄-R_{N-4} where R_{N-4} is as defined above,
- (27) -(CH₂)₀₋₄-O-CO-(C₁-C₆ alkyl),
- 25 (28) -(CH₂)₀₋₄-O-P(O)-(OR_{N-aryl-1})₂ where R_{N-aryl-1} is -H or C₁-C₄ alkyl,
- (29) -(CH₂)₀₋₄-O-CO-N(R_{N-5})₂ where R_{N-5} is as defined above,
- (30) -(CH₂)₀₋₄-O-CS-N(R_{N-5})₂ where R_{N-5} is as defined above,
- 30 (31) -(CH₂)₀₋₄-O-(R_{N-5})₂ where R_{N-5} is as defined above,
- (32) -(CH₂)₀₋₄-O-(R_{N-5})₂-COOH where R_{N-5} is as defined above,

(33) $-(\text{CH}_2)_{0-4}-\text{S}-(\text{R}_{\text{N}-5})_2$ where $\text{R}_{\text{N}-5}$ is as defined

above,

(34) $-(\text{CH}_2)_{0-4}-\text{O}-(\text{C}_1-\text{C}_6 \text{ alkyl optionally substituted with one, two, three, four, or five of } -\text{F}),$

5

(35) $\text{C}_3-\text{C}_7 \text{ cycloalkyl},$

(36) $\text{C}_2-\text{C}_6 \text{ alkenyl with one or two double bonds optionally substituted with } \text{C}_1-\text{C}_3 \text{ alkyl, } -\text{F, } -\text{Cl, } -\text{Br, } -\text{I, } -\text{OH, } -\text{SH, } -\text{C}\equiv\text{N, } -\text{CF}_3, \text{ C}_1-\text{C}_3 \text{ alkoxy, or } -\text{NR}_{1-\text{a}}\text{R}_{1-\text{b}} \text{ where } \text{R}_{1-\text{a}} \text{ and } \text{R}_{1-\text{b}} \text{ are as defined above,}$

10

(37) $\text{C}_2-\text{C}_6 \text{ alkynyl with one or two triple bonds optionally substituted with } \text{C}_1-\text{C}_3 \text{ alkyl, } -\text{F, } -\text{Cl, } -\text{Br, } -\text{I, } -\text{OH, } -\text{SH, } -\text{C}\equiv\text{N, } -\text{CF}_3, \text{ C}_1-\text{C}_3 \text{ alkoxy, or } -\text{NR}_{1-\text{a}}\text{R}_{1-\text{b}} \text{ where } \text{R}_{1-\text{a}} \text{ and } \text{R}_{1-\text{b}} \text{ are as defined above,}$

(38) $-(\text{CH}_2)_{0-4}-\text{N}(-\text{H or } \text{R}_{\text{N}-5})-\text{SO}_2-\text{R}_{\text{N}-2}$ where $\text{R}_{\text{N}-5}$ and $\text{R}_{\text{N}-2}$ can be the same or different and are as defined above, or

(39) $-(\text{CH}_2)_{0-4}-\text{C}_3-\text{C}_7 \text{ cycloalkyl},$

15

(C) $\text{R}_{\text{N-aryl}}-\text{W}-\text{R}_{\text{N-aryl}},$ where $\text{R}_{\text{N-aryl}}$ can be the same or different,

(D) $\text{R}_{\text{N-aryl}}-\text{W}-\text{R}_{\text{N-heteroaryl}},$

(E) $\text{R}_{\text{N-aryl}}-\text{W}-\text{R}_{\text{N-1-heterocycle}},$ wherein $\text{R}_{\text{N-1-heterocycle}}$ is the same as $\text{R}_{1-\text{heterocycle}},$ and $\text{R}_{1-\text{heterocycle}}$ is as defined above

20

(F) $\text{R}_{\text{N-heteroaryl}}-\text{W}-\text{R}_{\text{N-aryl}},$

(G) $\text{R}_{\text{N-heteroaryl}}-\text{W}-\text{R}_{\text{N-heteroaryl}},$

(H) $\text{R}_{\text{N-heteroaryl}}-\text{W}-\text{R}_{\text{N-1-heterocycle}},$

(I) $\text{R}_{\text{N-heterocycle}}-\text{W}-\text{R}_{\text{N-aryl}},$ wherein $\text{R}_{\text{N-heterocycle}}$ is the same as $\text{R}_{1-\text{heterocycle}},$ and $\text{R}_{1-\text{heterocycle}}$ is as defined above, and $\text{R}_{\text{N-aryl}}$ is as defined above,

25

(J) $\text{R}_{\text{N-heterocycle}}-\text{W}-\text{R}_{\text{N-heteroaryl}},$ and

(K) $\text{R}_{\text{N-heterocycle}}-\text{W}-\text{R}_{\text{N-1-heterocycle}},$

where W is

(17) $-(\text{CH}_2)_{0-4}-,$

(18) $-\text{O}-,$

30

(19) $-\text{S}(\text{O})_{0-2}-,$

(20) $-\text{N}(\text{R}_{\text{N}-5})-$ where $\text{R}_{\text{N}-5}$ is as defined

above, or

(5) $-\text{CO}-;$

(II) $-\text{CO}-(\text{C}_1\text{-C}_{10}\text{ alkyl})$ where alkyl is optionally substituted with one three substituents selected from the group consisting of:

- (A) $-\text{OH}$,
(B) $-\text{C}_1\text{-C}_6\text{ alkoxy}$,
5 (C) $-\text{C}_1\text{-C}_6\text{ thioalkoxy}$,
(D) $-\text{CO-O-R}_{\text{N-8}}$ where $\text{R}_{\text{N-8}}$ is $-\text{H}$, $\text{C}_1\text{-C}_6\text{ alkyl}$ or $-\text{phenyl}$,
(E) $-\text{CO-NR}_{\text{N-2}}\text{R}_{\text{N-3}}$ where $\text{R}_{\text{N-2}}$ and $\text{R}_{\text{N-3}}$ are the same or different and are as defined above,
(F) $-\text{CO-R}_{\text{N-4}}$ where $\text{R}_{\text{N-4}}$ is as defined above,
10 (G) $-\text{SO}_2-(\text{C}_1\text{-C}_8\text{ alkyl})$,
(H) $-\text{SO}_2\text{-NR}_{\text{N-2}}\text{R}_{\text{N-3}}$ where $\text{R}_{\text{N-2}}$ and $\text{R}_{\text{N-3}}$ are the same or different and are as defined above,
(I) $-\text{NH-CO}-(\text{C}_1\text{-C}_6\text{ alkyl})$,
(J) $-\text{NH-CO-O-R}_{\text{N-8}}$ where $\text{R}_{\text{N-8}}$ is as defined above,
15 (K) $-\text{NR}_{\text{N-2}}\text{R}_{\text{N-3}}$ where $\text{R}_{\text{N-2}}$ and $\text{R}_{\text{N-3}}$ are the same or different and are as defined above,
(L) $-\text{R}_{\text{N-4}}$ where $\text{R}_{\text{N-4}}$ is as defined above,
(M) $-\text{O-CO}-(\text{C}_1\text{-C}_6\text{ alkyl})$,
(N) $-\text{O-CO-NR}_{\text{N-8}}\text{R}_{\text{N-8}}$ where $\text{R}_{\text{N-8}}$ are the same or different
20 and are as defined above,
(O) $-\text{O}-(\text{C}_1\text{-C}_5\text{ alkyl})\text{-COOH}$,
(P) $-\text{O}-(\text{C}_1\text{-C}_6\text{ alkyl})$ optionally substituted with one, two, or three of $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, or $-\text{I}$,
(Q) $-\text{NH-SO}_2-(\text{C}_1\text{-C}_6\text{ alkyl})$, and
25 (R) $-\text{F}$, or $-\text{Cl}$,

(III) $-\text{CO}-(\text{C}_1\text{-C}_6\text{ alkyl})\text{-O}-(\text{C}_1\text{-C}_6\text{ alkyl})$ where alkyl is optionally substituted with one, two, or three substituents selected from the group consisting of:

- (A) $-\text{OH}$,
30 (B) $-\text{C}_1\text{-C}_6\text{ alkoxy}$,
(C) $-\text{C}_1\text{-C}_6\text{ thioalkoxy}$,
(D) $-\text{CO-O-R}_{\text{N-8}}$ where $\text{R}_{\text{N-8}}$ is $-\text{H}$, $\text{C}_1\text{-C}_6\text{ alkyl}$ or $-\text{phenyl}$,

(E) $-\text{CO}-\text{NR}_{\text{N}-2}\text{R}_{\text{N}-3}$ where $\text{R}_{\text{N}-2}$ and $\text{R}_{\text{N}-3}$ are the same or different and are as defined above,

(F) $-\text{CO}-\text{R}_{\text{N}-4}$ where $\text{R}_{\text{N}-4}$ is as defined above,

(G) $-\text{SO}_2-(\text{C}_1-\text{C}_8 \text{ alkyl})$,

5 (H) $-\text{SO}_2-\text{NR}_{\text{N}-2}\text{R}_{\text{N}-3}$ where $\text{R}_{\text{N}-2}$ and $\text{R}_{\text{N}-3}$ are the same or different and are as defined above,

(I) $-\text{NH}-\text{CO}-(\text{C}_1-\text{C}_6 \text{ alkyl})$,

(J) $-\text{NH}-\text{CO}-\text{O}-\text{R}_{\text{N}-8}$ where $\text{R}_{\text{N}-8}$ is as defined above,

(K) $-\text{NR}_{\text{N}-2}\text{R}_{\text{N}-3}$ where $\text{R}_{\text{N}-2}$ and $\text{R}_{\text{N}-3}$ are the same or different

10 and are as defined above,

(L) $-\text{R}_{\text{N}-4}$ where $\text{R}_{\text{N}-4}$ is as defined above,

(M) $-\text{O}-\text{CO}-(\text{C}_1-\text{C}_6 \text{ alkyl})$,

(N) $-\text{O}-\text{CO}-\text{NR}_{\text{N}-8}\text{R}_{\text{N}-8}$ where $\text{R}_{\text{N}-8}$ are the same or different

and are as defined above,

15 (O) $-\text{O}-(\text{C}_1-\text{C}_5 \text{ alkyl})-\text{COOH}$,

(P) $-\text{O}-(\text{C}_1-\text{C}_6 \text{ alkyl})$ optionally substituted with one, two, or three of -F, -Cl, -Br, or -I),

(Q) $-\text{NH}-\text{SO}_2-(\text{C}_1-\text{C}_6 \text{ alkyl})$, and

(R) -F, or -Cl,

20 (IV) $-\text{CO}-(\text{C}_1-\text{C}_6 \text{ alkyl})-\text{S}-(\text{C}_1-\text{C}_6 \text{ alkyl})$ where alkyl is optionally substituted with one, two, or three of substituents selected from the group consisting of:

(A) -OH,

(B) $-\text{C}_1-\text{C}_6 \text{ alkoxy}$,

25 (C) $-\text{C}_1-\text{C}_6 \text{ thioalkoxy}$,

(D) $-\text{CO}-\text{O}-\text{R}_{\text{N}-8}$ where $\text{R}_{\text{N}-8}$ is as defined above,

(E) $-\text{CO}-\text{NR}_{\text{N}-2}\text{R}_{\text{N}-3}$ where $\text{R}_{\text{N}-2}$ and $\text{R}_{\text{N}-3}$ are the same or different and are as defined above,

(F) $-\text{CO}-\text{R}_{\text{N}-4}$ where $\text{R}_{\text{N}-4}$ is as defined above,

30 (G) $-\text{SO}_2-(\text{C}_1-\text{C}_8 \text{ alkyl})$,

(H) $-\text{SO}_2-\text{NR}_{\text{N}-2}\text{R}_{\text{N}-3}$ where $\text{R}_{\text{N}-2}$ and $\text{R}_{\text{N}-3}$ are the same or different and are as defined above,

(I) $-\text{NH}-\text{CO}-(\text{C}_1-\text{C}_6 \text{ alkyl})$,

(J) $-\text{NH}-\text{CO}-\text{O}-\text{R}_{\text{N}-8}$ where $\text{R}_{\text{N}-8}$ is as defined above,

(K) $-\text{NR}_{\text{N}-2}\text{R}_{\text{N}-3}$ where $\text{R}_{\text{N}-2}$ and $\text{R}_{\text{N}-3}$ are the same or different

and are as defined above,

(L) $-\text{R}_{\text{N}-4}$ where $\text{R}_{\text{N}-4}$ is as defined above,

5 (M) $-\text{O}-\text{CO}-(\text{C}_1-\text{C}_6 \text{ alkyl})$,

(N) $-\text{O}-\text{CO}-\text{NR}_{\text{N}-8}\text{R}_{\text{N}-8}$ where $\text{R}_{\text{N}-8}$ are the same or different

and are as defined above,

(O) $-\text{O}-(\text{C}_1-\text{C}_5 \text{ alkyl})-\text{COOH}$,

(P) $-\text{O}-(\text{C}_1-\text{C}_6 \text{ alkyl})$ optionally substituted with one, two, or

10 three of $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, or $-\text{I}$,

(Q) $-\text{NH}-\text{SO}_2-(\text{C}_1-\text{C}_6 \text{ alkyl})$, and

(R) $-\text{F}$, or $-\text{Cl}$,

(V) $-\text{CO}-\text{CH}(-(\text{CH}_2)_{0-2}-\text{O}-\text{R}_{\text{N}-10})-(\text{CH}_2)_{0-2}-\text{R}_{\text{N-aryl}}/\text{R}_{\text{N-heteroaryl}}$ where

$\text{R}_{\text{N-aryl}}$ and $\text{R}_{\text{N-heteroaryl}}$ are as defined above, where $\text{R}_{\text{N}-10}$ is selected from the group

15 consisting of:

(A) $-\text{H}$,

(B) $\text{C}_1-\text{C}_6 \text{ alkyl}$,

(C) $\text{C}_3-\text{C}_7 \text{ cycloalkyl}$,

(D) $\text{C}_2-\text{C}_6 \text{ alkenyl}$ with one double bond,

20 (E) $\text{C}_2-\text{C}_6 \text{ alkynyl}$ with one triple bond,

(F) $\text{R}_{1-\text{aryl}}$ where $\text{R}_{1-\text{aryl}}$ is as defined above, and

(G) $\text{R}_{\text{N-heteroaryl}}$ where $\text{R}_{\text{N-heteroaryl}}$ is as defined above, or

(VI) $-\text{CO}-(\text{C}_3-\text{C}_8 \text{ cycloalkyl})$ where alkyl is optionally substituted

with one or two substituents selected from the group consisting of:

25 (A) $-(\text{CH}_2)_{0-4}-\text{OH}$,

(B) $-(\text{CH}_2)_{0-4}-\text{C}_1-\text{C}_6 \text{ alkoxy}$,

(C) $-(\text{CH}_2)_{0-4}-\text{C}_1-\text{C}_6 \text{ thioalkoxy}$,

(D) $-(\text{CH}_2)_{0-4}-\text{CO}-\text{O}-\text{R}_{\text{N}-8}$ where $\text{R}_{\text{N}-8}$ is $-\text{H}$, $\text{C}_1-\text{C}_6 \text{ alkyl}$ or

phenyl,

30 (E) $-(\text{CH}_2)_{0-4}-\text{CO}-\text{NR}_{\text{N}-2}\text{R}_{\text{N}-3}$ where $\text{R}_{\text{N}-2}$ and $\text{R}_{\text{N}-3}$ are the

same or different and are as defined above,

(F) $-(\text{CH}_2)_{0-4}-\text{CO}-\text{R}_{\text{N}-4}$ where $\text{R}_{\text{N}-4}$ is as defined above,

(G) $-(\text{CH}_2)_{0-4}-\text{SO}_2-(\text{C}_1-\text{C}_8 \text{ alkyl})$,

(H) $-(\text{CH}_2)_{0-4}-\text{SO}_2-\text{NR}_{\text{N}-2}\text{R}_{\text{N}-3}$ where $\text{R}_{\text{N}-2}$ and $\text{R}_{\text{N}-3}$ are the same or different and are as defined above,

(I) $-(\text{CH}_2)_{0-4}-\text{NH}-\text{CO}-(\text{C}_1-\text{C}_6 \text{ alkyl})$,

(J) $-\text{NH}-\text{CO}-\text{O}-\text{R}_{\text{N}-8}$ where $\text{R}_{\text{N}-8}$ is as defined above,

5 (K) $-(\text{CH}_2)_{0-4}-\text{NR}_{\text{N}-2}\text{R}_{\text{N}-3}$ where $\text{R}_{\text{N}-2}$ and $\text{R}_{\text{N}-3}$ are the same or different and are as defined above,

(L) $-(\text{CH}_2)_{0-4}-\text{R}_{\text{N}-4}$ where $\text{R}_{\text{N}-4}$ is as defined above,

(M) $-\text{O}-\text{CO}-(\text{C}_1-\text{C}_6 \text{ alkyl})$,

(N) $-\text{O}-\text{CO}-\text{NR}_{\text{N}-8}\text{R}_{\text{N}-8}$ where $\text{R}_{\text{N}-8}$ are the same or different
10 and are as defined above,

(O) $-\text{O}-(\text{C}_1-\text{C}_5 \text{ alkyl})-\text{COOH}$,

(P) $-\text{O}-(\text{C}_1-\text{C}_6 \text{ alkyl})$ optionally substituted with one, two, or three of $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, or $-\text{I}$,

(Q) $-\text{NH}-\text{SO}_2-(\text{C}_1-\text{C}_6 \text{ alkyl})$, and

15 (R) $-\text{F}$, or $-\text{Cl}$;

where R_A is:

(I) C_1-C_{10} alkyl optionally substituted with one, two or three substituents selected from the group consisting of C_1-C_3 alkyl, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, $-\text{OH}$,
20 $-\text{SH}$, $-\text{C}\equiv\text{N}$, $-\text{CF}_3$, C_1-C_6 alkoxy, $-\text{O}-\text{phenyl}$, $-\text{NR}_{1-\text{a}}\text{R}_{1-\text{b}}$ where $\text{R}_{1-\text{a}}$ and $\text{R}_{1-\text{b}}$ are as defined above, $-\text{OC}=\text{O NR}_{1-\text{a}}\text{R}_{1-\text{b}}$ where $\text{R}_{1-\text{a}}$ and $\text{R}_{1-\text{b}}$ are as defined above, $-\text{S}(=\text{O})_{0-2}\text{R}_{1-\text{a}}$ where $\text{R}_{1-\text{a}}$ is as defined above, $-\text{NR}_{1-\text{a}}\text{C}=\text{O NR}_{1-\text{a}}\text{R}_{1-\text{b}}$ where $\text{R}_{1-\text{a}}$ and $\text{R}_{1-\text{b}}$ are as defined above, $-\text{C}=\text{O NR}_{1-\text{a}}\text{R}_{1-\text{b}}$ where $\text{R}_{1-\text{a}}$ and $\text{R}_{1-\text{b}}$ are as defined above, and $-\text{S}(=\text{O})_2 \text{NR}_{1-\text{a}}\text{R}_{1-\text{b}}$ where $\text{R}_{1-\text{a}}$ and $\text{R}_{1-\text{b}}$ are as defined above,

25 (II) $-(\text{CH}_2)_{0-3}-(\text{C}_3-\text{C}_8)$ cycloalkyl where cycloalkyl can be optionally substituted with one, two or three substituents selected from the group consisting of C_1-C_3 alkyl, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, $-\text{OH}$, $-\text{SH}$, $-\text{C}\equiv\text{N}$, $-\text{CF}_3$, C_1-C_6 alkoxy, $-\text{O}-\text{phenyl}$, $-\text{CO}-\text{OH}$, $-\text{CO}-\text{O}-(\text{C}_1-\text{C}_4 \text{ alkyl})$, and $-\text{NR}_{1-\text{a}}\text{R}_{1-\text{b}}$ where $\text{R}_{1-\text{a}}$ and $\text{R}_{1-\text{b}}$ are as defined above,

30 (III) $-(\text{CR}_{\text{A}-\text{x}}\text{R}_{\text{A}-\text{y}})_{0-4}-\text{R}_{\text{A}-\text{aryl}}$ where $\text{R}_{\text{A}-\text{x}}$ and $\text{R}_{\text{A}-\text{y}}$ are

(A) $-\text{H}$,

(B) C_1-C_4 alkyl optionally substituted with one or two $-\text{OH}$,

(C) C₁-C₄ alkoxy optionally substituted with one, two, or three of -F,

(D) -(CH₂)₀₋₄-C₃-C₇ cycloalkyl,

(E) C₂-C₆ alkenyl containing one or two double bonds,

5 (F) C₂-C₆ alkynyl containing one or two triple bonds, or

(G) phenyl,

and where R_{A-x} and R_{A-y} are taken together with the carbon to which they are attached to form a carbocycle of three, four, five, six or seven carbon atoms, optionally where one carbon atom is replaced by a heteroatom selected from the

10 group consisting of -O-, -S-, -SO₂-, and -NR_{N-2}- and R_{A-aryl} is the same as R_{N-aryl},

(IV) -(CR_{A-x}R_{A-y})₀₋₄-R_{A-heteroaryl} where R_{A-heteroaryl} is the same as R_{N-heteroaryl} and R_{A-x} and R_{A-y} are as defined above,

(V) -(CR_{A-x}R_{A-y})₀₋₄-R_{A-aryl}-R_{A-aryl} where R_{A-aryl}, R_{A-x} and R_{A-y} are as defined above,

15 (VI) -(CR_{A-x}R_{A-y})₀₋₄-R_{A-aryl}-R_{A-heteroaryl} where R_{A-aryl}, R_{A-heteroaryl}, R_{A-x} and R_{A-y} are as defined above,

(VII) -(CR_{A-x}R_{A-y})₀₋₄-R_{A-heteroaryl}-R_{A-aryl} where R_{A-heteroaryl}, R_{A-aryl}, R_{A-x} and R_{A-y} are as defined above,

20 (VIII) -(CR_{A-x}R_{A-y})₀₋₄-R_{A-heteroaryl}-R_{A-heteroaryl} where R_{A-heteroaryl}, R_{A-x} and R_{A-y} are as defined above,

(IX) -(CR_{A-x}R_{A-y})₀₋₄-R_{A-aryl}-R_{A-heterocycle} where R_{A-heterocycle} is defined as R_{1-heterocycle}, and where R_{A-aryl}, R_{A-x} and R_{A-y} are as defined above,

(X) -(CR_{A-x}R_{A-y})₀₋₄-R_{A-heteroaryl}-R_{A-heterocycle} where R_{A-heteroaryl}, R_{A-heterocycle}, R_{A-x} and R_{A-y} are as defined above,

25 (XI) -(CR_{A-x}R_{A-y})₀₋₄-R_{A-heterocycle}-R_{A-aryl} where R_{A-heterocycle}, R_{A-aryl}, R_{A-x} and R_{A-y} are as defined above,

(XII) -(CR_{A-x}R_{A-y})₀₋₄-R_{A-heterocycle}-R_{A-heteroaryl} where R_{A-heterocycle}, R_{A-heteroaryl}, R_{A-x} and R_{A-y} are as defined above,

30 (XIII) -(CR_{A-x}R_{A-y})₀₋₄-R_{A-heterocycle}-R_{A-heterocycle} where R_{A-heterocycle}, R_{A-x} and R_{A-y} are as defined above,

(XIV) -(CR_{A-x}R_{A-y})₀₋₄-R_{A-heterocycle} where R_{A-heterocycle}, R_{A-x} and R_{A-y} are as defined above,

(XV) -[C(R_{A-1})(R_{A-2})]₁₋₃-CO-N-(R_{A-3})₂ where R_{A-1} and R_{A-2} are the same or different and are selected from the group consisting of:

(A) -H,

(B) -C₁-C₆ alkyl, optionally substituted with one, two or three substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(C) C₂-C₆ alkenyl with one or two double bonds, optionally substituted with one, two or three substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(D) C₂-C₆ alkynyl with one or two triple bonds, optionally substituted with one, two or three substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(E) -(CH₂)₁₋₂-S(O)₀₋₂-(C₁-C₆ alkyl),

(F) -(CH₂)₀₋₄-C₃-C₇ cycloalkyl, optionally substituted with one, two or three substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(G) -(C₁-C₄ alkyl)-R_{A'-aryl} where R_{A'-aryl} is as defined for R_{1-aryl},

(H) -(C₁-C₄ alkyl)-R_{A-heteroaryl} where R_{A-heteroaryl} is as defined above,

(I) -(C₁-C₄ alkyl)-R_{A-heterocycle} where R_{A-heterocycle} is as defined above,

(J) -R_{A-heteroaryl} where R_{A-heteroaryl} is as defined above,

(K) -R_{A-heterocycle} where R_{A-heterocycle} is as defined above,

(M) -(CH₂)₁₋₄-R_{A-4}-(CH₂)₀₋₄-R_{A'-aryl} where R_{A-4} is -O-, -S- or -NR_{A-5}- where R_{A-5} is C₁-C₆ alkyl, and where R_{A'-aryl} is defined above,

(N) -(CH₂)₁₋₄-R_{A-4}-(CH₂)₀₋₄-R_{A-heteroaryl} where R_{A-4} and R_{A-heteroaryl} are as defined above, and

(O) -R_{A'-aryl} where R_{A'-aryl} is as defined above,

and where R_{A-3} is the same or different and is:

(A) -H,

(B) $-C_1-C_6$ alkyl optionally substituted with one, two or three substituents selected from the group consisting of C_1-C_3 alkyl, $-F$, $-Cl$, $-Br$, $-I$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, C_1-C_6 alkoxy, $-O$ -phenyl, $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

5 (C) C_2-C_6 alkenyl with one or two double bonds, optionally substituted with one, two or three substituents selected from the group consisting of C_1-C_3 alkyl, $-F$, $-Cl$, $-Br$, $-I$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, C_1-C_6 alkoxy, $-O$ -phenyl, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

(D) C_2-C_6 alkynyl with one or two triple bonds, optionally substituted with one, two or three substituents selected from the group consisting of C_1-C_3 alkyl, $-F$, $-Cl$, $-Br$, $-I$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, C_1-C_6 alkoxy, $-O$ -phenyl, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

(E) $-(CH_2)_{0-4}-C_3-C_7$ cycloalkyl, optionally substituted with one, two or three substituents selected from the group consisting of C_1-C_3 alkyl, $-F$, $-Cl$, $-Br$, $-I$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, C_1-C_6 alkoxy, $-O$ -phenyl, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

(F) $-R_{A'-aryl}$ where $R_{A'-aryl}$ is as defined above,

(G) $-R_{A-heteroaryl}$ where $R_{A-heteroaryl}$ is as defined above,

(H) $-R_{A-heterocycle}$ where $R_{A-heterocycle}$ is as defined above,

20 (I) $-(C_1-C_4 \text{ alkyl})-R_{A'-aryl}$ where $R_{A'-aryl}$ is as defined above,

(J) $-(C_1-C_4 \text{ alkyl})-R_{A-heteroaryl}$ where $R_{A-heteroaryl}$ is as defined above,

(K) $-(C_1-C_4 \text{ alkyl})-R_{A-heterocycle}$ where $R_{A-heterocycle}$ is as defined above, or

(XVI) $-\text{CH}(R_{A-aryl})_2$ where R_{A-aryl} are the same or different and are as defined above,

(XVII) $-\text{CH}(R_{A-heteroaryl})_2$ where $R_{A-heteroaryl}$ are the same or different and are as defined above,

30 (XVIII) $-\text{CH}(R_{A-aryl})(R_{A-heteroaryl})$ where R_{A-aryl} and $R_{A-heteroaryl}$ are as defined above,

(XIX) $-\text{cyclopentyl}$, $-\text{cyclohexyl}$, or $-\text{cycloheptyl}$ ring fused to R_{A-aryl} , $R_{A-heteroaryl}$, $R_{A-heterocycle}$ where R_{A-aryl} or $R_{A-heteroaryl}$ or $R_{A-heterocycle}$ are as defined above

where one carbon of cyclopentyl, cyclohexyl, or -cycloheptyl is optionally replaced with NH, $\text{NR}_{\text{N-5}}$, O, or $\text{S}(=\text{O})_{0-2}$, and where cyclopentyl, cyclohexyl, or -cycloheptyl can be optionally substituted with one or two $\text{-C}_1\text{-C}_3$ alkyl, -F , -OH , -SH , $\text{-C}\equiv\text{N}$, -CF_3 , $\text{C}_1\text{-C}_6$ alkoxy, $=\text{O}$, or $\text{-NR}_{1-\text{a}}\text{R}_{1-\text{b}}$ where $\text{R}_{1-\text{a}}$ and $\text{R}_{1-\text{b}}$ are as defined above,

5 (XX) $\text{C}_2\text{-C}_{10}$ alkenyl containing one or two double bonds optionally substituted with one, two or three substituents selected from the group consisting of $\text{C}_1\text{-C}_3$ alkyl, -F , -Cl , -Br , -I , -OH , -SH , $\text{-C}\equiv\text{N}$, -CF_3 , $\text{C}_1\text{-C}_6$ alkoxy, -O-phenyl , and $\text{-NR}_{1-\text{a}}\text{R}_{1-\text{b}}$ where $\text{R}_{1-\text{a}}$ and $\text{R}_{1-\text{b}}$ are as defined above,

(XXI) $\text{C}_2\text{-C}_{10}$ alkynyl containing one or two triple bonds optionally substituted with one, two or three substituents selected from the group consisting of $\text{C}_1\text{-C}_3$ alkyl, -F , -Cl , -Br , -I , -OH , -SH , $\text{-C}\equiv\text{N}$, -CF_3 , $\text{C}_1\text{-C}_6$ alkoxy, -O-phenyl , $\text{-NR}_{1-\text{a}}\text{R}_{1-\text{b}}$ where $\text{R}_{1-\text{a}}$ and $\text{R}_{1-\text{b}}$ are as defined above,

(XXI) $\text{-(CH}_2\text{)}_{0-1}\text{-CHR}_{\text{A-6}}\text{-(CH}_2\text{)}_{0-1}\text{-R}_{\text{A-aryl}}$ where $\text{R}_{\text{A-aryl}}$ is as defined above and $\text{R}_{\text{A-6}}$ is $\text{-(CH}_2\text{)}_{0-6}\text{-OH}$,

15 (XXII) $\text{-(CH}_2\text{)}_{0-1}\text{-CHR}_{\text{A-6}}\text{-(CH}_2\text{)}_{0-1}\text{-R}_{\text{A-heteroaryl}}$ where $\text{R}_{\text{A-heteroaryl}}$ and $\text{R}_{\text{A-6}}$ is as defined above,

(XXIII) $\text{-CH(R}_{\text{A-aryl}}\text{ or R}_{\text{A-heteroaryl}}\text{)-CO-O(C}_1\text{-C}_4\text{ alkyl)}$ where $\text{R}_{\text{A-aryl}}$ and $\text{R}_{\text{A-heteroaryl}}$ are as defined above,

(XXIV) $\text{-CH(CH}_2\text{-OH)-CH(OH)-micro-NO}_2$,

20 (XXV) $\text{(C}_1\text{-C}_6\text{ alkyl)-O-(C}_1\text{-C}_6\text{ alkyl)-OH}$,

(XXVII) $\text{-CH}_2\text{-NH-CH}_2\text{-CH(O-CH}_2\text{-CH}_3\text{)}_2$,

(XXVIII) -H ,

(XXIX) $\text{-(CH}_2\text{)}_{0-6}\text{-C(=NR}_{1-\text{a}}\text{)(NR}_{1-\text{a}}\text{R}_{1-\text{b}})$ where $\text{R}_{1-\text{a}}$ and $\text{R}_{1-\text{b}}$ are as defined above; or

25 (XXX)

$\text{-C=OC(HR}_6\text{)NHR}_7$, where R_6 and R_7 are as defined below,

-C=OR_7 , where R_7 is as defined below,

-C=OOR_7 , where R_7 is as defined below, or

-SOOR_7 where R_7 is as defined below,

30 wherein R_6 is:

hydrogen,

$\text{C}_1\text{-C}_3$ alkyl,

phenyl,

thioalkoxyalkyl,
alkyl substituted aryl,
cycloalkyl,
cycloalkylalkyl,
5 hydroxyalkyl,
alkoxyalkyl,
aryloxyalkyl,
haloalkyl,
carboxyalkyl,
10 alkoxycarbonylalkyl,
aminoalkyl,
(N-protected)aminoalkyl,
alkylaminoalkyl,
((N-protected)(alkyl)amino)alkyl,
15 dialkylaminoalkyl,
guanidinoalkyl,
lower alkenyl,
heterocyclic,
(heterocyclic)alkyl),
20 arylthioalkyl,
arylsulfonylalkyl,
(heterocyclic)thioalkyl,
(heterocyclic)sulfonylalkyl,
(heterocyclic)oxyalkyl,
25 arylalkoxyalkyl,
arylthioalkoxyalkyl,
arylalkylsulfonylalkyl,
(heterocyclic))alkoxyalkyl,
(heterocyclic)thioalkoxyalkyl,
30 (heterocyclic)alkylsulfonylalkyl,
cycloalkyloxyalkyl,
cycloalkylthioalkyl,
cycloalkylsulfonylalkyl,
cycloalkylalkoxyalkyl,

- 5 cycloalkylthioalkoxyalkyl,
cycloalkylalkylsulfonylalkyl,
aminocarbonyl,
alkylaminocarbonyl,
dialkylaminocarbonyl,
aroalkyl,
(heterocyclic)carbonylalkyl,
polyhydroxyalkyl,
aminocarbonylalkyl,
10 alkylaminocarbonylalkyl,
dialkylaminocarbonylalkyl,
aryloxyalkyl, or
alkylsulfonylalkyl,
wherein heterocyclic is pyridyl, thiazolyl, isothiazolyl,
15 oxazolyl, isoxazolyl, furanyl, thienyl, tetrahydrofuranyl, tetrahydrothienyl and
tetrahydro[2H]pyranyl and wherein the heterocycle is unsubstituted or substituted
with one to three substituents independently selected from hydroxy, halo, amino,
alkylamino, dialkylamino, alkoxy, polyalkoxy, haloalkyl, cycloalkyl,
cycloalkylalkyl, aryl, arylalkyl, COOH, -SO₃H, lower alkenyl or lower alkyl;
20 wherein R₇ is:
C₁ - C₃ alkyl,
phenyl,
thioalkoxyalkyl,
(aryl)alkyl,
25 cycloalkyl,
cycloalkylalkyl,
hydroxyalkyl,
alkoxyalkyl,
aryloxyalkyl,
30 haloalkyl,
carboxyalkyl,
alkoxycarbonylalkyl,
aminoalkyl,
(N-protected)aminocalkyl,

alkylaminoalkyl,
((N-protected)(alkyl)amino)alkyl,
dialkylaminoalkyl,
guanidinoalkyl,
5 lower alkenyl,
heterocyclic,
(heterocyclic)alkyl),
arylthioalkyl,
arylsulfonylalkyl,
10 (heterocyclic)thioalkyl,
(heterocyclic)sulfonylalkyl,
(heterocyclic)oxyalkyl,
arylalkoxyalkyl,
arylthioalkoxyalkyl,
15 arylalkylsulfonylalkyl,
(heterocyclic)alkoxyalkyl,
(heterocyclic)thioalkoxyalkyl,
(heterocyclic)alkylsulfonylalkyl,
cycloalkyloxyalkyl,
20 cycloalkylthioalkyl,
cycloalkylsulfonylalkyl,
cycloalkylalkoxyalkyl,
cycloalkylthioalkoxyalkyl,
cycloalkylalkylsulfonylalkyl,
25 aminocarbonyl,
alkylaminocarbonyl,
dialkylaminocarbonyl,
aroylalkyl,
(heterocyclic)carbonylalkyl,
30 polyhydroxyalkyl,
aminocarbonylalkyl,
alkylaminocarbonylalkyl,
dialkylaminocarbonylalkyl,
aryloxyalkyl, or

alkylsulfonylalkyl,

wherein heterocyclic is pyridyl, thiazolyl, isothiazolyl,

oxazolyl, isoxazolyl, furanyl, thienyl, tetrahydrofuranyl, tetrahydrothienyl and tetrahydro[2H]pyranyl and wherein the heterocycle is unsubstituted or substituted

5 with one to three substituents independently selected from hydroxy, halo, amino, alkylamino, dialkylamino, alkoxy, polyalkoxy, haloalkyl, cycloalkyl, cycloalkylalkyl, aryl, arylalkyl, COOH, -SO₃H, lower alkenyl or lower alkyl; and

where PROTECTING GROUP is selected from the group consisting of *t*-

butoxycarbonyl, benzyloxycarbonyl, formyl, trityl, acetyl, trichloroacetyl,

10 dichloroacetyl, chloroacetyl, trifluoroacetyl, difluoroacetyl, fluoroacetyl, 4-

phenylbenzyloxycarbonyl, 2-methylbenzyloxycarbonyl, 4-

ethoxybenzyloxycarbonyl, 4-fluorobenzyloxycarbonyl, 4-chlorobenzyloxycarbonyl,

3-chlorobenzyloxycarbonyl, 2-chlorobenzyloxycarbonyl, 2,4-

dichlorobenzyloxycarbonyl, 4-bromobenzyloxycarbonyl, 3-

15 bromobenzyloxycarbonyl, 4-nitrobenzyloxycarbonyl, 4-cyanobenzyloxycarbonyl, 2-

(4-xenyl)isopropoxycarbonyl, 1,1-diphenyleth-1-yloxycarbonyl, 1,1-diphenylprop-

1-yloxycarbonyl, 2-phenylprop-2-yloxycarbonyl, 2-(*p*-toluyl)prop-2-yloxycarbonyl,

cyclopentanyloxycarbonyl, 1-methylcyclopentanyloxycarbonyl,

cyclohexanyloxycarbonyl, 1-methylcyclohexanyloxycarbonyl, 2-

20 methylcyclohexanyloxycarbonyl, 2-(4-toluylsulfonyl)ethoxycarbonyl, 2-

(methylsulfonyl)ethoxycarbonyl, 2-(triphenylphosphino)ethoxycarbonyl,

fluorenylmethoxycarbonyl, 2-(trimethylsilyl)ethoxycarbonyl, allyloxycarbonyl, 1-

(trimethylsilylmethyl)prop-1-enyloxycarbonyl, 5-benzisoxalylmethoxycarbonyl, 4-

acetoxycarbonyl, 2,2,2-trichloroethoxycarbonyl, 2-ethynyl-2-

25 propoxycarbonyl, cyclopropylmethoxycarbonyl, 4-(decyloxy)benzyloxycarbonyl,

isobornyloxycarbonyl and 1-piperidyloxycarbonyl, 9-fluorenylmethyl carbonate, -

CH-CH=CH₂ and phenyl-C(=N-)-H.

47. A protected compound according to claim 46,

30 where R₁ is:

-(CH₂)₀₋₁-(R_{1-aryl}), or

-(CH₂)_{n1}-(R_{1-heteroaryl});

where R_A is:

-C₁-C₈ alkyl,

- 5 $-(\text{CH}_2)_{0-3}-(\text{C}_3-\text{C}_7)$ cycloalkyl,
 $-(\text{CR}_{\text{A}-\text{x}}\text{R}_{\text{A}-\text{y}})_{0-4}-\text{R}_{\text{A-aryl}}$,
 $-(\text{CR}_{\text{A}-\text{x}}\text{R}_{\text{A}-\text{y}})_{0-4}-\text{R}_{\text{A-heteroaryl}}$,
 $-(\text{CR}_{\text{A}-\text{x}}\text{R}_{\text{A}-\text{y}})_{0-4}-\text{R}_{\text{A-heterocycle}}$,
 -cyclopentyl or -cyclohexyl ring fused to $\text{R}_{\text{A-aryl}}$ or $\text{R}_{\text{A-heteroaryl}}$ or $\text{R}_{\text{A-}}$
 heterocycle.

48. A protected compound according to claim 47

where R_1 is:

- 10 $-(\text{CH}_2)-(\text{R}_{1\text{-aryl}})$, or
 $-(\text{CH}_2)-(\text{R}_{1\text{-heteroaryl}})$;

where R_2 is $-\text{H}$;

where R_3 is $-\text{H}$;

where R_{A} is:

- 15 $-\text{C}_1-\text{C}_8$ alkyl,
 $-(\text{CH}_2)_{0-3}-(\text{C}_3-\text{C}_7)$ cycloalkyl,
 $-(\text{CR}_{\text{A}-\text{x}}\text{R}_{\text{A}-\text{y}})_{0-4}-\text{R}_{\text{A-aryl}}$,
 $-(\text{CR}_{\text{A}-\text{x}}\text{R}_{\text{A}-\text{y}})_{0-4}-\text{R}_{\text{A-heteroaryl}}$,
 $-(\text{CR}_{\text{A}-\text{x}}\text{R}_{\text{A}-\text{y}})_{0-4}-\text{R}_{\text{A-heterocycle}}$,
 20 -cyclopentyl or -cyclohexyl ring fused to $\text{R}_{\text{A-aryl}}$ or $\text{R}_{\text{A-heteroaryl}}$ or $\text{R}_{\text{A-}}$
 heterocycle.


49. A protected compound according to claim 46 where PROTECTING GROUP is *t*-butoxycarbonyl.

25

50. A protected compound according to claim 46 where PROTECTING GROUP is benzyloxycarbonyl.

51. A compound of the formula (XII)

30



XII

where R_1 is:

- (I) C_1 - C_6 alkyl, optionally substituted with one, two or three substituents selected from the group consisting of C_1 - C_3 alkyl, C_1 - C_7 alkyl (optionally substituted with C_1 - C_3 alkyl and C_1 - C_3 alkoxy), -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C_1 - C_3 alkoxy, -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are -H or C_1 - C_6 alkyl, and -OC=O NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,
- (II) -CH₂-S(O)₀₋₂-(C_1 - C_6 alkyl),
- (III) -CH₂-CH₂-S(O)₀₋₂-(C_1 - C_6 alkyl),
- (IV) C_2 - C_6 alkenyl with one or two double bonds, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, -C≡N, -CF₃, C_1 - C_3 alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are -H or C_1 - C_6 alkyl,
- (V) C_2 - C_6 alkynyl with one or two triple bonds, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, -C≡N, -CF₃, C_1 - C_3 alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are -H or C_1 - C_6 alkyl,
- (VI) -(CH₂)_{n1}-(R_{1-aryl}) where n₁ is zero or one and where R_{1-aryl} is phenyl, 1-naphthyl, 2-naphthyl and indanyl, indenyl, dihydronaphthalyl, or tetralinyl optionally substituted with one, two, three or four of the following substituents on the aryl ring:
- (A) C_1 - C_6 alkyl optionally substituted with one, two or three substituents selected from the group consisting of C_1 - C_3 alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, and C_1 - C_3 alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,
- (B) C_2 - C_6 alkenyl with one or two double bonds, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, -C≡N, -CF₃, C_1 - C_3 alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are -H or C_1 - C_6 alkyl,
- (C) C_2 - C_6 alkynyl with one or two triple bonds, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, -C≡N, -CF₃, C_1 - C_3 alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are -H or C_1 - C_6 alkyl,

(D) -F, Cl, -Br or -I,

(F) -C₁-C₆ alkoxy optionally substituted with one, two or three of - F,

(G) -NR_{N-2}R_{N-3} where R_{N-2} and R_{N-3} are as defined below,

5 (H) -OH,

(I) -C≡N,

(J) C₃-C₇ cycloalkyl, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, -C≡N, -CF₃, C₁-C₃ alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are -H or C₁-C₆ alkyl,

10 (K) -CO-(C₁-C₄ alkyl),

(L) -SO₂-NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(M) -CO-NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

or

(N) -SO₂-(C₁-C₄ alkyl),

15 (VII) -(CH₂)_{n1}-(R_{1-heteroaryl}) where n₁ is as defined above and where R_{1-heteroaryl} is selected from the group consisting of:

pyridinyl,

pyrimidinyl,

quinolinyl,

20 benzothienyl,

indolyl,

indolinyl,

pyridazinyl,

pyrazinyl,

25 isoquinolyl,

quinazolinyl,

quinoxalinyl,

phthalazinyl,

imidazolyl,

30 isoxazolyl,

pyrazolyl,

oxazolyl,

thiazolyl,

	indoliziny1,
	indazolyl,
	benzothiazolyl,
	benzimidazolyl,
5	benzofuranyl,
	furanyl,
	thienyl,
	pyrrolyl,
	oxadiazolyl,
10	thiadiazolyl,
	triazolyl,
	tetrazolyl,
	oxazolopyridinyl,
	imidazopyridinyl,
15	isothiazolyl,
	naphthyridinyl,
	cinnolinyl,
	carbazolyl,
	beta-carbolinyl,
20	isochromanyl,
	chromanyl,
	tetrahydroisoquinolinyl,
	isoindolinyl,
	isobenzotetrahydrofuranyl,
25	isobenzotetrahydrothienyl,
	isobenzothienyl,
	benzoxazolyl,
	pyridopyridinyl,
	benzotetrahydrofuranyl,
30	benzotetrahydrothienyl,
	purinyl,
	benzodioxolyl,
	triazinyl,
	phenoxazinyl,

	phenothiazinyl,
	pteridinyl,
	benzothiazolyl,
	imidazopyridinyl,
5	imidazothiazolyl,
	dihydrobenzisoxazinyl,
	benzisoxazinyl,
	benzoxazinyl,
	dihydrobenziso­thiazinyl,
10	benzopyranyl,
	benzothiopyranyl,
	coumarinyl,
	isocoumarinyl,
	chromonyl,
15	chromanonyl,
	pyridinyl-N-oxide,
	tetrahydroquinolinyl
	dihydroquinolinyl
	dihydroquinolinonyl
20	dihydroisoquinolinonyl
	dihydrocoumarinyl
	dihydroisocoumarinyl
	isoindolinonyl
	benzodioxanyl
25	benzoxazolinonyl
	pyrrolyl N-oxide,
	pyrimidinyl N-oxide,
	pyridazinyl N-oxide,
	pyrazinyl N-oxide,
30	quinolinyl N-oxide,
	indolyl N-oxide,
	indolinyl N-oxide,
	isoquinolyl N-oxide,
	quinazolinyl N-oxide,

quinoxaliny N-oxide,
 phthalaziny N-oxide,
 imidazoly N-oxide,
 isoxazoly N-oxide,
 5 oxazoly N-oxide,
 thiazoly N-oxide,
 indoliziny N-oxide,
 indazoly N-oxide,
 benzothiazoly N-oxide,
 10 benzimidazoly N-oxide,
 pyrroly N-oxide,
 oxadiazoly N-oxide,
 thiadiazoly N-oxide,
 triazoly N-oxide,
 15 tetrazoly N-oxide,
 benzothiopyrany S-oxide, and
 benzothiopyrany S,S-dioxide,

where the $R_{1\text{-heteroaryl}}$ group is bonded to $-(CH_2)_{n1}-$ by any ring atom
 of the parent $R_{1\text{-heteroaryl}}$ group substituted by hydrogen such that the new bond to the
 20 $R_{1\text{-heteroaryl}}$ group replaces the hydrogen atom and its bond, where heteroaryl is
 optionally substituted with one, two, three or four of:

(1) C_1-C_6 alkyl optionally substituted with one, two or
 three substituents selected from the group consisting of C_1-C_3 alkyl, -F, -Cl, -Br, -I,
 -OH, -SH, $-C\equiv N$, $-CF_3$, C_1-C_3 alkoxy, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as
 25 defined above,

(2) C_2-C_6 alkenyl with one or two double bonds,
 optionally substituted with one, two or three substituents selected from the group
 consisting of -F, -Cl, -OH, -SH, $-C\equiv N$, $-CF_3$, C_1-C_3 alkoxy, and $-NR_{1-a}R_{1-b}$ where R_{1-}
 a and R_{1-b} are -H or C_1-C_6 alkyl,

(3) C_2-C_6 alkynyl with one or two triple bonds,
 optionally substituted with one, two or three substituents selected from the group
 consisting of -F, -Cl, -OH, -SH, $-C\equiv N$, $-CF_3$, C_1-C_3 alkoxy, and $-NR_{1-a}R_{1-b}$ where R_{1-}
 a and R_{1-b} are -H or C_1-C_6 alkyl,

- (4) -F, Cl, -Br or -I,
(6) -C₁-C₆ alkoxy optionally substituted with one, two,
or three of -F,
(7) -NR_{N-2}R_{N-3} where R_{N-2} and R_{N-3} are as defined
5 below,
(8) -OH,
(9) -C≡N,
(10) C₃-C₇ cycloalkyl, optionally substituted with one,
two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, -
10 C≡N, -CF₃, C₁-C₃ alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are -H or C₁-C₆ alkyl,
(11) -CO-(C₁-C₄ alkyl),
(12) -SO₂-NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined
above,
(13) -CO-NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined
15 above, or
(14) -SO₂-(C₁-C₄ alkyl), with the proviso that when n₁
is zero R_{1-heteroaryl} is not bonded to the carbon chain by nitrogen; or
(VIII) -(CH₂)_{n1}-(R_{1-heterocycle}) where n₁ is as defined above and R_{1-heterocycle}
heterocycle is selected from the group consisting of:
20 morpholinyl,
thiomorpholinyl,
thiomorpholinyl S-oxide,
thiomorpholinyl S,S-dioxide,
piperazinyl,
25 homopiperazinyl,
pyrrolidinyl,
pyrrolinyl,
tetrahydropyranyl,
piperidinyl,
30 tetrahydrofuranyl,
tetrahydrothienyl,
homopiperidinyl,
homomorpholinyl,

homothiomorpholinyl,
 homothiomorpholinyl S,S-dioxide,
 oxazolidinonyl,
 dihydropyrazolyl,
 5 dihydropyrrolyl,
 dihydropyrazinyl,
 dihydropyridinyl,
 dihydropyrimidinyl,
 dihydrofuryl,
 10 dihydropyranyl,
 tetrahydrothienyl S-oxide,
 tetrahydrothienyl S,S-dioxide, and
 homothiomorpholinyl S-oxide,

where the $R_{1\text{-heterocycle}}$ group is bonded by any atom of the parent $R_{1\text{-heterocycle}}$ group substituted by hydrogen such that the new bond to the $R_{1\text{-heterocycle}}$ group replaces the hydrogen atom and its bond, where heterocycle is optionally substituted with one, two, three or four:

(1) $C_1\text{-}C_6$ alkyl optionally substituted with one, two or three substituents selected from the group consisting of $C_1\text{-}C_3$ alkyl, -F, -Cl, -Br, -I, -OH, -SH, $-C\equiv N$, $-CF_3$, $C_1\text{-}C_3$ alkoxy, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

(2) $C_2\text{-}C_6$ alkenyl with one or two double bonds, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, $-C\equiv N$, $-CF_3$, $C_1\text{-}C_3$ alkoxy, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are -H or $C_1\text{-}C_6$ alkyl,

(3) $C_2\text{-}C_6$ alkynyl with one or two triple bonds, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, $-C\equiv N$, $-CF_3$, $C_1\text{-}C_3$ alkoxy, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are -H or $C_1\text{-}C_6$ alkyl,

(4) -F, Cl, -Br or -I,

(5) $C_1\text{-}C_6$ alkoxy,

(6) $-C_1\text{-}C_6$ alkoxy optionally substituted with one,

two, or three -F,

(7) $-NR_{N-2}R_{N-3}$ where R_{N-2} and R_{N-3} are as defined

below,

(8) $-OH$,

(9) $-C\equiv N$,

5 (10) C_3-C_7 cycloalkyl, optionally substituted with one, two or three substituents selected from the group consisting of $-F$, $-Cl$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, C_1-C_3 alkoxy, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are $-H$ or C_1-C_6 alkyl,

(11) $-CO-(C_1-C_4 \text{ alkyl})$,

(12) $-SO_2-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined

10 above,

(13) $-CO-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined

above,

(14) $-SO_2-(C_1-C_4 \text{ alkyl})$, or

(15) $=O$, with the proviso that when n_1 is zero R_1 -

15 heterocycle is not bonded to the carbon chain by nitrogen;

where R_2 is:

(I) $-H$,

(II) C_1-C_6 alkyl, optionally substituted with one, two or three
20 substituents selected from the group consisting of C_1-C_3 alkyl, $-F$, $-Cl$, $-Br$, $-I$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, C_1-C_3 alkoxy, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

(III) $-(CH_2)_{0-4}-R_{2-1}$ where R_{2-1} is R_{1-aryl} or $R_{1-heteroaryl}$ where R_{1-aryl} and $R_{1-heteroaryl}$ are as defined above;

25 (IV) C_2-C_6 alkenyl with one or two double bonds, optionally substituted with one, two or three substituents selected from the group consisting of $-F$, $-Cl$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, C_1-C_3 alkoxy, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are $-H$ or C_1-C_6 alkyl,

(V) C_2-C_6 alkynyl with one or two triple bonds, optionally substituted
30 with one, two or three substituents selected from the group consisting of $-F$, $-Cl$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, C_1-C_3 alkoxy, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are $-H$ or C_1-C_6 alkyl, or

(VI) $-(CH_2)_{0-4}-C_3-C_7$ cycloalkyl, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, $-C\equiv N$, $-CF_3$, C_1-C_3 alkoxy, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are -H or C_1-C_6 alkyl;

5 where R_3 is:

(I)-H,

(II) C_1-C_6 alkyl, optionally substituted with one, two or three substituents selected from the group consisting of C_1-C_3 alkyl, -F, -Cl, -Br, -I, -OH, -SH, $-C\equiv N$, $-CF_3$, C_1-C_3 alkoxy, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined

10 above,

(III) $-(CH_2)_{0-4}-R_{2-1}$ where R_{2-1} is R_{1-aryl} or $R_{1-heteroaryl}$ where R_{1-aryl} and $R_{1-heteroaryl}$ are as defined above

(IV) C_2-C_6 alkenyl with one or two double bonds,

(V) C_2-C_6 alkynyl with one or two triple bonds; or

15 (VI) $-(CH_2)_{0-4}-C_3-C_7$ cycloalkyl, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, $-C\equiv N$, $-CF_3$, C_1-C_3 alkoxy, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are -H or C_1-C_6 alkyl,

and where R_2 and R_3 are taken together with the carbon to which they are attached to form a carbocycle of three, four, five, six, or seven carbon atoms,

20 optionally where one carbon atom is replaced by a heteroatom selected from the group consisting of -O-, -S-, $-SO_2-$, and $-NR_{N-2}-$, where R_{N-2} is selected from the group consisting of:

(a) -H,

(b) C_1-C_6 alkyl optionally substituted with one

25 substituent selected from the group consisting of:

(i) -OH, and

(ii) $-NH_2$,

(c) C_1-C_6 alkyl optionally substituted with one, two, or three -F, -Cl, -Br, or -I,

30 (d) C_3-C_7 cycloalkyl,

(e) $-(C_1-C_2 \text{ alkyl})-(C_3-C_7 \text{ cycloalkyl})$,

(f) $-(C_1-C_6 \text{ alkyl})-O-(C_1-C_3 \text{ alkyl})$,

bonds,
 (g) $-C_2-C_6$ alkenyl with one or two double

bonds,
 (h) $-C_2-C_6$ alkynyl with one or two triple

5 and one triple bond,
 (i) $-C_1-C_6$ alkyl chain with one double bond

(j) $-R_{1-aryl}$ where R_{1-aryl} is as defined above, and

(k) $-R_{1-heteroaryl}$ where $R_{1-heteroaryl}$ is as defined
 above;

10

where R_N is:

(I) $R_{N-1}-X_N$ where X_N is selected from the group consisting of:

(A) $-CO-$,

(B) $-SO_2-$,

15 (C) $-(CR'R'')_{1-6}$ where R' and R'' are the same or different and
 are $-H$ and C_1-C_4 alkyl,

(D) $-CO-(CR'R'')_{1-6}-X_{N-1}$ where X_{N-1} is selected from the
 group consisting of $-O-$, $-S-$ and $-NR'-$ and where R' and R'' are as defined above,
 and

20 (E) a single bond;

where R_{N-1} is selected from the group consisting of:

(A) R_{N-aryl} where R_{N-aryl} is phenyl, 1-naphthyl, 2-naphthyl,
 tetralinyl, indanyl, 6,7,8,9-tetrahydro-5H-benzo[a]cycloheptenyl, or dihydronaphthyl
 optionally substituted with one, two or three of the following substituents which can
 25 be the same or different and are:

(1) C_1-C_6 alkyl, optionally substituted with one, two or
 three substituents selected from the group consisting of C_1-C_3 alkyl, $-F$, $-Cl$, $-Br$, $-I$,
 $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, C_1-C_3 alkoxy, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as
 defined above,

30

(2) $-OH$,

(3) $-NO_2$,

(4) $-F$, $-Cl$, $-Br$, $-I$,

(5) $-CO-OH$,

(6) $-C\equiv N$,

(7) $-(CH_2)_{0-4}-CO-NR_{N-2}R_{N-3}$ where R_{N-2} and R_{N-3} are the same or different and are selected from the group consisting of:

(a) $-H$,

5 (b) $-C_1-C_6$ alkyl optionally substituted with one substituent selected from the group consisting of:

(i) $-OH$, and

(ii) $-NH_2$,

10 (c) $-C_1-C_6$ alkyl optionally substituted with one, two, or three $-F$, $-Cl$, $-Br$, or $-I$,

(d) $-C_3-C_7$ cycloalkyl,

(e) $-(C_1-C_2 \text{ alkyl})-(C_3-C_7 \text{ cycloalkyl})$,

(f) $-(C_1-C_6 \text{ alkyl})-O-(C_1-C_3 \text{ alkyl})$,

15 (g) $-C_2-C_6$ alkenyl with one or two double bonds,

(h) $-C_2-C_6$ alkynyl with one or two triple

bonds,

(i) $-C_1-C_6$ alkyl chain with one double bond

and one triple bond,

20 (j) $-R_{1-aryl}$ where R_{1-aryl} is as defined above, and

(k) $-R_{1-heteroaryl}$ where $R_{1-heteroaryl}$ is as defined

above,

(8) $-(CH_2)_{0-4}-CO-(C_1-C_{12} \text{ alkyl})$,

25 (9) $-(CH_2)_{0-4}-CO-(C_2-C_{12} \text{ alkenyl with one, two or three double bonds})$,

(10) $-(CH_2)_{0-4}-CO-(C_2-C_{12} \text{ alkynyl with one, two or$

three triple bonds),

(11) $-(CH_2)_{0-4}-CO-(C_3-C_7 \text{ cycloalkyl})$,

(12) $-(CH_2)_{0-4}-CO-R_{1-aryl}$ where R_{1-aryl} is as defined

30 above,

(13) $-(CH_2)_{0-4}-CO-R_{1-heteroaryl}$ where $R_{1-heteroaryl}$ is as

defined above,

(14) $-(\text{CH}_2)_{0-4}-\text{CO}-\text{R}_{1-\text{heterocycle}}$ where $\text{R}_{1-\text{heterocycle}}$ is as defined above,

(15) $-(\text{CH}_2)_{0-4}-\text{CO}-\text{R}_{\text{N}-4}$ where $\text{R}_{\text{N}-4}$ is selected from the group consisting of morpholinyl, thiomorpholinyl, piperazinyl, piperidinyl, homomorpholinyl, homothiomorpholinyl, homothiomorpholinyl S-oxide, homothiomorpholinyl S,S-dioxide, pyrrolinyl and pyrrolidinyl where each group is optionally substituted with one, two, three, or four of C_1-C_6 alkyl,

(16) $-(\text{CH}_2)_{0-4}-\text{CO}-\text{O}-\text{R}_{\text{N}-5}$ where $\text{R}_{\text{N}-5}$ is selected from the group consisting of:

- (a) C_1-C_6 alkyl,
- (b) $-(\text{CH}_2)_{0-2}-(\text{R}_{1-\text{aryl}})$ where $\text{R}_{1-\text{aryl}}$ is as defined above,
- (c) C_2-C_6 alkenyl containing one or two double bonds,
- (d) C_2-C_6 alkynyl containing one or two triple bonds,
- (e) C_3-C_7 cycloalkyl, and
- (f) $-(\text{CH}_2)_{0-2}-(\text{R}_{1-\text{heteroaryl}})$ where $\text{R}_{1-\text{heteroaryl}}$ is as defined above,

(17) $-(\text{CH}_2)_{0-4}-\text{SO}_2-\text{NR}_{\text{N}-2}\text{R}_{\text{N}-3}$ where $\text{R}_{\text{N}-2}$ and $\text{R}_{\text{N}-3}$ are as defined above,

(18) $-(\text{CH}_2)_{0-4}-\text{SO}-(\text{C}_1-\text{C}_8 \text{ alkyl}),$

(19) $-(\text{CH}_2)_{0-4}-\text{SO}_2-(\text{C}_1-\text{C}_{12} \text{ alkyl}),$

(20) $-(\text{CH}_2)_{0-4}-\text{SO}_2-(\text{C}_3-\text{C}_7 \text{ cycloalkyl}),$

(21) $-(\text{CH}_2)_{0-4}-\text{N}(\text{H or } \text{R}_{\text{N}-5})-\text{CO}-\text{O}-\text{R}_{\text{N}-5}$ where $\text{R}_{\text{N}-5}$ can be the same or different and is as defined above,

(22) $-(\text{CH}_2)_{0-4}-\text{N}(\text{H or } \text{R}_{\text{N}-5})-\text{CO}-\text{N}(\text{R}_{\text{N}-5})_2,$ where $\text{R}_{\text{N}-5}$ can be the same or different and is as defined above,

(23) $-(\text{CH}_2)_{0-4}-\text{N}-\text{CS}-\text{N}(\text{R}_{\text{N}-5})_2,$ where $\text{R}_{\text{N}-5}$ can be the same or different and is as defined above,

(24) $-(\text{CH}_2)_{0-4}-\text{N}(-\text{H or } \text{R}_{\text{N}-5})-\text{CO}-\text{R}_{\text{N}-2}$ where $\text{R}_{\text{N}-5}$ and $\text{R}_{\text{N}-2}$ can be the same or different and are as defined above,

(25) $-(\text{CH}_2)_{0-4}-\text{NR}_{\text{N}-2}\text{R}_{\text{N}-3}$ where $\text{R}_{\text{N}-2}$ and $\text{R}_{\text{N}-3}$ can be the same or different and are as defined above,

- (26) $-(\text{CH}_2)_{0-4}-\text{R}_{\text{N}-4}$ where $\text{R}_{\text{N}-4}$ is as defined above,
- (27) $-(\text{CH}_2)_{0-4}-\text{O}-\text{CO}-(\text{C}_1-\text{C}_6 \text{ alkyl})$,
- (28) $-(\text{CH}_2)_{0-4}-\text{O}-\text{P}(\text{O})-(\text{OR}_{\text{N-aryl-1}})_2$ where $\text{R}_{\text{N-aryl-1}}$ is —
 H or C_1-C_4 alkyl,
- 5 (29) $-(\text{CH}_2)_{0-4}-\text{O}-\text{CO}-\text{N}(\text{R}_{\text{N-5}})_2$ where $\text{R}_{\text{N-5}}$ is as
 defined above,
- (30) $-(\text{CH}_2)_{0-4}-\text{O}-\text{CS}-\text{N}(\text{R}_{\text{N-5}})_2$ where $\text{R}_{\text{N-5}}$ is as defined
 above,
- (31) $-(\text{CH}_2)_{0-4}-\text{O}-(\text{R}_{\text{N-5}})_2$ where $\text{R}_{\text{N-5}}$ is as defined
 10 above,
- (32) $-(\text{CH}_2)_{0-4}-\text{O}-(\text{R}_{\text{N-5}})_2-\text{COOH}$ where $\text{R}_{\text{N-5}}$ is as
 defined above,
- (33) $-(\text{CH}_2)_{0-4}-\text{S}-(\text{R}_{\text{N-5}})_2$ where $\text{R}_{\text{N-5}}$ is as defined
 above,
- 15 (34) $-(\text{CH}_2)_{0-4}-\text{O}-(\text{C}_1-\text{C}_6 \text{ alkyl optionally substituted}$
 with one, two, three, four, or five of $-\text{F}$),
- (35) C_3-C_7 cycloalkyl,
- (36) C_2-C_6 alkenyl with one or two double bonds
 optionally substituted with C_1-C_3 alkyl, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, $-\text{OH}$, $-\text{SH}$, $-\text{C}\equiv\text{N}$, $-\text{CF}_3$, C_1-
 20 C_3 alkoxy, or $-\text{NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,
- (37) C_2-C_6 alkynyl with one or two triple bonds
 optionally substituted with C_1-C_3 alkyl, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, $-\text{OH}$, $-\text{SH}$, $-\text{C}\equiv\text{N}$, $-\text{CF}_3$, C_1-
 C_3 alkoxy, or $-\text{NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,
- (38) $-(\text{CH}_2)_{0-4}-\text{N}(-\text{H or } \text{R}_{\text{N-5}})-\text{SO}_2-\text{R}_{\text{N-2}}$ where $\text{R}_{\text{N-5}}$ and
 25 $\text{R}_{\text{N-2}}$ can be the same or different and are as described above, or
- (39) $-(\text{CH}_2)_{0-4}-\text{C}_3-\text{C}_7$ cycloalkyl,
- (B) $-\text{R}_{\text{N-heteroaryl}}$ where $\text{R}_{\text{N-heteroaryl}}$ is selected from the group
 consisting of:
- 30 pyridinyl,
 pyrimidinyl,
 quinolinyl,
 benzothienyl,
 indolyl,

	indolinyI,
	pyridazinyI,
	pyrazinyI,
	isoindolyI,
5	isoquinolyI,
	quinazolinyl,
	quinoxalinyI,
	phthalazinyI,
	imidazolyl,
10	isoxazolyl,
	pyrazolyl,
	oxazolyl,
	thiazolyl,
	indolizinyI,
15	indazolyl,
	benzothiazolyl,
	benzimidazolyl,
	benzofuranyl,
	furanyl,
20	thienyl,
	pyrrolyl,
	oxadiazolyl,
	thiadiazolyl,
	triazolyl,
25	tetrazolyl,
	oxazolopyridinyI,
	imidazopyridinyI,
	isothiazolyl,
	naphthyridinyI,
30	cinnolinyI,
	carbazolyl,
	beta-carbolinyI,
	isochromanyl,
	chromanyl,

	tetrahydroisoquinolinyI,
	isoindolinyI,
	isobenzotetrahydrofuranyI,
	isobenzotetrahydrothienyI,
5	isobenzothiényI,
	benzoxazolyl,
	pyridopyridinyI,
	benzotetrahydrofuranyI,
	benzotetrahydrothienyI,
10	purinyI,
	benzodioxolyl,
	triazinyI,
	phenoxazinyI,
	phenothiazinyI,
15	pteridinyI,
	benzothiazolyl,
	imidazopyridinyI,
	imidazothiazolyl,
	dihydrobenzisoXazinyI,
20	benzisoXazinyI,
	benzoxazinyI,
	dihydrobenzisothiazinyI,
	benzopyranyI,
	benzothiopyranyI,
25	coumarinyI,
	isocoumarinyI,
	chromonyI,
	chromanonyI,
	pyridinyI-N-oxide,
30	tetrahydroquinolinyI,
	dihydroquinolinyI,
	dihydroquinolinonyI,
	dihydroisoquinolinonyI,
	dihydrocoumarinyI,

	dihydroisocoumarinyl,
	isoindolinonyl,
	benzodioxanyl,
	benzoxazolinonyl,
5	pyrrolyl N-oxide,
	pyrimidinyl N-oxide,
	pyridazinyl N-oxide,
	pyrazinyl N-oxide,
	quinolinyl N-oxide,
10	indolyl N-oxide,
	indolinyl N-oxide,
	isoquinolyl N-oxide,
	quinazolinyl N-oxide,
	quinoxalinyl N-oxide,
15	phthalazinyl N-oxide,
	imidazolyl N-oxide,
	isoxazolyl N-oxide,
	oxazolyl N-oxide,
	thiazolyl N-oxide,
20	indoliziny N-oxide,
	indazolyl N-oxide,
	benzothiazolyl N-oxide,
	benzimidazolyl N-oxide,
	pyrrolyl N-oxide,
25	oxadiazolyl N-oxide,
	thiadiazolyl N-oxide,
	triazolyl N-oxide,
	tetrazolyl N-oxide,
	benzothiopyranyl S-oxide, and
30	benzothiopyranyl S,S-dioxide

where the $R_{N\text{-heteroaryl}}$ group is bonded by any atom of the parent $R_{N\text{-heteroaryl}}$ group substituted by hydrogen such that the new bond to the $R_{N\text{-heteroaryl}}$ group replaces the hydrogen atom and its bond, where heteroaryl is optionally substituted with one, two, three, or four of:

- (1) C₁-C₆ alkyl, optionally substituted with one, two or three substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₃ alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,
- 5 (2) -OH,
 (3) -NO₂,
 (4) -F, -Cl, -Br, or -I,
 (5) -CO-OH,
 (6) -C≡N,
- 10 (7) -(CH₂)₀₋₄-CO-NR_{N-2}R_{N-3} where R_{N-2} and R_{N-3} are the same or different and are selected from the group consisting of:
- (a) -H,
 (b) -C₁-C₆ alkyl optionally substituted with one substituent selected from the group consisting of:
- 15 (i) -OH, and
 (ii) -NH₂,
 (c) -C₁-C₆ alkyl optionally substituted with one, two, or three -F, -Cl, -Br, -I,
 (d) -C₃-C₇ cycloalkyl,
 (e) -(C₁-C₂ alkyl)-(C₃-C₇ cycloalkyl),
 (f) -(C₁-C₆ alkyl)-O-(C₁-C₃ alkyl),
 (g) -C₂-C₆ alkenyl with one or two double bonds,
 (h) -C₂-C₆ alkynyl with one or two triple bonds,
 (i) -C₁-C₆ alkyl chain with one double bond and one triple bond,
 (j) -R_{1-aryl} where R_{1-aryl} is as defined above,
 (k) -R_{1-heteroaryl} where R_{1-heteroaryl} is as defined above,
- 20
 25
 30
- (8) -(CH₂)₀₋₄-CO-(C₁-C₁₂ alkyl),
 (9) -(CH₂)₀₋₄-CO-(C₂-C₁₂ alkenyl with one, two or three double bonds),

- (10) $-(CH_2)_{0-4}-CO-(C_2-C_{12}$ alkynyl with one, two or three triple bonds),
- (11) $-(CH_2)_{0-4}-CO-(C_3-C_7$ cycloalkyl),
- (12) $-(CH_2)_{0-4}-CO-R_{1-aryl}$ where R_{1-aryl} is as defined above,
- (13) $-(CH_2)_{0-4}-CO-R_{1-heteroaryl}$ where $R_{1-heteroaryl}$ is as defined above,
- (14) $-(CH_2)_{0-4}-CO-R_{1-heterocycle}$ where $R_{1-heterocycle}$ is as defined above,
- (15) (15) $-(CH_2)_{0-4}-CO-R_{N-4}$ where R_{N-4} is selected from the group consisting of morpholinyl, thiomorpholinyl, piperazinyl, piperidinyl, homomorpholinyl, homothiomorpholinyl, homothiomorpholinyl S-oxide, homothiomorpholinyl S,S-dioxide, pyrrolinyl and pyrrolidinyl where each group is optionally substituted with one, two, three, or four of C_1-C_6 alkyl,
- (16) $-(CH_2)_{0-4}-CO-O-R_{N-5}$ where R_{N-5} is selected from the group consisting of:
- (a) C_1-C_6 alkyl,
- (b) $-(CH_2)_{0-2}-(R_{1-aryl})$ where R_{1-aryl} is as defined above,
- (c) C_2-C_6 alkenyl containing one or two double bonds,
- (d) C_2-C_6 alkynyl containing one or two triple bonds,
- (e) C_3-C_7 cycloalkyl, and
- (f) $-(CH_2)_{0-2}-(R_{1-heteroaryl})$ where $R_{1-heteroaryl}$ is as defined above,
- (17) $-(CH_2)_{0-4}-SO_2-NR_{N-2}R_{N-3}$ where R_{N-2} and R_{N-3} are as defined above,
- (18) $-(CH_2)_{0-4}-SO-(C_1-C_8$ alkyl),
- (19) $-(CH_2)_{0-4}-SO_2-(C_1-C_{12}$ alkyl),
- (20) $-(CH_2)_{0-4}-SO_2-(C_3-C_7$ cycloalkyl),
- (21) $-(CH_2)_{0-4}-N(H \text{ or } R_{N-5})-CO-O-R_{N-5}$ where R_{N-5} can be the same or different and is as defined above,

- (22) $-(\text{CH}_2)_{0-4}-\text{N}(\text{H or } \text{R}_{\text{N}-5})-\text{CO}-\text{N}(\text{R}_{\text{N}-5})_2$, where $\text{R}_{\text{N}-5}$ can be the same or different and is as defined above,
- (23) $-(\text{CH}_2)_{0-4}-\text{N}-\text{CS}-\text{N}(\text{R}_{\text{N}-5})_2$, where $\text{R}_{\text{N}-5}$ can be the same or different and is as defined above,
- 5 (24) $-(\text{CH}_2)_{0-4}-\text{N}(\text{H or } \text{R}_{\text{N}-5})-\text{CO}-\text{R}_{\text{N}-2}$ where $\text{R}_{\text{N}-5}$ and $\text{R}_{\text{N}-2}$ can be the same or different and are as defined above,
- (25) $-(\text{CH}_2)_{0-4}-\text{NR}_{\text{N}-2}\text{R}_{\text{N}-3}$ where $\text{R}_{\text{N}-2}$ and $\text{R}_{\text{N}-3}$ can be the same or different and are as defined above,
- (26) $-(\text{CH}_2)_{0-4}-\text{R}_{\text{N}-4}$ where $\text{R}_{\text{N}-4}$ is as defined above,
- 10 (27) $-(\text{CH}_2)_{0-4}-\text{O}-\text{CO}-(\text{C}_1-\text{C}_6 \text{ alkyl})$,
- (28) $-(\text{CH}_2)_{0-4}-\text{O}-\text{P}(\text{O})-(\text{OR}_{\text{N-aryl-1}})_2$ where $\text{R}_{\text{N-aryl-1}}$ is $\text{H or C}_1-\text{C}_4 \text{ alkyl}$,
- (29) $-(\text{CH}_2)_{0-4}-\text{O}-\text{CO}-\text{N}(\text{R}_{\text{N}-5})_2$ where $\text{R}_{\text{N}-5}$ is as defined above,
- 15 (30) $-(\text{CH}_2)_{0-4}-\text{O}-\text{CS}-\text{N}(\text{R}_{\text{N}-5})_2$ where $\text{R}_{\text{N}-5}$ is as defined above,
- (31) $-(\text{CH}_2)_{0-4}-\text{O}-(\text{R}_{\text{N}-5})_2$ where $\text{R}_{\text{N}-5}$ is as defined above,
- (32) $-(\text{CH}_2)_{0-4}-\text{O}-(\text{R}_{\text{N}-5})_2-\text{COOH}$ where $\text{R}_{\text{N}-5}$ is as defined above,
- 20 (33) $-(\text{CH}_2)_{0-4}-\text{S}-(\text{R}_{\text{N}-5})_2$ where $\text{R}_{\text{N}-5}$ is as defined above,
- (34) $-(\text{CH}_2)_{0-4}-\text{O}-(\text{C}_1-\text{C}_6 \text{ alkyl optionally substituted with one, two, three, four, or five of } -\text{F})$,
- 25 (35) $\text{C}_3-\text{C}_7 \text{ cycloalkyl}$,
- (36) $\text{C}_2-\text{C}_6 \text{ alkenyl with one or two double bonds optionally substituted with } \text{C}_1-\text{C}_3 \text{ alkyl, } -\text{F, } -\text{Cl, } -\text{Br, } -\text{I, } -\text{OH, } -\text{SH, } -\text{C}\equiv\text{N, } -\text{CF}_3, \text{ C}_1-\text{C}_3 \text{ alkoxy, or } -\text{NR}_{1-a}\text{R}_{1-b} \text{ where } \text{R}_{1-a} \text{ and } \text{R}_{1-b} \text{ are as defined above,}$
- (37) $\text{C}_2-\text{C}_6 \text{ alkynyl with one or two triple bonds optionally substituted with } \text{C}_1-\text{C}_3 \text{ alkyl, } -\text{F, } -\text{Cl, } -\text{Br, } -\text{I, } -\text{OH, } -\text{SH, } -\text{C}\equiv\text{N, } -\text{CF}_3, \text{ C}_1-\text{C}_3 \text{ alkoxy, or } -\text{NR}_{1-a}\text{R}_{1-b} \text{ where } \text{R}_{1-a} \text{ and } \text{R}_{1-b} \text{ are as defined above,}$
- 30 (38) $-(\text{CH}_2)_{0-4}-\text{N}(\text{H or } \text{R}_{\text{N}-5})-\text{SO}_2-\text{R}_{\text{N}-2}$ where $\text{R}_{\text{N}-5}$ and $\text{R}_{\text{N}-2}$ can be the same or different and are as defined above, or

(39) $-(\text{CH}_2)_{0-4}-\text{C}_3-\text{C}_7$ cycloalkyl,

(C) $\text{R}_{\text{N-aryl}}-\text{W}-\text{R}_{\text{N-aryl}}$, where $\text{R}_{\text{N-aryl}}$ can be the same or different,

(D) $\text{R}_{\text{N-aryl}}-\text{W}-\text{R}_{\text{N-heteroaryl}}$,

5 (E) $\text{R}_{\text{N-aryl}}-\text{W}-\text{R}_{\text{N-1-heterocycle}}$, wherein $\text{R}_{\text{N-1-heterocycle}}$ is the same as $\text{R}_{\text{1-heterocycle}}$, and $\text{R}_{\text{1-heterocycle}}$ is as defined above

(F) $\text{R}_{\text{N-heteroaryl}}-\text{W}-\text{R}_{\text{N-aryl}}$,

(G) $\text{R}_{\text{N-heteroaryl}}-\text{W}-\text{R}_{\text{N-heteroaryl}}$,

(H) $\text{R}_{\text{N-heteroaryl}}-\text{W}-\text{R}_{\text{N-1-heterocycle}}$,

10 (I) $\text{R}_{\text{N-heterocycle}}-\text{W}-\text{R}_{\text{N-aryl}}$, wherein $\text{R}_{\text{N-heterocycle}}$ is the same as $\text{R}_{\text{1-heterocycle}}$, and $\text{R}_{\text{1-heterocycle}}$ is as defined above, and $\text{R}_{\text{N-aryl}}$ is as defined above,

(J) $\text{R}_{\text{N-heterocycle}}-\text{W}-\text{R}_{\text{N-heteroaryl}}$, and

(K) $\text{R}_{\text{N-heterocycle}}-\text{W}-\text{R}_{\text{N-1-heterocycle}}$,

where W is

15 (21) $-(\text{CH}_2)_{0-4}-$,

(22) $-\text{O}-$,

(23) $-\text{S}(\text{O})_{0-2}-$,

(24) $-\text{N}(\text{R}_{\text{N-5}})-$ where $\text{R}_{\text{N-5}}$ is as defined

above, or

20 (5) $-\text{CO}-$;

(II) $-\text{CO}-(\text{C}_1-\text{C}_{10} \text{ alkyl})$ where alkyl is optionally substituted with one three substituents selected from the group consisting of:

(A) $-\text{OH}$,

(B) $-\text{C}_1-\text{C}_6$ alkoxy,

25 (C) $-\text{C}_1-\text{C}_6$ thioalkoxy,

(D) $-\text{CO}-\text{O}-\text{R}_{\text{N-8}}$ where $\text{R}_{\text{N-8}}$ is $-\text{H}$, C_1-C_6 alkyl or $-\text{phenyl}$,

(E) $-\text{CO}-\text{NR}_{\text{N-2}}\text{R}_{\text{N-3}}$ where $\text{R}_{\text{N-2}}$ and $\text{R}_{\text{N-3}}$ are the same or

different and are as defined above,

(F) $-\text{CO}-\text{R}_{\text{N-4}}$ where $\text{R}_{\text{N-4}}$ is as defined above,

30 (G) $-\text{SO}_2-(\text{C}_1-\text{C}_8 \text{ alkyl})$,

(H) $-\text{SO}_2-\text{NR}_{\text{N-2}}\text{R}_{\text{N-3}}$ where $\text{R}_{\text{N-2}}$ and $\text{R}_{\text{N-3}}$ are the same or

different and are as defined above,

(I) $-\text{NH}-\text{CO}-(\text{C}_1-\text{C}_6 \text{ alkyl})$,

(J) $-\text{NH}-\text{CO}-\text{O}-\text{R}_{\text{N-8}}$ where $\text{R}_{\text{N-8}}$ is as defined above,

(K) $-NR_{N-2}R_{N-3}$ where R_{N-2} and R_{N-3} are the same or different and are as defined above,

(L) $-R_{N-4}$ where R_{N-4} is as defined above,

(M) $-O-CO-(C_1-C_6 \text{ alkyl})$,

5 (N) $-O-CO-NR_{N-8}R_{N-8}$ where R_{N-8} are the same or different and are as defined above,

(O) $-O-(C_1-C_5 \text{ alkyl})-COOH$,

(P) $-O-(C_1-C_6 \text{ alkyl})$ optionally substituted with one, two, or three of $-F$, $-Cl$, $-Br$, or $-I$,

10 (Q) $-NH-SO_2-(C_1-C_6 \text{ alkyl})$, and

(R) $-F$, or $-Cl$,

(III) $-CO-(C_1-C_6 \text{ alkyl})-O-(C_1-C_6 \text{ alkyl})$ where alkyl is optionally substituted with one, two, or three substituents selected from the group consisting of:

15 (A) $-OH$,

(B) $-C_1-C_6 \text{ alkoxy}$,

(C) $-C_1-C_6 \text{ thioalkoxy}$,

(D) $-CO-O-R_{N-8}$ where R_{N-8} is $-H$, $C_1-C_6 \text{ alkyl}$ or $-phenyl$,

(E) $-CO-NR_{N-2}R_{N-3}$ where R_{N-2} and R_{N-3} are the same or
20 different and are as defined above,

(F) $-CO-R_{N-4}$ where R_{N-4} is as defined above,

(G) $-SO_2-(C_1-C_8 \text{ alkyl})$,

(H) $-SO_2-NR_{N-2}R_{N-3}$ where R_{N-2} and R_{N-3} are the same or different and are as defined above,

25 (I) $-NH-CO-(C_1-C_6 \text{ alkyl})$,

(J) $-NH-CO-O-R_{N-8}$ where R_{N-8} is as defined above,

(K) $-NR_{N-2}R_{N-3}$ where R_{N-2} and R_{N-3} are the same or different and are as defined above,

(L) $-R_{N-4}$ where R_{N-4} is as defined above,

30 (M) $-O-CO-(C_1-C_6 \text{ alkyl})$,

(N) $-O-CO-NR_{N-8}R_{N-8}$ where R_{N-8} are the same or different and are as defined above,

(O) $-O-(C_1-C_5 \text{ alkyl})-COOH$,

(P) $-O-(C_1-C_6 \text{ alkyl optionally substituted with one, two, or three of } -F, -Cl, -Br, \text{ or } -I),$

(Q) $-NH-SO_2-(C_1-C_6 \text{ alkyl}),$ and

(R) $-F, \text{ or } -Cl,$

5 (IV) $-CO-(C_1-C_6 \text{ alkyl})-S-(C_1-C_6 \text{ alkyl})$ where alkyl is optionally substituted with one, two, or three of substituents selected from the group consisting of:

(A) $-OH,$

(B) $-C_1-C_6 \text{ alkoxy},$

10 (C) $-C_1-C_6 \text{ thioalkoxy},$

(D) $-CO-O-R_{N-8}$ where R_{N-8} is as defined above,

(E) $-CO-NR_{N-2}R_{N-3}$ where R_{N-2} and R_{N-3} are the same or different and are as defined above,

(F) $-CO-R_{N-4}$ where R_{N-4} is as defined above,

15 (G) $-SO_2-(C_1-C_8 \text{ alkyl}),$

(H) $-SO_2-NR_{N-2}R_{N-3}$ where R_{N-2} and R_{N-3} are the same or different and are as defined above,

(I) $-NH-CO-(C_1-C_6 \text{ alkyl}),$

(J) $-NH-CO-O-R_{N-8}$ where R_{N-8} is as defined above,

20 (K) $-NR_{N-2}R_{N-3}$ where R_{N-2} and R_{N-3} are the same or different and are as defined above,

(L) $-R_{N-4}$ where R_{N-4} is as defined above,

(M) $-O-CO-(C_1-C_6 \text{ alkyl}),$

(N) $-O-CO-NR_{N-8}R_{N-8}$ where R_{N-8} are the same or different

25 and are as defined above,

(O) $-O-(C_1-C_5 \text{ alkyl})-COOH,$

(P) $-O-(C_1-C_6 \text{ alkyl optionally substituted with one, two, or three of } -F, -Cl, -Br, \text{ or } -I),$

(Q) $-NH-SO_2-(C_1-C_6 \text{ alkyl}),$ and

30 (R) $-F, \text{ or } -Cl,$

(V) $-CO-CH(-(CH_2)_{0-2}-O-R_{N-10})-(CH_2)_{0-2}-R_{N-aryl}/R_{N-heteroaryl})$ where R_{N-aryl} and $R_{N-heteroaryl}$ are as defined above, where R_{N-10} is selected from the group consisting of:

- (A) -H,
 (B) C₁-C₆ alkyl,
 (C) C₃-C₇ cycloalkyl,
 (D) C₂-C₆ alkenyl with one double bond,
 5 (E) C₂-C₆ alkynyl with one triple bond,
 (F) R_{1-aryl} where R_{1-aryl} is as defined above, and
 (G) R_{N-heteroaryl} where R_{N-heteroaryl} is as defined above, or
 (VI) -CO-(C₃-C₈ cycloalkyl) where alkyl is optionally substituted
 with one or two substituents selected from the group consisting of:
- 10 (A) -(CH₂)₀₋₄-OH,
 (B) -(CH₂)₀₋₄-C₁-C₆ alkoxy,
 (C) -(CH₂)₀₋₄-C₁-C₆ thioalkoxy,
 (D) -(CH₂)₀₋₄-CO-O-R_{N-8} where R_{N-8} is -H, C₁-C₆ alkyl or
 phenyl,
 15 (E) -(CH₂)₀₋₄-CO-NR_{N-2}R_{N-3} where R_{N-2} and R_{N-3} are the
 same or different and are as defined above,
 (F) -(CH₂)₀₋₄-CO-R_{N-4} where R_{N-4} is as defined above,
 (G) -(CH₂)₀₋₄-SO₂-(C₁-C₈ alkyl),
 (H) -(CH₂)₀₋₄-SO₂-NR_{N-2}R_{N-3} where R_{N-2} and R_{N-3} are the
 20 same or different and are as defined above,
 (I) -(CH₂)₀₋₄-NH-CO-(C₁-C₆ alkyl),
 (J) -NH-CO-O-R_{N-8} where R_{N-8} is as defined above,
 (K) -(CH₂)₀₋₄-NR_{N-2}R_{N-3} where R_{N-2} and R_{N-3} are the same or
 different and are as defined above,
 25 (L) -(CH₂)₀₋₄-R_{N-4} where R_{N-4} is as defined above,
 (M) -O-CO-(C₁-C₆ alkyl),
 (N) -O-CO-NR_{N-8}R_{N-8} where R_{N-8} are the same or different
 and are as defined above,
 (O) -O-(C₁-C₅ alkyl)-COOH,
 30 (P) -O-(C₁-C₆ alkyl optionally substituted with one, two, or
 three of -F, -Cl, -Br, or -I),
 (Q) -NH-SO₂-(C₁-C₆ alkyl), and

(R) -F, or -Cl;

where R_A is:

(I) C_1-C_{10} alkyl optionally substituted with one, two or three
 5 substituents selected from the group consisting of C_1-C_3 alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C \equiv N, -CF₃, C_1-C_6 alkoxy, -O-phenyl, -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above, -OC=O NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above, -S(=O)₀₋₂ R_{1-a} where R_{1-a} is as defined above, -NR_{1-a}C=O NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above, -C=O NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above, and -
 10 S(=O)₂ NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(II) -(CH₂)₀₋₃-(C₃-C₈) cycloalkyl where cycloalkyl can be optionally substituted with one, two or three substituents selected from the group consisting of C_1-C_3 alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C \equiv N, -CF₃, C_1-C_6 alkoxy, -O-phenyl, -CO-OH, -CO-O-(C₁-C₄ alkyl), and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined
 15 above,

(III) -(CR_{A-x}R_{A-y})₀₋₄-R_{A-aryl} where R_{A-x} and R_{A-y} are

(A) -H,

(B) C_1-C_4 alkyl optionally substituted with one or two -OH,

(C) C_1-C_4 alkoxy optionally substituted with one, two, or

20 three of -F,

(D) -(CH₂)₀₋₄-C₃-C₇ cycloalkyl,

(E) C_2-C_6 alkenyl containing one or two double bonds,

(F) C_2-C_6 alkynyl containing one or two triple bonds, or

(G) phenyl,

25 and where R_{A-x} and R_{A-y} are taken together with the carbon to which they are attached to form a carbocycle of three, four, five, six or seven carbon atoms, optionally where one carbon atom is replaced by a heteroatom selected from the group consisting of -O-, -S-, -SO₂-, and -NR_{N-2}- and R_{A-aryl} is the same as R_{N-aryl},

(IV) -(CR_{A-x}R_{A-y})₀₋₄-R_{A-heteroaryl} where R_{A-heteroaryl} is the same as R_{N-heteroaryl} and R_{A-x} and R_{A-y} are as defined above,
 30

(V) -(CR_{A-x}R_{A-y})₀₋₄-R_{A-aryl}-R_{A-aryl} where R_{A-aryl}, R_{A-x} and R_{A-y} are as defined above,

(VI) $-(\text{CR}_{\text{A-x}}\text{R}_{\text{A-y}})_{0-4}\text{-R}_{\text{A-aryl}}\text{-R}_{\text{A-heteroaryl}}$ where $\text{R}_{\text{A-aryl}}$, $\text{R}_{\text{A-heteroaryl}}$, $\text{R}_{\text{A-x}}$ and $\text{R}_{\text{A-y}}$ are as defined above,

(VII) $-(\text{CR}_{\text{A-x}}\text{R}_{\text{A-y}})_{0-4}\text{-R}_{\text{A-heteroaryl}}\text{-R}_{\text{A-aryl}}$ where $\text{R}_{\text{A-heteroaryl}}$, $\text{R}_{\text{A-aryl}}$, $\text{R}_{\text{A-x}}$ and $\text{R}_{\text{A-y}}$ are as defined above,

5 (VIII) $-(\text{CR}_{\text{A-x}}\text{R}_{\text{A-y}})_{0-4}\text{-R}_{\text{A-heteroaryl}}\text{-R}_{\text{A-heteroaryl}}$ where $\text{R}_{\text{A-heteroaryl}}$, $\text{R}_{\text{A-x}}$ and $\text{R}_{\text{A-y}}$ are as defined above,

(IX) $-(\text{CR}_{\text{A-x}}\text{R}_{\text{A-y}})_{0-4}\text{-R}_{\text{A-aryl}}\text{-R}_{\text{A-heterocycle}}$ where $\text{R}_{\text{A-heterocycle}}$ is defined as $\text{R}_{1\text{-heterocycle}}$, and where $\text{R}_{\text{A-aryl}}$, $\text{R}_{\text{A-x}}$ and $\text{R}_{\text{A-y}}$ are as defined above,

(X) $-(\text{CR}_{\text{A-x}}\text{R}_{\text{A-y}})_{0-4}\text{-R}_{\text{A-heteroaryl}}\text{-R}_{\text{A-heterocycle}}$ where $\text{R}_{\text{A-heteroaryl}}$, $\text{R}_{\text{A-heterocycle}}$, $\text{R}_{\text{A-x}}$ and $\text{R}_{\text{A-y}}$ are as defined above,

(XI) $-(\text{CR}_{\text{A-x}}\text{R}_{\text{A-y}})_{0-4}\text{-R}_{\text{A-heterocycle}}\text{-R}_{\text{A-aryl}}$ where $\text{R}_{\text{A-heterocycle}}$, $\text{R}_{\text{A-aryl}}$, $\text{R}_{\text{A-x}}$ and $\text{R}_{\text{A-y}}$ are as defined above,

(XII) $-(\text{CR}_{\text{A-x}}\text{R}_{\text{A-y}})_{0-4}\text{-R}_{\text{A-heterocycle}}\text{-R}_{\text{A-heteroaryl}}$ where $\text{R}_{\text{A-heterocycle}}$, $\text{R}_{\text{A-heteroaryl}}$, $\text{R}_{\text{A-x}}$ and $\text{R}_{\text{A-y}}$ are as defined above,

15 (XIII) $-(\text{CR}_{\text{A-x}}\text{R}_{\text{A-y}})_{0-4}\text{-R}_{\text{A-heterocycle}}\text{-R}_{\text{A-heterocycle}}$ where $\text{R}_{\text{A-heterocycle}}$, $\text{R}_{\text{A-x}}$ and $\text{R}_{\text{A-y}}$ are as defined above,

(XIV) $-(\text{CR}_{\text{A-x}}\text{R}_{\text{A-y}})_{0-4}\text{-R}_{\text{A-heterocycle}}$ where $\text{R}_{\text{A-heterocycle}}$, $\text{R}_{\text{A-x}}$ and $\text{R}_{\text{A-y}}$ are as defined above,

(XV) $-\text{[C(R}_{\text{A-1}}\text{)(R}_{\text{A-2}}\text{)]}_{1-3}\text{-CO-N-(R}_{\text{A-3}}\text{)}_2$ where $\text{R}_{\text{A-1}}$ and $\text{R}_{\text{A-2}}$ are the same or different and are selected from the group consisting of:

(A) -H,

(B) $\text{-C}_1\text{-C}_6$ alkyl, optionally substituted with one, two or three substituents selected from the group consisting of $\text{C}_1\text{-C}_3$ alkyl, -F, -Cl, -Br, -I, -OH, -SH, $\text{-C}\equiv\text{N}$, -CF_3 , $\text{C}_1\text{-C}_6$ alkoxy, -O-phenyl, and $\text{-NR}_{1\text{-a}}\text{R}_{1\text{-b}}$ where $\text{R}_{1\text{-a}}$ and $\text{R}_{1\text{-b}}$ are as defined above,

(C) $\text{C}_2\text{-C}_6$ alkenyl with one or two double bonds, optionally substituted with one, two or three substituents selected from the group consisting of $\text{C}_1\text{-C}_3$ alkyl, -F, -Cl, -Br, -I, -OH, -SH, $\text{-C}\equiv\text{N}$, -CF_3 , $\text{C}_1\text{-C}_6$ alkoxy, -O-phenyl, and $\text{-NR}_{1\text{-a}}\text{R}_{1\text{-b}}$ where $\text{R}_{1\text{-a}}$ and $\text{R}_{1\text{-b}}$ are as defined above,

30 (D) $\text{C}_2\text{-C}_6$ alkynyl with one or two triple bonds, optionally substituted with one, two or three substituents selected from the group consisting of $\text{C}_1\text{-C}_3$ alkyl, -F, -Cl, -Br, -I, -OH, -SH, $\text{-C}\equiv\text{N}$, -CF_3 , $\text{C}_1\text{-C}_6$ alkoxy, -O-phenyl, and $\text{-NR}_{1\text{-a}}\text{R}_{1\text{-b}}$ where $\text{R}_{1\text{-a}}$ and $\text{R}_{1\text{-b}}$ are as defined above,

(E) $-(\text{CH}_2)_{1-2}-\text{S}(\text{O})_{0-2}-(\text{C}_1-\text{C}_6 \text{ alkyl})$,

(F) $-(\text{CH}_2)_{0-4}-\text{C}_3-\text{C}_7 \text{ cycloalkyl}$, optionally substituted with one, two or three substituents selected from the group consisting of $\text{C}_1-\text{C}_3 \text{ alkyl}$, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, $-\text{OH}$, $-\text{SH}$, $-\text{C}\equiv\text{N}$, $-\text{CF}_3$, $\text{C}_1-\text{C}_6 \text{ alkoxy}$, $-\text{O-phenyl}$, and $-\text{NR}_{1-a}\text{R}_{1-b}$ where

5 R_{1-a} and R_{1-b} are as defined above,

(G) $-(\text{C}_1-\text{C}_4 \text{ alkyl})-\text{R}_{\text{A}'\text{-aryl}}$ where $\text{R}_{\text{A}'\text{-aryl}}$ is as defined for R_{1-}

aryl,

(H) $-(\text{C}_1-\text{C}_4 \text{ alkyl})-\text{R}_{\text{A-heteroaryl}}$ where $\text{R}_{\text{A-heteroaryl}}$ is as defined

above,

10 (I) $-(\text{C}_1-\text{C}_4 \text{ alkyl})-\text{R}_{\text{A-heterocycle}}$ where $\text{R}_{\text{A-heterocycle}}$ is as defined above,

(J) $-\text{R}_{\text{A-heteroaryl}}$ where $\text{R}_{\text{A-heteroaryl}}$ is as defined above,

(K) $-\text{R}_{\text{A-heterocycle}}$ where $\text{R}_{\text{A-heterocycle}}$ is as defined above,

(M) $-(\text{CH}_2)_{1-4}-\text{R}_{\text{A-4}}-(\text{CH}_2)_{0-4}-\text{R}_{\text{A}'\text{-aryl}}$ where $\text{R}_{\text{A-4}}$ is $-\text{O}-$, $-\text{S}-$ or
15 $-\text{NR}_{\text{A-5}}-$ where $\text{R}_{\text{A-5}}$ is $\text{C}_1-\text{C}_6 \text{ alkyl}$, and where $\text{R}_{\text{A}'\text{-aryl}}$ is defined above,

(N) $-(\text{CH}_2)_{1-4}-\text{R}_{\text{A-4}}-(\text{CH}_2)_{0-4}-\text{R}_{\text{A-heteroaryl}}$ where $\text{R}_{\text{A-4}}$ and $\text{R}_{\text{A-heteroaryl}}$ are as defined above, and

(O) $-\text{R}_{\text{A}'\text{-aryl}}$ where $\text{R}_{\text{A}'\text{-aryl}}$ is as defined above,

and where $\text{R}_{\text{A-3}}$ is the same or different and is:

20 (A) $-\text{H}$,

(B) $\text{C}_1-\text{C}_6 \text{ alkyl}$ optionally substituted with one, two or three substituents selected from the group consisting of $\text{C}_1-\text{C}_3 \text{ alkyl}$, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, $-\text{OH}$, $-\text{SH}$, $-\text{C}\equiv\text{N}$, $-\text{CF}_3$, $\text{C}_1-\text{C}_6 \text{ alkoxy}$, $-\text{O-phenyl}$, $-\text{NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

25 (C) $\text{C}_2-\text{C}_6 \text{ alkenyl}$ with one or two double bonds, optionally substituted with one, two or three substituents selected from the group consisting of $\text{C}_1-\text{C}_3 \text{ alkyl}$, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, $-\text{OH}$, $-\text{SH}$, $-\text{C}\equiv\text{N}$, $-\text{CF}_3$, $\text{C}_1-\text{C}_6 \text{ alkoxy}$, $-\text{O-phenyl}$, and $-\text{NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

(D) $\text{C}_2-\text{C}_6 \text{ alkynyl}$ with one or two triple bonds, optionally
30 substituted with one, two or three substituents selected from the group consisting of $\text{C}_1-\text{C}_3 \text{ alkyl}$, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, $-\text{OH}$, $-\text{SH}$, $-\text{C}\equiv\text{N}$, $-\text{CF}_3$, $\text{C}_1-\text{C}_6 \text{ alkoxy}$, $-\text{O-phenyl}$, and $-\text{NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

(E) $-(\text{CH}_2)_{0-4}-\text{C}_3-\text{C}_7$ cycloalkyl, optionally substituted with one, two or three substituents selected from the group consisting of C_1-C_3 alkyl, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, $-\text{OH}$, $-\text{SH}$, $-\text{C}\equiv\text{N}$, $-\text{CF}_3$, C_1-C_6 alkoxy, $-\text{O}$ -phenyl, and $-\text{NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

5 (F) $-\text{R}_{\text{A}'\text{-aryl}}$ where $\text{R}_{\text{A}'\text{-aryl}}$ is as defined above,

(G) $-\text{R}_{\text{A-heteroaryl}}$ where $\text{R}_{\text{A-heteroaryl}}$ is as defined above,

(H) $-\text{R}_{\text{A-heterocycle}}$ where $\text{R}_{\text{A-heterocycle}}$ is as defined above,

(I) $-(\text{C}_1-\text{C}_4 \text{ alkyl})-\text{R}_{\text{A}'\text{-aryl}}$ where $\text{R}_{\text{A}'\text{-aryl}}$ is as defined above,

10 (J) $-(\text{C}_1-\text{C}_4 \text{ alkyl})-\text{R}_{\text{A-heteroaryl}}$ where $\text{R}_{\text{A-heteroaryl}}$ is as defined above,

(K) $-(\text{C}_1-\text{C}_4 \text{ alkyl})-\text{R}_{\text{A-heterocycle}}$ where $\text{R}_{\text{A-heterocycle}}$ is as defined above, or

(XVI) $-\text{CH}(\text{R}_{\text{A-aryl}})_2$ where $\text{R}_{\text{A-aryl}}$ are the same or different and are as defined above,

(XVII) $-\text{CH}(\text{R}_{\text{A-heteroaryl}})_2$ where $\text{R}_{\text{A-heteroaryl}}$ are the same or different and are as defined above,

(XVIII) $-\text{CH}(\text{R}_{\text{A-aryl}})(\text{R}_{\text{A-heteroaryl}})$ where $\text{R}_{\text{A-aryl}}$ and $\text{R}_{\text{A-heteroaryl}}$ are as defined above,

20 (XIX) -cyclopentyl, -cyclohexyl, or -cycloheptyl ring fused to $\text{R}_{\text{A-aryl}}$, $\text{R}_{\text{A-heteroaryl}}$, $\text{R}_{\text{A-heterocycle}}$ where $\text{R}_{\text{A-aryl}}$ or $\text{R}_{\text{A-heteroaryl}}$ or $\text{R}_{\text{A-heterocycle}}$ are as defined above where one carbon of cyclopentyl, cyclohexyl, or -cycloheptyl is optionally replaced with NH , $\text{NR}_{\text{N-5}}$, O , or $\text{S}(=\text{O})_{0-2}$, and where cyclopentyl, cyclohexyl, or -cycloheptyl can be optionally substituted with one or two $-\text{C}_1-\text{C}_3$ alkyl, $-\text{F}$, $-\text{OH}$, $-\text{SH}$, $-\text{C}\equiv\text{N}$, $-\text{CF}_3$, C_1-C_6 alkoxy, $=\text{O}$, or $-\text{NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

(XX) C_2-C_{10} alkenyl containing one or two double bonds optionally substituted with one, two or three substituents selected from the group consisting of C_1-C_3 alkyl, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, $-\text{OH}$, $-\text{SH}$, $-\text{C}\equiv\text{N}$, $-\text{CF}_3$, C_1-C_6 alkoxy, $-\text{O}$ -phenyl, and $-\text{NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

30 (XXI) C_2-C_{10} alkynyl containing one or two triple bonds optionally substituted with one, two or three substituents selected from the group consisting of C_1-C_3 alkyl, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, $-\text{OH}$, $-\text{SH}$, $-\text{C}\equiv\text{N}$, $-\text{CF}_3$, C_1-C_6 alkoxy, $-\text{O}$ -phenyl, $-\text{NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

(XXI) $-(\text{CH}_2)_{0-1}-\text{CHR}_{\text{A-6}}-(\text{CH}_2)_{0-1}-\text{R}_{\text{A-aryl}}$ where $\text{R}_{\text{A-aryl}}$ is as defined above and $\text{R}_{\text{A-6}}$ is $-(\text{CH}_2)_{0-6}-\text{OH}$,

(XXII) $-(\text{CH}_2)_{0-1}-\text{CHR}_{\text{A-6}}-(\text{CH}_2)_{0-1}-\text{R}_{\text{A-heteroaryl}}$ where $\text{R}_{\text{A-heteroaryl}}$ and $\text{R}_{\text{A-6}}$ is as defined above,

5 (XXIII) $-\text{CH}(-\text{R}_{\text{A-aryl}}$ or $\text{R}_{\text{A-heteroaryl}})-\text{CO}-\text{O}(\text{C}_1-\text{C}_4 \text{ alkyl})$ where $\text{R}_{\text{A-aryl}}$ and $\text{R}_{\text{A-heteroaryl}}$ are as defined above,

(XXIV) $-\text{CH}(-\text{CH}_2-\text{OH})-\text{CH}(-\text{OH})-\text{micro-NO}_2$,

(XXV) $(\text{C}_1-\text{C}_6 \text{ alkyl})-\text{O}-(\text{C}_1-\text{C}_6 \text{ alkyl})-\text{OH}$,

(XXVII) $-\text{CH}_2-\text{NH}-\text{CH}_2-\text{CH}(-\text{O}-\text{CH}_2-\text{CH}_3)_2$,

10 (XXVIII) $-\text{H}$,

(XXIX) $-(\text{CH}_2)_{0-6}-\text{C}(=\text{NR}_{1-a})(\text{NR}_{1-a}\text{R}_{1-b})$ where R_{1-a} and R_{1-b} are as defined above; or

(XXX)

15 $-\text{C}=\text{OC}(\text{HR}_6)\text{NHR}_7$, where R_6 and R_7 are as defined below,
 $-\text{C}=\text{OR}_7$, where R_7 is as defined below,
 $-\text{C}=\text{OOR}_7$, where R_7 is as defined below, or
 $-\text{SOOR}_7$ where R_7 is as defined below,

wherein R_6 is:

20 hydrogen,
 $\text{C}_1 - \text{C}_3 \text{ alkyl}$,
phenyl,
thioalkoxyalkyl,
alkyl substituted aryl,
cycloalkyl,
25 cycloalkylalkyl,
hydroxyalkyl,
alkoxyalkyl,
aryloxyalkyl,
haloalkyl,
30 carboxyalkyl,
alkoxycarbonylalkyl,
aminoalkyl,
(N-protected)aminoalkyl,
alkylaminoalkyl,

5 ((N-protected)(alkyl)amino)alkyl,
dialkylaminoalkyl,
guanidinoalkyl,
lower alkenyl,
heterocyclic,
(heterocyclic)alkyl),
arylthioalkyl,
arylsulfonylalkyl,
(heterocyclic)thioalkyl,
10 (heterocyclic)sulfonylalkyl,
(heterocyclic)oxyalkyl,
arylalkoxyalkyl,
arylthioalkoxyalkyl,
arylalkylsulfonylalkyl,
15 (heterocyclic))alkoxyalkyl,
(heterocyclic)thioalkoxyalkyl,
(heterocyclic)alkylsulfonylalkyl,
cycloalkyloxyalkyl,
cycloalkylthioalkyl,
20 cycloalkylsulfonylalkyl,
cycloalkylalkoxyalkyl,
cycloalkylthioalkoxyalkyl,
cycloalkylalkylsulfonylalkyl,
aminocarbonyl,
25 alkylaminocarbonyl,
dialkylaminocarbonyl,
aroalkyl,
(heterocyclic)carbonylalkyl,
polyhydroxyalkyl,
30 aminocarbonylalkyl,
alkylaminocarbonylalkyl,
dialkylaminocarbonylalkyl,
aryloxyalkyl, or
alkylsulfonylalkyl,

wherein heterocyclic is pyridyl, thiazolyl, isothiazolyl, oxazolyl, isoxazolyl, furanyl, thienyl, tetrahydrofuranyl, tetrahydrothienyl and tetrahydro[2H]pyranyl and wherein the heterocycle is unsubstituted or substituted with one to three substituents independently selected from hydroxy, halo, amino, alkylamino, dialkylamino, alkoxy, polyalkoxy, haloalkyl, cycloalkyl, 5 cycloalkylalkyl, aryl, arylalkyl, COOH, -SO₃H, lower alkenyl or lower alkyl;


wherein R₇ is:

C₁ - C₃ alkyl,
phenyl,
10 thioalkoxyalkyl,
(aryl)alkyl,
cycloalkyl,
cycloalkylalkyl,
hydroxyalkyl,
15 alkoxyalkyl,
aryloxyalkyl,
haloalkyl,
carboxyalkyl,
alkoxycarbonylalkyl,
20 aminoalkyl,
(N-protected)aminocalkyl,
alkylaminoalkyl,
((N-protected)(alkyl)amino)alkyl,
dialkylaminoalkyl,
25 guanidinoalkyl,
lower alkenyl,
heterocyclic,
(heterocyclic)alkyl),
arylthioalkyl,
30 arylsulfonyalkyl,
(heterocyclic)thioalkyl,
(heterocyclic)sulfonylalkyl,
(heterocyclic)oxyalkyl,
arylalkoxyalkyl,


arylthioalkoxyalkyl,
 arylalkylsulfonylalkyl,
 (heterocyclic)alkoxyalkyl,
 (heterocyclic)thioalkoxyalkyl,
 5 (heterocyclic)alkylsulfonylalkyl,
 cycloalkyloxyalkyl,
 cycloalkylthioalkyl,
 cycloalkylsulfonylalkyl,
 cycloalkylalkoxyalkyl,
 10 cycloalkylthioalkoxyalkyl,
 cycloalkylalkylsulfonylalkyl,
 aminocarbonyl,
 alkylaminocarbonyl,
 dialkylaminocarbonyl,
 15 aroylalkyl,
 (heterocyclic)carbonylalkyl,
 polyhydroxyalkyl,
 aminocarbonylalkyl,
 alkylaminocarbonylalkyl,
 20 dialkylaminocarbonylalkyl,
 aryloxyalkyl, or
 alkylsulfonylalkyl,

wherein heterocyclic is pyridyl, thiazolyl, isothiazolyl,
 oxazolyl, isoxazolyl, furanyl, thienyl, tetrahydrofuranyl, tetrahydrothienyl and
 25 tetrahydro[2H]pyranyl and wherein the heterocycle is unsubstituted or substituted
 with one to three substituents independently selected from hydroxy, halo, amino,
 alkylamino, dialkylamino, alkoxy, polyalkoxy, haloalkyl, cycloalkyl,
 cycloalkylalkyl, aryl, arylalkyl, COOH, -SO₃H, lower alkenyl or lower alkyl.

30 52 A compound according to claim 51 where R₁ is:



where R_A is:



- 5 $-(\text{CH}_2)_{0-3}-(\text{C}_3-\text{C}_7)$ cycloalkyl,
 $-(\text{CR}_{\text{A}-x}\text{R}_{\text{A}-y})_{0-4}-\text{R}_{\text{A-aryl}}$,
 $-(\text{CR}_{\text{A}-x}\text{R}_{\text{A}-y})_{0-4}-\text{R}_{\text{A-heteroaryl}}$,
 $-(\text{CR}_{\text{A}-x}\text{R}_{\text{A}-y})_{0-4}-\text{R}_{\text{A-heterocycle}}$,
 -cyclopentyl or -cyclohexyl ring fused to $\text{R}_{\text{A-aryl}}$ or $\text{R}_{\text{A-heteroaryl}}$ or $\text{R}_{\text{A-}}$
 heterocycle.

53. A protected compound according to claim 51

where R_1 is:

- 10 $-(\text{CH}_2)-(\text{R}_{1\text{-aryl}})$, or
 $-(\text{CH}_2)-(\text{R}_{1\text{-heteroaryl}})$;

where R_2 is $-\text{H}$;


where R_3 is $-\text{H}$;

where R_{A} is:

- 15 $-\text{C}_1-\text{C}_8$ alkyl,
 $-(\text{CH}_2)_{0-3}-(\text{C}_3-\text{C}_7)$ cycloalkyl,
 $-(\text{CR}_{\text{A}-x}\text{R}_{\text{A}-y})_{0-4}-\text{R}_{\text{A-aryl}}$,
 $-(\text{CR}_{\text{A}-x}\text{R}_{\text{A}-y})_{0-4}-\text{R}_{\text{A-heteroaryl}}$,
 $-(\text{CR}_{\text{A}-x}\text{R}_{\text{A}-y})_{0-4}-\text{R}_{\text{A-heterocycle}}$,
 20 -cyclopentyl or -cyclohexyl ring fused to $\text{R}_{\text{A-aryl}}$ or $\text{R}_{\text{A-heteroaryl}}$ or $\text{R}_{\text{A-}}$
 heterocycle.

54. A method of treating a patient who has, or in preventing a patient from getting,
 a disease or condition selected from the group consisting of Alzheimer's disease, for
 25 helping prevent or delay the onset of Alzheimer's disease, for treating patients with
 mild cognitive impairment (MCI) and preventing or delaying the onset of
 Alzheimer's disease in those who would progress from MCI to AD, for treating
 Down's syndrome, for treating humans who have Hereditary Cerebral Hemorrhage
 with Amyloidosis of the Dutch-Type, for treating cerebral amyloid angiopathy and
 30 preventing its potential consequences, i.e. single and recurrent lobar hemorrhages,
 for treating other degenerative dementias, including dementias of mixed vascular
 and degenerative origin, dementia associated with Parkinson's disease, dementia
 associated with progressive supranuclear palsy, dementia associated with cortical
 basal degeneration, diffuse Lewy body type of Alzheimer's disease and who is in

need of such treatment which comprises administration of a therapeutically effective amount of a compound selected from the group consisting of a substituted amine of formula (XV)



5

where R_1 is:

- (I) C_1 - C_6 alkyl, optionally substituted with one, two or three substituents selected from the group consisting of C_1 - C_3 alkyl, C_1 - C_7 alkyl (optionally substituted with C_1 - C_3 alkyl and C_1 - C_3 alkoxy), -F, -Cl, -Br, -I, -OH, -SH, - $C\equiv N$, - CF_3 , C_1 - C_3 alkoxy, - $NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are -H or C_1 - C_6 alkyl, and - $OC=O$ $NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

(II) $-CH_2-S(O)_{0-2}-(C_1-C_6 \text{ alkyl})$,

(III) $-CH_2-CH_2-S(O)_{0-2}-(C_1-C_6 \text{ alkyl})$,

- (IV) C_2 - C_6 alkenyl with one or two double bonds, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, - $C\equiv N$, - CF_3 , C_1 - C_3 alkoxy, and - $NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are -H or C_1 - C_6 alkyl,

- (V) C_2 - C_6 alkynyl with one or two triple bonds, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, - $C\equiv N$, - CF_3 , C_1 - C_3 alkoxy, and - $NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are -H or C_1 - C_6 alkyl,

- (VI) $-(CH_2)_{n1}-(R_{1-aryl})$ where n_1 is zero or one and where R_{1-aryl} is phenyl, 1-naphthyl, 2-naphthyl and indanyl, indenyl, dihydronaphthalyl, or tetralinyl optionally substituted with one, two, three or four of the following substituents on the aryl ring:

(A) C_1 - C_6 alkyl optionally substituted with one, two or three substituents selected from the group consisting of C_1 - C_3 alkyl, -F, -Cl, -Br, -I, -OH, -SH, - $C\equiv N$, - CF_3 , and C_1 - C_3 alkoxy, and - $NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

- (B) C₂-C₆ alkenyl with one or two double bonds, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, -C≡N, -CF₃, C₁-C₃ alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are -H or C₁-C₆ alkyl,
- 5 (C) C₂-C₆ alkynyl with one or two triple bonds, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, -C≡N, -CF₃, C₁-C₃ alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are -H or C₁-C₆ alkyl,
- (D) -F, Cl, -Br or -I,
- 10 (F) -C₁-C₆ alkoxy optionally substituted with one, two or three of -F,
- (G) -NR_{N-2}R_{N-3} where R_{N-2} and R_{N-3} are as defined below,
- (H) -OH,
- (I) -C≡N,
- 15 (J) C₃-C₇ cycloalkyl, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, -C≡N, -CF₃, C₁-C₃ alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are -H or C₁-C₆ alkyl,
- (K) -CO-(C₁-C₄ alkyl),
- (L) -SO₂-NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,
- 20 (M) -CO-NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,
- or
- (N) -SO₂-(C₁-C₄ alkyl),
- (VII) -(CH₂)_{n1}-(R_{1-heteroaryl}) where n₁ is as defined above and where R_{1-heteroaryl} is selected from the group consisting of:
- 25 pyridinyl,
- pyrimidinyl,
- quinolinyl,
- benzothienyl,
- indolyl,
- 30 indolinyl,
- pyridazinyl,
- pyrazinyl,
- isoquinolyl,

	quinazoliny1,
	quinoxaliny1,
	phthalaziny1,
	imidazolyl,
5	isoxazolyl,
	pyrazolyl,
	oxazolyl,
	thiazolyl,
	indoliziny1,
10	indazolyl,
	benzothiazolyl,
	benzimidazolyl,
	benzofuranyl,
	furanyl,
15	thienyl,
	pyrrolyl,
	oxadiazolyl,
	thiadiazolyl,
	triazolyl,
20	tetrazolyl,
	oxazolopyridiny1,
	imidazopyridiny1,
	isothiazolyl,
	naphthyridiny1,
25	cinnoliny1,
	carbazolyl,
	beta-carboliny1,
	isochromanyl,
	chromanyl,
30	tetrahydroisoquinoliny1,
	isoindoliny1,
	isobenzotetrahydrofuranyl,
	isobenzotetrahydrothienyl,
	isobenzothieryl,

5 benzoxazolyl,
 pyridopyridinyl,
 benzotetrahydrofuranyl,
 benzotetrahydrothienyl,
 purinyl,
 benzodioxolyl,
 triazinyl,
 phenoxazinyl,
 phenothiazinyl,
10 pteridinyl,
 benzothiazolyl,
 imidazopyridinyl,
 imidazothiazolyl,
 dihydrobenzisoaxazinyl,
15 benzisoaxazinyl,
 benzoxazinyl,
 dihydrobenzisothiazinyl,
 benzopyranyl,
 benzothiopyranyl,
20 coumarinyl,
 isocoumarinyl,
 chromonyl,
 chromanonyl,
 pyridinyl-N-oxide,
25 tetrahydroquinolinyl
 dihydroquinolinyl
 dihydroquinolinonyl
 dihydroisoquinolinonyl
 dihydrocoumarinyl
30 dihydroisocoumarinyl
 isoindolinonyl
 benzodioxanyl
 benzoxazolinonyl
 pyrrolyl N-oxide,

5 pyrimidinyl N-oxide,
 pyridazinyl N-oxide,
 pyrazinyl N-oxide,
 quinolinyl N-oxide,
 indolyl N-oxide,
 indolinyln N-oxide,
 isoquinolyl N-oxide,
 quinazolinyl N-oxide,
 quinoxalinyln N-oxide,
 10 phthalazinyl N-oxide,
 imidazolyl N-oxide,
 isoxazolyl N-oxide,
 oxazolyl N-oxide,
 thiazolyl N-oxide,
 15 indolizinyln N-oxide,
 indazolyl N-oxide,
 benzothiazolyl N-oxide,
 benzimidazolyl N-oxide,
 pyrrolyl N-oxide,
 20 oxadiazolyl N-oxide,
 thiadiazolyl N-oxide,
 triazolyl N-oxide,
 tetrazolyl N-oxide,
 benzothiopyranyl S-oxide, and
 25 benzothiopyranyl S,S-dioxide,

where the $R_{1\text{-heteroaryl}}$ group is bonded to $-(CH_2)_{n1}-$ by any ring atom
 of the parent $R_{1\text{-heteroaryl}}$ group substituted by hydrogen such that the new bond to the
 $R_{1\text{-heteroaryl}}$ group replaces the hydrogen atom and its bond, where heteroaryl is
 optionally substituted with one, two, three or four of:

30 (1) C_1 - C_6 alkyl optionally substituted with one, two or
 three substituents selected from the group consisting of C_1 - C_3 alkyl, -F, -Cl, -Br, -I,
 -OH, -SH, $-C\equiv N$, $-CF_3$, C_1 - C_3 alkoxy, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as
 defined above,

- (2) C₂-C₆ alkenyl with one or two double bonds, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, -C≡N, -CF₃, C₁-C₃ alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are -H or C₁-C₆ alkyl,
- 5 (3) C₂-C₆ alkynyl with one or two triple bonds, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, -C≡N, -CF₃, C₁-C₃ alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are -H or C₁-C₆ alkyl,
- (4) -F, Cl, -Br or -I,
- 10 (6) -C₁-C₆ alkoxy optionally substituted with one, two, or three of -F,
- (7) -NR_{N-2}R_{N-3} where R_{N-2} and R_{N-3} are as defined below,
- (8) -OH,
- 15 (9) -C≡N,
- (10) C₃-C₇ cycloalkyl, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, -C≡N, -CF₃, C₁-C₃ alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are -H or C₁-C₆ alkyl,
- (11) -CO-(C₁-C₄ alkyl),
- 20 (12) -SO₂-NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,
- (13) -CO-NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above, or
- (14) -SO₂-(C₁-C₄ alkyl), with the proviso that when n₁
- 25 is zero R_{1-heteroaryl} is not bonded to the carbon chain by nitrogen; or
- (VIII) -(CH₂)_{n1}-(R_{1-heterocycle}) where n₁ is as defined above and R_{1-heterocycle} is selected from the group consisting of:
- morpholinyl,
- thiomorpholinyl,
- 30 thiomorpholinyl S-oxide,
- thiomorpholinyl S,S-dioxide,
- piperazinyl,
- homopiperazinyl,

- pyrrolidinyl,
 pyrrolinyl,
 tetrahydropyranyl,
 piperidinyl,
 5 tetrahydrofuranyl,
 tetrahydrothienyl,
 homopiperidinyl,
 homomorpholinyl,
 homothiomorpholinyl,
 10 homothiomorpholinyl S,S-dioxide,
 oxazolidinonyl,
 dihydropyrazolyl,
 dihydropyrrolyl,
 dihydropyrazinyl,
 15 dihydropyridinyl,
 dihydropyrimidinyl,
 dihydrofuryl,
 dihydropyranyl,
 tetrahydrothienyl S-oxide,
 20 tetrahydrothienyl S,S-dioxide, and
 homothiomorpholinyl S-oxide,

where the R_{1-heterocycle} group is bonded by any atom of the parent R₁.

- heterocycle group substituted by hydrogen such that the new bond to the R_{1-heterocycle}
 group replaces the hydrogen atom and its bond, where heterocycle is optionally
 25 substituted with one, two, three or four:

- (1) C₁-C₆ alkyl optionally substituted with one, two or
 three substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I,
 -OH, -SH, -C≡N, -CF₃, C₁-C₃ alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as
 defined above,
 30 (2) C₂-C₆ alkenyl with one or two double bonds,
 optionally substituted with one, two or three substituents selected from the group
 consisting of -F, -Cl, -OH, -SH, -C≡N, -CF₃, C₁-C₃ alkoxy, and -NR_{1-a}R_{1-b} where R₁₋
a and R{1-b} are -H or C₁-C₆ alkyl,

- (3) C₂-C₆ alkynyl with one or two triple bonds, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, -C≡N, -CF₃, C₁-C₃ alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are -H or C₁-C₆ alkyl,
- 5 (4) -F, Cl, -Br or -I,
 (5) C₁-C₆ alkoxy,
 (6) -C₁-C₆ alkoxy optionally substituted with one, two, or three -F,
 (7) -NR_{N-2}R_{N-3} where R_{N-2} and R_{N-3} are as defined
- 10 below,
 (8) -OH,
 (9) -C≡N,
 (10) C₃-C₇ cycloalkyl, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, -
- 15 C≡N, -CF₃, C₁-C₃ alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are -H or C₁-C₆ alkyl,
 (11) -CO-(C₁-C₄ alkyl),
 (12) -SO₂-NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,
 (13) -CO-NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined
- 20 above,
 (14) -SO₂-(C₁-C₄ alkyl), or
 (15) =O, with the proviso that when n₁ is zero R_{1-heterocycle} is not bonded to the carbon chain by nitrogen;
- 25 where R₂ is:
 (I)-H,
 (II) C₁-C₆ alkyl, optionally substituted with one, two or three substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₃ alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined
- 30 above,
 (III) -(CH₂)₀₋₄-R₂₋₁ where R₂₋₁ is R_{1-aryl} or R_{1-heteroaryl} where R_{1-aryl} and R_{1-heteroaryl} are as defined above;

(IV) C₂-C₆ alkenyl with one or two double bonds, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, -C≡N, -CF₃, C₁-C₃ alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are -H or C₁-C₆ alkyl,

5 (V) C₂-C₆ alkynyl with one or two triple bonds, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, -C≡N, -CF₃, C₁-C₃ alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are -H or C₁-C₆ alkyl, or

10 (VI) -(CH₂)₀₋₄- C₃-C₇ cycloalkyl, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, -C≡N, -CF₃, C₁-C₃ alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are -H or C₁-C₆ alkyl;

where R₃ is:

(I)-H,

15 (II) C₁-C₆ alkyl, optionally substituted with one, two or three substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₃ alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

20 (III) -(CH₂)₀₋₄-R₂₋₁ where R₂₋₁ is R_{1-aryl} or R_{1-heteroaryl} where R_{1-aryl} and R_{1-heteroaryl} are as defined above

(IV) C₂-C₆ alkenyl with one or two double bonds,

(V) C₂-C₆ alkynyl with one or two triple bonds; or

25 (VI) -(CH₂)₀₋₄- C₃-C₇ cycloalkyl, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, -C≡N, -CF₃, C₁-C₃ alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are -H or C₁-C₆ alkyl,

and where R₂ and R₃ are taken together with the carbon to which they are attached to form a carbocycle of three, four, five, six, or seven carbon atoms, optionally where one carbon atom is replaced by a heteroatom selected from the group consisting of -O-, -S-, -SO₂-, and -NR_{N-2}-, where R_{N-2} is selected from the group consisting of:

(a) -H,

(b) -C₁-C₆ alkyl optionally substituted with one substituent selected from the group consisting of:

350

(i) -OH, and

(ii) -NH₂,(c) -C₁-C₆ alkyl optionally substituted with

one, two, or three -F, -Cl, -Br, or -I,

5

(d) -C₃-C₇ cycloalkyl,(e) -(C₁-C₂ alkyl)-(C₃-C₇ cycloalkyl),(f) -(C₁-C₆ alkyl)-O-(C₁-C₃ alkyl),(g) -C₂-C₆ alkenyl with one or two double

bonds,

10

(h) -C₂-C₆ alkynyl with one or two triple

bonds,

(i) -C₁-C₆ alkyl chain with one double bond

and one triple bond,

(j) -R_{1-aryl} where R_{1-aryl} is as defined above, and

15

(k) -R_{1-heteroaryl} where R_{1-heteroaryl} is as defined

above;

where R_N is:(I) R_{N-1}-X_N- where X_N is selected from the group consisting of:

20

(A) -CO-,

(B) -SO₂-,(C) -(CR'R'')₁₋₆ where R' and R'' are the same or different andare -H and C₁-C₄ alkyl,(D) -CO-(CR'R'')₁₋₆-X_{N-1} where X_{N-1} is selected from the

25

group consisting of -O-, -S- and -NR'- and where R' and R'' are as defined above,

and

(E) a single bond;

where R_{N-1} is selected from the group consisting of:(A) R_{N-aryl} where R_{N-aryl} is phenyl, 1-naphthyl, 2-naphthyl,

30

tetralinyl, indanyl, 6,7,8,9-tetrahydro-5H-benzo[a]cycloheptenyl, or dihydronaphthyl optionally substituted with one, two or three of the following substituents which can be the same or different and are:

(1) C₁-C₆ alkyl, optionally substituted with one, two orthree substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I,

-OH, -SH, $-C\equiv N$, $-CF_3$, C_1-C_3 alkoxy, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

(2) -OH,

(3) $-NO_2$,

5 (4) -F, -Cl, -Br, -I,

(5) -CO-OH,

(6) $-C\equiv N$,

(7) $-(CH_2)_{0-4}-CO-NR_{N-2}R_{N-3}$ where R_{N-2} and R_{N-3} are

the same or different and are selected from the group consisting of:

10 (a) -H,

(b) $-C_1-C_6$ alkyl optionally substituted with one substituent selected from the group consisting of:

(i) -OH, and

(ii) $-NH_2$,

15 (c) $-C_1-C_6$ alkyl optionally substituted with one, two, or three -F, -Cl, -Br, or -I,

(d) $-C_3-C_7$ cycloalkyl,

(e) $-(C_1-C_2 \text{ alkyl})-(C_3-C_7 \text{ cycloalkyl})$,

(f) $-(C_1-C_6 \text{ alkyl})-O-(C_1-C_3 \text{ alkyl})$,

20 (g) $-C_2-C_6$ alkenyl with one or two double bonds,

(h) $-C_2-C_6$ alkynyl with one or two triple

bonds,

(i) $-C_1-C_6$ alkyl chain with one double bond

25 and one triple bond,

(j) $-R_{1-aryl}$ where R_{1-aryl} is as defined above, and

(k) $-R_{1-heteroaryl}$ where $R_{1-heteroaryl}$ is as defined

above,

(8) $-(CH_2)_{0-4}-CO-(C_1-C_{12} \text{ alkyl})$,

30 (9) $-(CH_2)_{0-4}-CO-(C_2-C_{12} \text{ alkenyl with one, two or three double bonds})$,

(10) $-(CH_2)_{0-4}-CO-(C_2-C_{12} \text{ alkynyl with one, two or$

three triple bonds),

(11) $-(\text{CH}_2)_{0-4}-\text{CO}-(\text{C}_3-\text{C}_7 \text{ cycloalkyl})$,

(12) $-(\text{CH}_2)_{0-4}-\text{CO}-\text{R}_{1-\text{aryl}}$ where $\text{R}_{1-\text{aryl}}$ is as defined

above,

(13) $-(\text{CH}_2)_{0-4}-\text{CO}-\text{R}_{1-\text{heteroaryl}}$ where $\text{R}_{1-\text{heteroaryl}}$ is as

5 defined above,

(14) $-(\text{CH}_2)_{0-4}-\text{CO}-\text{R}_{1-\text{heterocycle}}$ where $\text{R}_{1-\text{heterocycle}}$ is as

defined above,

(15) $-(\text{CH}_2)_{0-4}-\text{CO}-\text{R}_{\text{N}-4}$ where $\text{R}_{\text{N}-4}$ is selected from

the group consisting of morpholinyl, thiomorpholinyl, piperazinyl, piperidinyl,

10 homomorpholinyl, homothiomorpholinyl, homothiomorpholinyl S-oxide,

homothiomorpholinyl S,S-dioxide, pyrrolinyl and pyrrolidinyl where each group is

optionally substituted with one, two, three, or four of C_1-C_6 alkyl,

(16) $-(\text{CH}_2)_{0-4}-\text{CO}-\text{O}-\text{R}_{\text{N}-5}$ where $\text{R}_{\text{N}-5}$ is

selected from the group consisting of:

15

(a) C_1-C_6 alkyl,

(b) $-(\text{CH}_2)_{0-2}-(\text{R}_{1-\text{aryl}})$ where $\text{R}_{1-\text{aryl}}$ is as defined

above,

(c) C_2-C_6 alkenyl containing one or two double

bonds,

20

(d) C_2-C_6 alkynyl containing one or two triple

bonds,

(e) C_3-C_7 cycloalkyl, and

(f) $-(\text{CH}_2)_{0-2}-(\text{R}_{1-\text{heteroaryl}})$ where $\text{R}_{1-\text{heteroaryl}}$ is as

defined above,

25

(17) $-(\text{CH}_2)_{0-4}-\text{SO}_2-\text{NR}_{\text{N}-2}\text{R}_{\text{N}-3}$ where $\text{R}_{\text{N}-2}$ and $\text{R}_{\text{N}-3}$ are

as defined above,

(18) $-(\text{CH}_2)_{0-4}-\text{SO}-(\text{C}_1-\text{C}_8 \text{ alkyl})$,

(19) $-(\text{CH}_2)_{0-4}-\text{SO}_2-(\text{C}_1-\text{C}_{12} \text{ alkyl})$,

(20) $-(\text{CH}_2)_{0-4}-\text{SO}_2-(\text{C}_3-\text{C}_7 \text{ cycloalkyl})$,

30

(21) $-(\text{CH}_2)_{0-4}-\text{N}(\text{H or } \text{R}_{\text{N}-5})-\text{CO}-\text{O}-\text{R}_{\text{N}-5}$ where $\text{R}_{\text{N}-5}$

can be the same or different and is as defined above,

(22) $-(\text{CH}_2)_{0-4}-\text{N}(\text{H or } \text{R}_{\text{N}-5})-\text{CO}-\text{N}(\text{R}_{\text{N}-5})_2$, where $\text{R}_{\text{N}-5}$

can be the same or different and is as defined above,

(23) $-(CH_2)_{0-4}-N-CS-N(R_{N-5})_2$, where R_{N-5} can be the same or different and is as defined above,

(24) $-(CH_2)_{0-4}-N(-H \text{ or } R_{N-5})-CO-R_{N-2}$ where R_{N-5} and R_{N-2} can be the same or different and are as defined above,

5 (25) $-(CH_2)_{0-4}-NR_{N-2}R_{N-3}$ where R_{N-2} and R_{N-3} can be the same or different and are as defined above,

(26) $-(CH_2)_{0-4}-R_{N-4}$ where R_{N-4} is as defined above,

(27) $-(CH_2)_{0-4}-O-CO-(C_1-C_6 \text{ alkyl})$,

(28) $-(CH_2)_{0-4}-O-P(O)-(OR_{N-aryl-1})_2$ where $R_{N-aryl-1}$ is –
10 H or C_1-C_4 alkyl,

(29) $-(CH_2)_{0-4}-O-CO-N(R_{N-5})_2$ where R_{N-5} is as defined above,

(30) $-(CH_2)_{0-4}-O-CS-N(R_{N-5})_2$ where R_{N-5} is as defined above,

15 (31) $-(CH_2)_{0-4}-O-(R_{N-5})_2$ where R_{N-5} is as defined above,

(32) $-(CH_2)_{0-4}-O-(R_{N-5})_2-COOH$ where R_{N-5} is as defined above,

(33) $-(CH_2)_{0-4}-S-(R_{N-5})_2$ where R_{N-5} is as defined
20 above,

(34) $-(CH_2)_{0-4}-O-(C_1-C_6 \text{ alkyl optionally substituted with one, two, three, four, or five of } -F)$,

(35) C_3-C_7 cycloalkyl,

(36) C_2-C_6 alkenyl with one or two double bonds
25 optionally substituted with C_1-C_3 alkyl, $-F$, $-Cl$, $-Br$, $-I$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, C_1-C_3 alkoxy, or $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

(37) C_2-C_6 alkynyl with one or two triple bonds
optionally substituted with C_1-C_3 alkyl, $-F$, $-Cl$, $-Br$, $-I$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, C_1-C_3 alkoxy, or $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

30 (38) $-(CH_2)_{0-4}-N(-H \text{ or } R_{N-5})-SO_2-R_{N-2}$ where R_{N-5} and R_{N-2} can be the same or different and are as described above, or

(39) $-(CH_2)_{0-4}-C_3-C_7$ cycloalkyl,

(B) -R_{N-heteroaryl} where R_{N-heteroaryl} is selected from the group

consisting of:

- | | |
|----|-------------------|
| | pyridinyl, |
| | pyrimidinyl, |
| 5 | quinolinyl, |
| | benzothienyl, |
| | indolyl, |
| | indolinyl, |
| | pyridazinyl, |
| 10 | pyrazinyl, |
| | isoindolyl, |
| | isoquinolyl, |
| | quinazolinyl, |
| | quinoxalinyl, |
| 15 | phthalazinyl, |
| | imidazolyl, |
| | isoxazolyl, |
| | pyrazolyl, |
| | oxazolyl, |
| 20 | thiazolyl, |
| | indolizinyl, |
| | indazolyl, |
| | benzothiazolyl, |
| | benzimidazolyl, |
| 25 | benzofuranyl, |
| | furanyl, |
| | thienyl, |
| | pyrrolyl, |
| | oxadiazolyl, |
| 30 | thiadiazolyl, |
| | triazolyl, |
| | tetrazolyl, |
| | oxazolopyridinyl, |
| | imidazopyridinyl, |

	isothiazolyl,
	naphthyridinyl,
	cinnoliny1,
	carbazolyl,
5	beta-carboliny1,
	isochromanyl,
	chromanyl,
	tetrahydroisoquinoliny1,
	isoindoliny1,
10	isobenzotetrahydrofuranyl,
	isobenzotetrahydrothienyl,
	isobenzothienyl,
	benzoxazolyl,
	pyridopyridiny1,
15	benzotetrahydrofuranyl,
	benzotetrahydrothienyl,
	puriny1,
	benzodioxolyl,
	triaziny1,
20	phenoxaziny1,
	phenothiaziny1,
	pteridiny1,
	benzothiazolyl,
	imidazopyridiny1,
25	imidazothiazolyl,
	dihydrobenzisoiaziny1,
	benzisoiaziny1,
	benzoxaziny1,
	dihydrobenzisothiaziny1,
30	benzopyranyl,
	benzothiopyranyl,
	coumariny1,
	isocoumariny1,
	chromony1,

chromanonyl,
pyridinyl-N-oxide,
tetrahydroquinolinyl,
dihydroquinolinyl,
5 dihydroquinolinonyl,
dihydroisoquinolinonyl,
dihydrocoumarinyl,
dihydroisocoumarinyl,
isoindolinonyl,
10 benzodioxanyl,
benzoxazolinonyl,
pyrrolyl N-oxide,
pyrimidinyl N-oxide,
pyridazinyl N-oxide,
15 pyrazinyl N-oxide,
quinolinyl N-oxide,
indolyl N-oxide,
indolinyl N-oxide,
isoquinolyl N-oxide,
20 quinazolinyl N-oxide,
quinoxalinyl N-oxide,
phthalazinyl N-oxide,
imidazolyl N-oxide,
isoxazolyl N-oxide,
25 oxazolyl N-oxide,
thiazolyl N-oxide,
indoliziny N-oxide,
indazolyl N-oxide,
benzothiazolyl N-oxide,
30 benzimidazolyl N-oxide,
pyrrolyl N-oxide,
oxadiazolyl N-oxide,
thiadiazolyl N-oxide,
triazolyl N-oxide,

tetrazolyl N-oxide,
 benzothiopyranyl S-oxide, and
 benzothiopyranyl S,S-dioxide

where the $R_{N\text{-heteroaryl}}$ group is bonded by any atom of the
 5 parent $R_{N\text{-heteroaryl}}$ group substituted by hydrogen such that the new bond to the $R_{N\text{-heteroaryl}}$ group replaces the hydrogen atom and its bond, where heteroaryl is optionally substituted with one, two, three, or four of:

(1) $C_1\text{-}C_6$ alkyl, optionally substituted with one, two or
 three substituents selected from the group consisting of $C_1\text{-}C_3$ alkyl, -F, -Cl, -Br, -I,
 10 -OH, -SH, $-C\equiv N$, $-CF_3$, $C_1\text{-}C_3$ alkoxy, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as
 defined above,

(2) -OH,

(3) $-NO_2$,

(4) -F, -Cl, -Br, or -I,

15 (5) $-CO-OH$,

(6) $-C\equiv N$,

(7) $-(CH_2)_{0-4}-CO-NR_{N-2}R_{N-3}$ where R_{N-2} and R_{N-3} are
 the same or different and are selected from the group consisting of:

(a) -H,

20 (b) $-C_1\text{-}C_6$ alkyl optionally substituted with one
 substituent selected from the group consisting of:

(i) -OH, and

(ii) $-NH_2$,

(c) $-C_1\text{-}C_6$ alkyl optionally substituted with
 25 one, two, or three -F, -Cl, -Br, -I,

(d) $-C_3\text{-}C_7$ cycloalkyl,

(e) $-(C_1\text{-}C_2 \text{ alkyl})-(C_3\text{-}C_7 \text{ cycloalkyl})$,

(f) $-(C_1\text{-}C_6 \text{ alkyl})-O-(C_1\text{-}C_3 \text{ alkyl})$,

30 (g) $-C_2\text{-}C_6$ alkenyl with one or two double
 bonds,

(h) $-C_2\text{-}C_6$ alkynyl with one or two triple

bonds,

- and one triple bond,
- (i) $-C_1-C_6$ alkyl chain with one double bond
- (j) $-R_{1-aryl}$ where R_{1-aryl} is as defined above,
- (k) $-R_{1-heteroaryl}$ where $R_{1-heteroaryl}$ is as defined
- 5 above,
- (8) $-(CH_2)_{0-4}-CO-(C_1-C_{12}$ alkyl),
- (9) $-(CH_2)_{0-4}-CO-(C_2-C_{12}$ alkenyl with one, two or
- three double bonds),
- (10) $-(CH_2)_{0-4}-CO-(C_2-C_{12}$ alkynyl with one, two or
- 10 three triple bonds),
- (11) $-(CH_2)_{0-4}-CO-(C_3-C_7$ cycloalkyl),
- (12) $-(CH_2)_{0-4}-CO-R_{1-aryl}$ where R_{1-aryl} is as defined
- above,
- (13) $-(CH_2)_{0-4}-CO-R_{1-heteroaryl}$ where $R_{1-heteroaryl}$ is as
- 15 defined above,
- (14) $-(CH_2)_{0-4}-CO-R_{1-heterocycle}$ where $R_{1-heterocycle}$ is as
- defined above,
- (15) $-(CH_2)_{0-4}-CO-R_{N-4}$ where R_{N-4} is selected from
- the group consisting of morpholinyl, thiomorpholinyl, piperazinyl, piperidinyl,
- 20 homomorpholinyl, homothiomorpholinyl, homothiomorpholinyl S-oxide,
- homothiomorpholinyl S,S-dioxide, pyrrolinyl and pyrrolidinyl where each group is
- optionally substituted with one, two, three, or four of C_1-C_6 alkyl,
- (16) $-(CH_2)_{0-4}-CO-O-R_{N-5}$ where R_{N-5} is selected from
- the group consisting of:
- 25 (a) C_1-C_6 alkyl,
- (b) $-(CH_2)_{0-2}-(R_{1-aryl})$ where R_{1-aryl} is as defined
- above,
- (c) C_2-C_6 alkenyl containing one or two double
- bonds,
- 30 (d) C_2-C_6 alkynyl containing one or two triple
- bonds,
- (e) C_3-C_7 cycloalkyl, and
- (f) $-(CH_2)_{0-2}-(R_{1-heteroaryl})$ where $R_{1-heteroaryl}$ is as
- defined above,

- (17) $-(\text{CH}_2)_{0-4}-\text{SO}_2-\text{NR}_{\text{N}-2}\text{R}_{\text{N}-3}$ where $\text{R}_{\text{N}-2}$ and $\text{R}_{\text{N}-3}$ are as defined above,
- (18) $-(\text{CH}_2)_{0-4}-\text{SO}-(\text{C}_1-\text{C}_8 \text{ alkyl})$,
- (19) $-(\text{CH}_2)_{0-4}-\text{SO}_2-(\text{C}_1-\text{C}_{12} \text{ alkyl})$,
- 5 (20) $-(\text{CH}_2)_{0-4}-\text{SO}_2-(\text{C}_3-\text{C}_7 \text{ cycloalkyl})$,
- (21) $-(\text{CH}_2)_{0-4}-\text{N}(\text{H or } \text{R}_{\text{N}-5})-\text{CO}-\text{O}-\text{R}_{\text{N}-5}$ where $\text{R}_{\text{N}-5}$ can be the same or different and is as defined above,
- (22) $-(\text{CH}_2)_{0-4}-\text{N}(\text{H or } \text{R}_{\text{N}-5})-\text{CO}-\text{N}(\text{R}_{\text{N}-5})_2$, where $\text{R}_{\text{N}-5}$ can be the same or different and is as defined above,
- 10 (23) $-(\text{CH}_2)_{0-4}-\text{N}-\text{CS}-\text{N}(\text{R}_{\text{N}-5})_2$, where $\text{R}_{\text{N}-5}$ can be the same or different and is as defined above,
- (24) $-(\text{CH}_2)_{0-4}-\text{N}(\text{-H or } \text{R}_{\text{N}-5})-\text{CO}-\text{R}_{\text{N}-2}$ where $\text{R}_{\text{N}-5}$ and $\text{R}_{\text{N}-2}$ can be the same or different and are as defined above,
- (25) $-(\text{CH}_2)_{0-4}-\text{NR}_{\text{N}-2}\text{R}_{\text{N}-3}$ where $\text{R}_{\text{N}-2}$ and $\text{R}_{\text{N}-3}$ can be
- 15 the same or different and are as defined above,
- (26) $-(\text{CH}_2)_{0-4}-\text{R}_{\text{N}-4}$ where $\text{R}_{\text{N}-4}$ is as defined above,
- (27) $-(\text{CH}_2)_{0-4}-\text{O}-\text{CO}-(\text{C}_1-\text{C}_6 \text{ alkyl})$,
- (28) $-(\text{CH}_2)_{0-4}-\text{O}-\text{P}(\text{O})-(\text{OR}_{\text{N-aryl-1}})_2$ where $\text{R}_{\text{N-aryl-1}}$ is –
- 20 H or $\text{C}_1-\text{C}_4 \text{ alkyl}$,
- (29) $-(\text{CH}_2)_{0-4}-\text{O}-\text{CO}-\text{N}(\text{R}_{\text{N}-5})_2$ where $\text{R}_{\text{N}-5}$ is as defined above,
- (30) $-(\text{CH}_2)_{0-4}-\text{O}-\text{CS}-\text{N}(\text{R}_{\text{N}-5})_2$ where $\text{R}_{\text{N}-5}$ is as defined above,
- (31) $-(\text{CH}_2)_{0-4}-\text{O}-(\text{R}_{\text{N}-5})_2$ where $\text{R}_{\text{N}-5}$ is as defined
- 25 above,
- (32) $-(\text{CH}_2)_{0-4}-\text{O}-(\text{R}_{\text{N}-5})_2-\text{COOH}$ where $\text{R}_{\text{N}-5}$ is as defined above,
- (33) $-(\text{CH}_2)_{0-4}-\text{S}-(\text{R}_{\text{N}-5})_2$ where $\text{R}_{\text{N}-5}$ is as defined above,
- 30 (34) $-(\text{CH}_2)_{0-4}-\text{O}-(\text{C}_1-\text{C}_6 \text{ alkyl optionally substituted with one, two, three, four, or five of } -\text{F})$,
- (35) $\text{C}_3-\text{C}_7 \text{ cycloalkyl}$,

(36) C₂-C₆ alkenyl with one or two double bonds optionally substituted with C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₃ alkoxy, or -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(37) C₂-C₆ alkynyl with one or two triple bonds
 5 optionally substituted with C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₃ alkoxy, or -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(38) -(CH₂)₀₋₄-N(-H or R_{N-5})-SO₂-R_{N-2} where R_{N-5} and R_{N-2} can be the same or different and are as defined above, or

(39) -(CH₂)₀₋₄-C₃-C₇ cycloalkyl,
 10 (C) R_{N-aryl}-W-R_{N-aryl}, where R_{N-aryl} can be the same or different,

(D) R_{N-aryl}-W-R_{N-heteroaryl},

(E) R_{N-aryl}-W-R_{N-1-heterocycle}, wherein R_{N-1-heterocycle} is the same as R_{1-heterocycle}, and R_{1-heterocycle} is as defined above

15 (F) R_{N-heteroaryl}-W-R_{N-aryl},

(G) R_{N-heteroaryl}-W-R_{N-heteroaryl},

(H) R_{N-heteroaryl}-W-R_{N-1-heterocycle},

(I) R_{N-heterocycle}-W-R_{N-aryl}, wherein R_{N-heterocycle} is the same as R_{1-heterocycle}, and R_{1-heterocycle} is as defined above, and R_{N-aryl} is as defined above,

20 (J) R_{N-heterocycle}-W-R_{N-heteroaryl}, and

(K) R_{N-heterocycle}-W-R_{N-1-heterocycle},

where W is

(25) -(CH₂)₀₋₄-,

(26) -O-,

25 (27) -S(O)₀₋₂-,

(28) -N(R_{N-5})- where R_{N-5} is as defined

above, or

(5) -CO-;

(II) -CO-(C₁-C₁₀ alkyl) where alkyl is optionally substituted with one
 30 three substituents selected from the group consisting of:

(A) -OH,

(B) -C₁-C₆ alkoxy,

(C) -C₁-C₆ thioalkoxy,

- (D) $-\text{CO}-\text{O}-\text{R}_{\text{N}-8}$ where $\text{R}_{\text{N}-8}$ is $-\text{H}$, C_1-C_6 alkyl or $-\text{phenyl}$,
- (E) $-\text{CO}-\text{NR}_{\text{N}-2}\text{R}_{\text{N}-3}$ where $\text{R}_{\text{N}-2}$ and $\text{R}_{\text{N}-3}$ are the same or different and are as defined above,
- (F) $-\text{CO}-\text{R}_{\text{N}-4}$ where $\text{R}_{\text{N}-4}$ is as defined above,
- 5 (G) $-\text{SO}_2-(\text{C}_1-\text{C}_8 \text{ alkyl})$,
- (H) $-\text{SO}_2-\text{NR}_{\text{N}-2}\text{R}_{\text{N}-3}$ where $\text{R}_{\text{N}-2}$ and $\text{R}_{\text{N}-3}$ are the same or different and are as defined above,
- (I) $-\text{NH}-\text{CO}-(\text{C}_1-\text{C}_6 \text{ alkyl})$,
- (J) $-\text{NH}-\text{CO}-\text{O}-\text{R}_{\text{N}-8}$ where $\text{R}_{\text{N}-8}$ is as defined above,
- 10 (K) $-\text{NR}_{\text{N}-2}\text{R}_{\text{N}-3}$ where $\text{R}_{\text{N}-2}$ and $\text{R}_{\text{N}-3}$ are the same or different and are as defined above,
- (L) $-\text{R}_{\text{N}-4}$ where $\text{R}_{\text{N}-4}$ is as defined above,
- (M) $-\text{O}-\text{CO}-(\text{C}_1-\text{C}_6 \text{ alkyl})$,
- (N) $-\text{O}-\text{CO}-\text{NR}_{\text{N}-8}\text{R}_{\text{N}-8}$ where $\text{R}_{\text{N}-8}$ are the same or different
- 15 and are as defined above,
- (O) $-\text{O}-(\text{C}_1-\text{C}_5 \text{ alkyl})-\text{COOH}$,
- (P) $-\text{O}-(\text{C}_1-\text{C}_6 \text{ alkyl})$ optionally substituted with one, two, or three of $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, or $-\text{I}$,
- (Q) $-\text{NH}-\text{SO}_2-(\text{C}_1-\text{C}_6 \text{ alkyl})$, and
- 20 (R) $-\text{F}$, or $-\text{Cl}$,
- (III) $-\text{CO}-(\text{C}_1-\text{C}_6 \text{ alkyl})-\text{O}-(\text{C}_1-\text{C}_6 \text{ alkyl})$ where alkyl is optionally substituted with one, two, or three substituents selected from the group consisting of:
- (A) $-\text{OH}$,
- 25 (B) $-\text{C}_1-\text{C}_6$ alkoxy,
- (C) $-\text{C}_1-\text{C}_6$ thioalkoxy,
- (D) $-\text{CO}-\text{O}-\text{R}_{\text{N}-8}$ where $\text{R}_{\text{N}-8}$ is $-\text{H}$, C_1-C_6 alkyl or $-\text{phenyl}$,
- (E) $-\text{CO}-\text{NR}_{\text{N}-2}\text{R}_{\text{N}-3}$ where $\text{R}_{\text{N}-2}$ and $\text{R}_{\text{N}-3}$ are the same or different and are as defined above,
- 30 (F) $-\text{CO}-\text{R}_{\text{N}-4}$ where $\text{R}_{\text{N}-4}$ is as defined above,
- (G) $-\text{SO}_2-(\text{C}_1-\text{C}_8 \text{ alkyl})$,
- (H) $-\text{SO}_2-\text{NR}_{\text{N}-2}\text{R}_{\text{N}-3}$ where $\text{R}_{\text{N}-2}$ and $\text{R}_{\text{N}-3}$ are the same or different and are as defined above,

- (I) -NH-CO-(C₁-C₆ alkyl),
(J) -NH-CO-O-R_{N-8} where R_{N-8} is as defined above,
(K) -NR_{N-2}R_{N-3} where R_{N-2} and R_{N-3} are the same or different
and are as defined above,
5 (L) -R_{N-4} where R_{N-4} is as defined above,
(M) -O-CO-(C₁-C₆ alkyl),
(N) -O-CO-NR_{N-8}R_{N-8} where R_{N-8} are the same or different
and are as defined above,
(O) -O-(C₁-C₅ alkyl)-COOH,
10 (P) -O-(C₁-C₆ alkyl optionally substituted with one, two, or
three of -F, -Cl, -Br, or -I),
(Q) -NH-SO₂-(C₁-C₆ alkyl), and
(R) -F, or -Cl,
(IV) -CO-(C₁-C₆ alkyl)-S-(C₁-C₆ alkyl) where alkyl is optionally
15 substituted with one, two, or three of substituents selected from the group consisting
of:
(A) -OH,
(B) -C₁-C₆ alkoxy,
(C) -C₁-C₆ thioalkoxy,
20 (D) -CO-O-R_{N-8} where R_{N-8} is as defined above,
(E) -CO-NR_{N-2}R_{N-3} where R_{N-2} and R_{N-3} are the same or
different and are as defined above,
(F) -CO-R_{N-4} where R_{N-4} is as defined above,
(G) -SO₂-(C₁-C₈ alkyl),
25 (H) -SO₂-NR_{N-2}R_{N-3} where R_{N-2} and R_{N-3} are the same or
different and are as defined above,
(I) -NH-CO-(C₁-C₆ alkyl),
(J) -NH-CO-O-R_{N-8} where R_{N-8} is as defined above,
(K) -NR_{N-2}R_{N-3} where R_{N-2} and R_{N-3} are the same or different
30 and are as defined above,
(L) -R_{N-4} where R_{N-4} is as defined above,
(M) -O-CO-(C₁-C₆ alkyl),

(N) $-O-CO-NR_{N-8}R_{N-8}$ where R_{N-8} are the same or different and are as defined above,

(O) $-O-(C_1-C_5 \text{ alkyl})-COOH$,

(P) $-O-(C_1-C_6 \text{ alkyl optionally substituted with one, two, or three of } -F, -Cl, -Br, \text{ or } -I)$,

(Q) $-NH-SO_2-(C_1-C_6 \text{ alkyl})$, and

(R) $-F$, or $-Cl$,

(V) $-CO-CH(-(CH_2)_{0-2}-O-R_{N-10})-(CH_2)_{0-2}-R_{N-aryl}/R_{N-heteroaryl}$ where R_{N-aryl} and $R_{N-heteroaryl}$ are as defined above, where R_{N-10} is selected from the group consisting of:

(A) $-H$,

(B) $C_1-C_6 \text{ alkyl}$,

(C) $C_3-C_7 \text{ cycloalkyl}$,

(D) $C_2-C_6 \text{ alkenyl with one double bond}$,

(E) $C_2-C_6 \text{ alkynyl with one triple bond}$,

(F) R_{1-aryl} where R_{1-aryl} is as defined above, and

(G) $R_{N-heteroaryl}$ where $R_{N-heteroaryl}$ is as defined above, or

(VI) $-CO-(C_3-C_8 \text{ cycloalkyl})$ where alkyl is optionally substituted with one or two substituents selected from the group consisting of:

(A) $-(CH_2)_{0-4}-OH$,

(B) $-(CH_2)_{0-4}-C_1-C_6 \text{ alkoxy}$,

(C) $-(CH_2)_{0-4}-C_1-C_6 \text{ thioalkoxy}$,

(D) $-(CH_2)_{0-4}-CO-O-R_{N-8}$ where R_{N-8} is $-H$, $C_1-C_6 \text{ alkyl}$ or phenyl,

(E) $-(CH_2)_{0-4}-CO-NR_{N-2}R_{N-3}$ where R_{N-2} and R_{N-3} are the same or different and are as defined above,

(F) $-(CH_2)_{0-4}-CO-R_{N-4}$ where R_{N-4} is as defined above,

(G) $-(CH_2)_{0-4}-SO_2-(C_1-C_8 \text{ alkyl})$,

(H) $-(CH_2)_{0-4}-SO_2-NR_{N-2}R_{N-3}$ where R_{N-2} and R_{N-3} are the same or different and are as defined above,

(I) $-(CH_2)_{0-4}-NH-CO-(C_1-C_6 \text{ alkyl})$,

(J) $-NH-CO-O-R_{N-8}$ where R_{N-8} is as defined above,

(K) $-(CH_2)_{0-4}-NR_{N-2}R_{N-3}$ where R_{N-2} and R_{N-3} are the same or different and are as defined above,

(L) $-(CH_2)_{0-4}-R_{N-4}$ where R_{N-4} is as defined above,

(M) $-O-CO-(C_1-C_6 \text{ alkyl})$,

5 (N) $-O-CO-NR_{N-8}R_{N-8}$ where R_{N-8} are the same or different and are as defined above,

(O) $-O-(C_1-C_5 \text{ alkyl})-COOH$,

(P) $-O-(C_1-C_6 \text{ alkyl})$ optionally substituted with one, two, or three of $-F$, $-Cl$, $-Br$, or $-I$,

10 (Q) $-NH-SO_2-(C_1-C_6 \text{ alkyl})$, and

(R) $-F$, or $-Cl$;

where R_A is:

(I) $-C_1-C_{10}$ alkyl optionally substituted with one, two or three
 15 substituents selected from the group consisting of C_1-C_3 alkyl, $-F$, $-Cl$, $-Br$, $-I$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, C_1-C_6 alkoxy, $-O$ -phenyl, $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined above, $-OC=O NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined above, $-S(=O)_{0-2}R_{1-a}$ where R_{1-a} is as defined above, $-NR_{1-a}C=O NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined above, $-C=O NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined above, and $-S(=O)_2 NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

(II) $-(CH_2)_{0-3}-(C_3-C_8)$ cycloalkyl where cycloalkyl can be optionally substituted with one, two or three substituents selected from the group consisting of C_1-C_3 alkyl, $-F$, $-Cl$, $-Br$, $-I$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, C_1-C_6 alkoxy, $-O$ -phenyl, $-CO-OH$, $-CO-O-(C_1-C_4 \text{ alkyl})$, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined
 25 above,

(III) $-(CR_{A-x}R_{A-y})_{0-4}-R_{A-aryl}$ where R_{A-x} and R_{A-y} are

(A) $-H$,

(B) C_1-C_4 alkyl optionally substituted with one or two $-OH$,

(C) C_1-C_4 alkoxy optionally substituted with one, two, or
 30 three of $-F$,

(D) $-(CH_2)_{0-4}-C_3-C_7$ cycloalkyl,

(E) C_2-C_6 alkenyl containing one or two double bonds,

(F) C_2-C_6 alkynyl containing one or two triple bonds, or

(G) phenyl,

and where R_{A-x} and R_{A-y} are taken together with the carbon to which they are attached to form a carbocycle of three, four, five, six or seven carbon atoms, optionally where one carbon atom is replaced by a heteroatom selected from the

5 group consisting of $-O-$, $-S-$, $-SO_2-$, and $-NR_{N-2}-$ and R_{A-aryl} is the same as R_{N-aryl} ,

(IV) $-(CR_{A-x}R_{A-y})_{0-4}-R_{A-heteroaryl}$ where $R_{A-heteroaryl}$ is the same as $R_{N-heteroaryl}$ and R_{A-x} and R_{A-y} are as defined above,

(V) $-(CR_{A-x}R_{A-y})_{0-4}-R_{A-aryl}-R_{A-aryl}$ where R_{A-aryl} , R_{A-x} and R_{A-y} are as defined above,

10 (VI) $-(CR_{A-x}R_{A-y})_{0-4}-R_{A-aryl}-R_{A-heteroaryl}$ where R_{A-aryl} , $R_{A-heteroaryl}$, R_{A-x} and R_{A-y} are as defined above,

(VII) $-(CR_{A-x}R_{A-y})_{0-4}-R_{A-heteroaryl}-R_{A-aryl}$ where $R_{A-heteroaryl}$, R_{A-aryl} , R_{A-x} and R_{A-y} are as defined above,

(VIII) $-(CR_{A-x}R_{A-y})_{0-4}-R_{A-heteroaryl}-R_{A-heteroaryl}$ where $R_{A-heteroaryl}$, R_{A-x} and R_{A-y} are as defined above,

(IX) $-(CR_{A-x}R_{A-y})_{0-4}-R_{A-aryl}-R_{A-heterocycle}$ where $R_{A-heterocycle}$ is defined as $R_{1-heterocycle}$, and where R_{A-aryl} , R_{A-x} and R_{A-y} are as defined above,

(X) $-(CR_{A-x}R_{A-y})_{0-4}-R_{A-heteroaryl}-R_{A-heterocycle}$ where $R_{A-heteroaryl}$, $R_{A-heterocycle}$, R_{A-x} and R_{A-y} are as defined above,

20 (XI) $-(CR_{A-x}R_{A-y})_{0-4}-R_{A-heterocycle}-R_{A-aryl}$ where $R_{A-heterocycle}$, R_{A-aryl} , R_{A-x} and R_{A-y} are as defined above,

(XII) $-(CR_{A-x}R_{A-y})_{0-4}-R_{A-heterocycle}-R_{A-heteroaryl}$ where $R_{A-heterocycle}$, $R_{A-heteroaryl}$, R_{A-x} and R_{A-y} are as defined above,

(XIII) $-(CR_{A-x}R_{A-y})_{0-4}-R_{A-heterocycle}-R_{A-heterocycle}$ where $R_{A-heterocycle}$, R_{A-x} and R_{A-y} are as defined above,

25 (XIV) $-(CR_{A-x}R_{A-y})_{0-4}-R_{A-heterocycle}$ where $R_{A-heterocycle}$, R_{A-x} and R_{A-y} are as defined above,

(XV) $-[C(R_{A-1})(R_{A-2})]_{1-3}-CO-N(R_{A-3})_2$ where R_{A-1} and R_{A-2} are the same or different and are selected from the group consisting of:

30 (A) $-H$,

(B) $-C_1-C_6$ alkyl, optionally substituted with one, two or three substituents selected from the group consisting of C_1-C_3 alkyl, $-F$, $-Cl$, $-Br$, $-I$, $-OH$,

-SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(C) C₂-C₆ alkenyl with one or two double bonds, optionally substituted with one, two or three substituents selected from the group consisting of
 5 C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(D) C₂-C₆ alkynyl with one or two triple bonds, optionally substituted with one, two or three substituents selected from the group consisting of
 10 C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(E) -(CH₂)₁₋₂-S(O)₀₋₂-(C₁-C₆ alkyl),

(F) -(CH₂)₀₋₄-C₃-C₇ cycloalkyl, optionally substituted with one, two or three substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, and -NR_{1-a}R_{1-b} where
 15 R_{1-a} and R_{1-b} are as defined above,

(G) -(C₁-C₄ alkyl)-R_{A'-aryl} where R_{A'-aryl} is as defined for R_{1-aryl},

aryl,

(H) -(C₁-C₄ alkyl)-R_{A-heteroaryl} where R_{A-heteroaryl} is as defined above,

(I) -(C₁-C₄ alkyl)-R_{A-heterocycle} where R_{A-heterocycle} is as defined above,

(J) -R_{A-heteroaryl} where R_{A-heteroaryl} is as defined above,

(K) -R_{A-heterocycle} where R_{A-heterocycle} is as defined above,

(M) -(CH₂)₁₋₄-R_{A-4}-(CH₂)₀₋₄-R_{A'-aryl} where R_{A-4} is -O-, -S- or
 25 -NR_{A-5}- where R_{A-5} is C₁-C₆ alkyl, and where R_{A'-aryl} is defined above,

(N) -(CH₂)₁₋₄-R_{A-4}-(CH₂)₀₋₄-R_{A-heteroaryl} where R_{A-4} and R_{A-heteroaryl} are as defined above, and

(O) -R_{A'-aryl} where R_{A'-aryl} is as defined above,

and where R_{A-3} is the same or different and is:

(A) -H,

(B) -C₁-C₆ alkyl optionally substituted with one, two or three substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH,

-SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(C) C₂-C₆ alkenyl with one or two double bonds, optionally substituted with one, two or three substituents selected from the group consisting of
 5 C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(D) C₂-C₆ alkynyl with one or two triple bonds, optionally substituted with one, two or three substituents selected from the group consisting of
 10 C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(E) -(CH₂)₀₋₄-C₃-C₇ cycloalkyl, optionally substituted with one, two or three substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

15 (F) -R_{A'-aryl} where R_{A'-aryl} is as defined above,

(G) -R_{A-heteroaryl} where R_{A-heteroaryl} is as defined above,

(H) -R_{A-heterocycle} where R_{A-heterocycle} is as defined above,

(I) -(C₁-C₄ alkyl)-R_{A'-aryl} where R_{A'-aryl} is as defined above,

20 (J) -(C₁-C₄ alkyl)-R_{A-heteroaryl} where R_{A-heteroaryl} is as defined above,

(K) -(C₁-C₄ alkyl)-R_{A-heterocycle} where R_{A-heterocycle} is as defined above, or

(XVI) -CH(R_{A-aryl})₂ where R_{A-aryl} are the same or different and are as
 25 defined above,

(XVII) -CH(R_{A-heteroaryl})₂ where R_{A-heteroaryl} are the same or different and are as defined above,

(XVIII) -CH(R_{A-aryl})(R_{A-heteroaryl}) where R_{A-aryl} and R_{A-heteroaryl} are as defined above,

30 (XIX) -cyclopentyl, -cyclohexyl, or -cycloheptyl ring fused to R_{A-aryl}, R_{A-heteroaryl}, R_{A-heterocycle} where R_{A-aryl} or R_{A-heteroaryl} or R_{A-heterocycle} are as defined above where one carbon of cyclopentyl, cyclohexyl, or -cycloheptyl is optionally replaced with NH, NR_{N-5}, O, or S(=O)₀₋₂, and where cyclopentyl, cyclohexyl, or -cycloheptyl

can be optionally substituted with one or two -C₁-C₃ alkyl, -F, -OH, -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, =O, or -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(XX) C₂-C₁₀ alkenyl containing one or two double bonds optionally substituted with one, two or three substituents selected from the group consisting of
 5 C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(XXI) C₂-C₁₀ alkynyl containing one or two triple bonds optionally substituted with one, two or three substituents selected from the group consisting of
 10 C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(XXI) -(CH₂)₀₋₁-CHR_{A-6}-(CH₂)₀₋₁-R_{A-aryl} where R_{A-aryl} is as defined above and R_{A-6} is -(CH₂)₀₋₆-OH,

(XXII) -(CH₂)₀₋₁-CHR_{A-6}-(CH₂)₀₋₁-R_{A-heteroaryl} where R_{A-heteroaryl} and R_{A-6} is as defined above,

15 (XXIII) -CH(-R_{A-aryl} or R_{A-heteroaryl})-CO-O(C₁-C₄ alkyl) where R_{A-aryl} and R_{A-heteroaryl} are as defined above,

(XXIV) -CH(-CH₂-OH)-CH(-OH)-micro-NO₂,

(XXV) (C₁-C₆ alkyl)-O-(C₁-C₆ alkyl)-OH,

(XXVII) -CH₂-NH-CH₂-CH(-O-CH₂-CH₃)₂,

20 (XXVIII) -H,

(XXIX) -(CH₂)₀₋₆-C(=NR_{1-a})(NR_{1-a}R_{1-b}) where R_{1-a} and R_{1-b} are as defined above; or

(XXX)

-C=OC(HR₆)NHR₇, where R₆ and R₇ are as defined below,

25 -C=OR₇, where R₇ is as defined below,

-C=OOR₇, where R₇ is as defined below, or

-SOOR₇ where R₇ is as defined below,

wherein R₆ is:

hydrogen,

30 C₁ - C₃ alkyl,

phenyl,

thioalkoxyalkyl,

alkyl substituted aryl,

	cycloalkyl,
	cycloalkylalkyl,
	hydroxyalkyl,
	alkoxyalkyl,
5	aryloxyalkyl,
	haloalkyl,
	carboxyalkyl,
	alkoxycarbonylalkyl,
	aminoalkyl,
10	(N-protected)aminoalkyl,
	alkylaminoalkyl,
	((N-protected)(alkyl)amino)alkyl,
	dialkylaminoalkyl,
	guanidinoalkyl,
15	lower alkenyl,
	heterocyclic,
	(heterocyclic)alkyl),
	arylthioalkyl,
	arylsulfonylalkyl,
20	(heterocyclic)thioalkyl,
	(heterocyclic)sulfonylalkyl,
	(heterocyclic)oxyalkyl,
	arylalkoxyalkyl,
	arylthioalkoxyalkyl,
25	arylalkylsulfonylalkyl,
	(heterocyclic))alkoxyalkyl,
	(heterocyclic)thioalkoxyalkyl,
	(heterocyclic)alkylsulfonylalkyl,
	cycloalkyloxyalkyl,
30	cycloalkylthioalkyl,
	cycloalkylsulfonylalkyl,
	cycloalkylalkoxyalkyl,
	cycloalkylthioalkoxyalkyl,
	cycloalkylalkylsulfonylalkyl,

aminocarbonyl,
alkylaminocarbonyl,
dialkylaminocarbonyl,
aroalkyl,
5 (heterocyclic)carbonylalkyl,
polyhydroxyalkyl,
aminocarbonylalkyl,
alkylaminocarbonylalkyl,
dialkylaminocarbonylalkyl,
10 aryloxyalkyl, or
alkylsulfonylalkyl,
wherein heterocyclic is pyridyl, thiazolyl, isothiazolyl,
oxazolyl, isoxazolyl, furanyl, thienyl, tetrahydrofuranyl, tetrahydrothienyl and
tetrahydro[2H]pyranyl and wherein the heterocycle is unsubstituted or substituted
15 with one to three substituents independently selected from hydroxy, halo, amino,
alkylamino, dialkylamino, alkoxy, polyalkoxy, haloalkyl, cycloalkyl,
cycloalkylalkyl, aryl, arylalkyl, COOH, -SO₃H, lower alkenyl or lower alkyl;
wherein R₇ is:
C₁ - C₃ alkyl,
20 phenyl,
thioalkoxyalkyl,
(aryl)alkyl,
cycloalkyl,
cycloalkylalkyl,
25 hydroxyalkyl,
alkoxyalkyl,
aryloxyalkyl,
haloalkyl,
carboxyalkyl,
30 alkoxycarbonylalkyl,
aminoalkyl,
(N-protected)aminocalkyl,
alkylaminoalkyl,
((N-protected)(alkyl)amino)alkyl,

5 dialkylaminoalkyl,
 guanidinoalkyl,
 lower alkenyl,
 heterocyclic,
 (heterocyclic)alkyl),
 arylthioalkyl,
 arylsulfonylalkyl,
 (heterocyclic)thioalkyl,
 (heterocyclic)sulfonylalkyl,
10 (heterocyclic)oxyalkyl,
 arylalkoxyalkyl,
 arylthioalkoxyalkyl,
 arylalkylsulfonylalkyl,
 (heterocyclic))alkoxyalkyl,
15 (heterocyclic)thioalkoxyalkyl,
 (heterocyclic)alkylsulfonylalkyl,
 cycloalkyloxyalkyl,
 cycloalkylthioalkyl,
 cycloalkylsulfonylalkyl,
20 cycloalkylalkoxyalkyl,
 cycloalkylthioalkoxyalkyl,
 cycloalkylalkylsulfonylalkyl,
 aminocarbonyl,
 alkylaminocarbonyl,
25 dialkylaminocarbonyl,
 aroylalkyl,
 (heterocyclic)carbonylalkyl,
 polyhydroxyalkyl,
 aminocarbonylalkyl,
30 alkylaminocarbonylalkyl,
 dialkylaminocarbonylalkyl,
 aryloxyalkyl, or
 alkylsulfonylalkyl,

wherein heterocyclic is pyridyl, thiazolyl, isothiazolyl, oxazolyl, isoxazolyl, furanyl, thienyl, tetrahydrofuranyl, tetrahydrothienyl and tetrahydro[2H]pyranyl and wherein the heterocycle is unsubstituted or substituted with one to three substituents independently selected from hydroxy, halo, amino, alkylamino, dialkylamino, alkoxy, polyalkoxy, haloalkyl, cycloalkyl, cycloalkylalkyl, aryl, arylalkyl, COOH, -SO₃H, lower alkenyl or lower alkyl;

where X is -N, or -O, with the proviso that when X is O, R_B is absent; and when X is N,

R_B is:

(I)-C₁-C₁₀ alkyl optionally substituted with one, two or three substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above, -OC=O NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above, -S(=O)₀₋₂ R_{1-a} where R_{1-a} is as defined above, -NR_{1-a}C=O NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above, -C=O NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above, and -S(=O)₂ NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(II) -(CH₂)₀₋₃-(C₃-C₈) cycloalkyl where cycloalkyl can be optionally substituted with one, two or three substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, -COOH, -CO-O-(C₁-C₄ alkyl), and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(III) -(CR_{B-x}R_{B-y})₀₋₄-R_{B-aryl} where R_{B-x} and R_{B-y} are

(A) -H,
(B) C₁-C₄ alkyl optionally substituted with one or two -OH,
(C) C₁-C₄ alkoxy optionally substituted with one, two, or three of -F,

(D) -(CH₂)₀₋₄-C₃-C₇ cycloalkyl,
(E) C₂-C₆ alkenyl containing one or two double bonds,
(F) C₂-C₆ alkynyl containing one or two triple bonds,
(G) phenyl,

and where R_{B-x} and R_{B-y} are taken together with the carbon to which they are attached to form a carbocycle of three, four, five, six or seven carbon atoms, optionally where one carbon atom is replaced by a heteroatom selected from

the group consisting of -O-, -S-, -SO₂-, and -NR_{N-2} where R_{N-2} is as defined above, and R_{B-aryl} is the same as R_{N-aryl} and is defined above

(IV) -(CR_{B-x}R_{B-y})₀₋₄-R_{B-heteroaryl} where R_{B-heteroaryl} is the same as R_{N-heteroaryl}, R_{B-x}, and R_{B-y} are as defined above,

5 (V) -(CR_{B-x}R_{B-y})₀₋₄-R_{B-aryl}-R_{B-aryl} where R_{B-aryl}, R_{B-x}, and R_{B-y} are as defined above,

(VI) -(CR_{B-x}R_{B-y})₀₋₄-R_{B-aryl}-R_{B-heteroaryl} where R_{B-aryl}, R_{B-heteroaryl}, R_{B-x} and R_{B-y} are as defined above,

(VII) -(CR_{B-x}R_{B-y})₀₋₄-R_{B-heteroaryl}-R_{B-aryl} where R_{B-heteroaryl}, R_{B-aryl}, R_{B-x} and R_{B-y} are as defined above,

(VIII) -(CR_{B-x}R_{B-y})₀₋₄-R_{B-heteroaryl}-R_{B-heteroaryl} where R_{B-heteroaryl}, R_{B-x} and R_{B-y} are as defined above,

(IX) -(CR_{B-x}R_{B-y})₀₋₄-R_{B-aryl}-R_{B-heterocycle} where R_{B-heterocycle} is defined as R_{1-heterocycle}, and where R_{B-aryl}, R_{B-x} and R_{B-y} are as defined above,

15 (X) -(CR_{B-x}R_{B-y})₀₋₄-R_{B-heteroaryl}-R_{B-heterocycle} where R_{B-heteroaryl}, R_{B-heterocycle}, R_{B-x} and R_{B-y} are as defined above,

(XI) -(CR_{B-x}R_{B-y})₀₋₄-R_{B-heterocycle}-R_{B-aryl} where R_{B-heterocycle}, R_{B-aryl}, R_{B-x} and R_{B-y} are as defined above,

(XII) -(CR_{B-x}R_{B-y})₀₋₄-R_{B-heterocycle}-R_{B-heteroaryl} where R_{B-heterocycle}, R_{B-heteroaryl}, R_{B-x} and R_{B-y} are as defined above,

(XIII) -(CR_{B-x}R_{B-y})₀₋₄-R_{B-heterocycle}-R_{B-heterocycle} where R_{B-heterocycle}, R_{B-x} and R_{B-y} are as defined above,

(XIV) -(CR_{B-x}R_{B-y})₀₋₄-R_{B-heterocycle} where R_{B-heterocycle}, R_{B-x} and R_{B-y} are as defined above,

25 (XV) -[C(R_{B-1})(R_{B-2})]₁₋₃-CO-N-(R_{B-3})₂ where R_{B-1} and R_{B-2} are the same or different and are selected from the group consisting of:

(A) -H,

(B) -C₁-C₆ alkyl, optionally substituted with one, two or three substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(C) C₂-C₆ alkenyl with one or two double bonds, optionally substituted with one, two or three substituents selected from the group consisting of

C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(D) C₂-C₆ alkynyl with one or two triple bonds, optionally substituted with one, two or three substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(E) -(CH₂)₁₋₂-S(O)₀₋₂-(C₁-C₆ alkyl),

(F) -(CH₂)₀₋₄-C₃-C₇ cycloalkyl, optionally substituted with one, two or three substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(G) -(C₁-C₄ alkyl)-R_{B'-aryl} where R_{B'-aryl} is as defined above for R_{1-aryl},

(H) -(C₁-C₄ alkyl)-R_{B-heteroaryl} where R_{B-heteroaryl} is as defined above,

(I) -(C₁-C₄ alkyl)-R_{B-heterocycle} where R_{B-heterocycle} is as defined above,

(J) -R_{B-heteroaryl} where R_{B-heteroaryl} is as defined above,

(K) -R_{B-heterocycle} where R_{B-heterocycle} is as defined above,

(M) -(CH₂)₁₋₄-R_{B-4}-(CH₂)₀₋₄-R_{B'-aryl} where R_{B-4} is -O-, -S- or -NR_{B-5}- where R_{B-5} is C₁-C₆ alkyl, and where R_{B'-aryl} is defined above,

(N) -(CH₂)₁₋₄-R_{B-4}-(CH₂)₀₋₄-R_{B-heteroaryl} where R_{B-4} and R_{B-heteroaryl} are as defined above, and

(O) -R_{B'-aryl} where R_{B'-aryl} is as defined above, and where R_{B-3} is the same or different and is:

(A) -H,

(B) -C₁-C₆ alkyl optionally substituted with one, two or three substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(C) C₂-C₆ alkenyl with one or two double bonds, optionally substituted with one, two or three substituents selected from the group consisting of

C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(D) C₂-C₆ alkynyl with one or two triple bonds, optionally substituted with one, two or three substituents selected from the group consisting of
 5 C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(E) -(CH₂)₀₋₄-C₃-C₇ cycloalkyl, optionally substituted with one, two or three substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, and -NR_{1-a}R_{1-b} where
 10 R_{1-a} and R_{1-b} are as defined above,

(F) -R_{B'-aryl} where R_{B'-aryl} is as defined above,

(G) -R_{B-heteroaryl} where R_{B-heteroaryl} is as defined above,

(H) -R_{B-heterocycle} where R_{B-heterocycle} is as defined above,

(I) -(C₁-C₄ alkyl)-R_{B'-aryl} where R_{B'-aryl} is as defined
 15 above,

(J) -(C₁-C₄ alkyl)-R_{B-heteroaryl} where R_{B-heteroaryl} is as defined above,

(K) -(C₁-C₄ alkyl)-R_{B-heterocycle} where R_{B-heterocycle} is as defined above, or

20 (XVI) -CH(R_{B-aryl})₂ where R_{B-aryl} are the same or different and are as defined above,

(XVII) -CH(R_{B-heteroaryl})₂ where R_{B-heteroaryl} are the same or different and are as defined above,

(XVIII) -CH(R_{B-aryl})(R_{B-heteroaryl}) where R_{B-aryl} and R_{B-heteroaryl} are as
 25 defined above,

(XIX) -cyclopentyl, -cyclohexyl, or -cycloheptyl ring fused to R_{B-aryl} or R_{B-heteroaryl} or R_{B-heterocycle} where R_{B-aryl} or R_{B-heteroaryl} or R_{B-heterocycle} are as defined above where one carbon of cyclopentyl, cyclohexyl, or -cycloheptyl is optionally replaced with NH, NR_{N-5}, O, or S(=O)₀₋₂, and where cyclopentyl, cyclohexyl, or -
 30 cycloheptyl can be optionally substituted with one or two -C₁-C₃ alkyl, -F, -OH, -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, =O, or -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(XX) C₂-C₁₀ alkenyl containing one or two double bonds optionally substituted with one, two or three substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

5 (XXI) C₂-C₁₀ alkynyl containing one or two triple bonds optionally substituted with one, two or three substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(XXI) -(CH₂)₀₋₁-CHR_{C-6}-(CH₂)₀₋₁-R_{B-aryl} where R_{B-aryl} is as defined
10 above and R_{C-6} is -(CH₂)₀₋₆-OH,

(XXII) -(CH₂)₀₋₁-CHR_{B-6}-(CH₂)₀₋₁-R_{B-heteroaryl} where R_{B-heteroaryl} and R_{C-6} is as defined above,

(XXIII) -CH(-R_{B-aryl} or R_{B-heteroaryl})-CO-O(C₁-C₄ alkyl) where R_{B-aryl} and R_{B-heteroaryl} are as defined above,

15 (XXIV) -CH(-CH₂-OH)-CH(-OH)-micro-NO₂,

(XXV) (C₁-C₆ alkyl)-O-(C₁-C₆ alkyl)-OH,

(XXVII) -CH₂-NH-CH₂-CH(-O-CH₂-CH₃)₂,

(XXVIII) -H, or

(XXIX) -(CH₂)₀₋₆-C(=NR_{1-a})(NR_{1-a}R_{1-b}) where R_{1-a} and R_{1-b} are as
20 defined above;

or a pharmaceutically acceptable salt thereof.

55. A method of treatment according to claim 54 where the disease is Alzheimer's disease.

25



56. A method of treatment according to claim 54 where the method is helping prevent or delay the onset of Alzheimer's disease.

57. A method of treatment according to claim 54 where the disease is mild
30 cognitive impairment.

58. A method of treatment according to claim 54 where the disease is Down's syndrome.


59. A method of treatment according to claim 54 where the disease is Hereditary Cerebral Hemorrhage with Amyloidosis of the Dutch-Type.
- 5 60. A method of treatment according to claim 54 where the disease is cerebral amyloid angiopathy.
61. A method of treatment according to claim 54 where the disease is degenerative dementias.
- 10 62. A method of treatment according to claim 54 where the disease is diffuse Lewy body type of Alzheimer's disease.
63. A method of treatment according to claim 54 where the method is treating an
15 existing disease.
64. A method of treatment according to claim 54 where the method is preventing a disease from developing.
- 20 65. A method of treatment according to claim 54 where the therapeutically effective amount for oral administration is from about 0.1 mg/day to about 1,000 mg/day; for parenteral, sublingual, intranasal, intrathecal administration is from about 0.5 to about 100 mg/day; for depo administration and implants is from about 0.5 mg/day to about 50 mg/day; for topical administration is from about 0.5 mg/day to about 200
25 mg/day; for rectal administration is from about 0.5 mg to about 500 mg.
66. A method of treatment according to claim 65 where the therapeutically effective amount is for oral administration is from about 1 mg/day to about 100 mg/day and for parenteral administration is from about 5 to about 50 mg daily.
- 30 67. A method of treatment according to claim 66 where the therapeutically effective amount for oral administration is from about 5 mg/day to about 50 mg/day.
68. A method of treatment according to claim 54:

where R_1 is:




where R_N is:


5 $R_{N-1}-X_N$, where X_N is selected from the group consisting of:





where R_{N-1} is selected from the group consisting of:






10 $-R_{N\text{-heteroaryl}}, \text{ or}$



where R_A and R_B are each independently:




15 $-(CR_{A-x}R_{A-y})_{0-4}-R_{A\text{-aryl}},$





heterocycle; and

20 where X is:



69. A method of treatment according to claim 68:


25 where R_1 is:




where R_2 is $-H$;

where R_3 is $-H$;

30 where R_N is:



where R_{N-1} is selected from the group consisting of:



$-R_{N\text{-heteroaryl}}$;

where R_A and R_B are each independently:

$-(CH_2)_{0-3}-(C_3-C_7)$ cycloalkyl,

$-(CR_{A-x}R_{A-y})_{0-4}-R_{A\text{-aryl}}$,

5 $-(CR_{A-x}R_{A-y})_{0-4}-R_{A\text{-heteroaryl}}$,

$-(CR_{A-x}R_{A-y})_{0-4}-R_{A\text{-heterocycle}}$, or

$-\text{cyclopentyl}$ or $-\text{cyclohexyl}$ ring fused to a $R_{A\text{-aryl}}$ or $R_{A\text{-heteroaryl}}$ or $R_{A\text{-heterocycle}}$; and

where X is:

10 $-N$, or

$-O$ with the proviso that if X is O, R_B is absent.

70. A method of treatment according to claim 69 where R_A and R_B are each independently:

15 $-(CR_{C-x}R_{C-y})_{0-4}-R_{C\text{-aryl}}$,

$-(CR_{C-x}R_{C-y})_{0-4}-R_{C\text{-heteroaryl}}$,

$-\text{cyclopentyl}$ or $-\text{cyclohexyl}$ ring fused to a $R_{C\text{-aryl}}$ or $R_{C\text{-heteroaryl}}$ or $R_{C\text{-heterocycle}}$.

71. A method of treatment according to claim 54 where R_1 is:

20 $-(CH_2)-(R_{1\text{-aryl}})$ where $R_{1\text{-aryl}}$ is phenyl.

72. A method of treatment according to claim 71 where R_1 is:

$-(CH_2)-(R_{1\text{-aryl}})$ where $R_{1\text{-aryl}}$ is phenyl substituted with two -F.

25 73. A method of treatment according to claim 72 where the -F substitution is 3,5-difluorobenzyl.

74. A method of treatment according to claim 54 where R_2 is -H.

30 75. A method of treatment according to claim 54 where R_3 is -H.

76. A method of treatment according to claim 54 where R_N is

$R_{N-1}-X_N-$ where X_N is -CO-, where R_{N-1} is $R_{N\text{-aryl}}$ where $R_{N\text{-aryl}}$ is phenyl substituted with one -CO-NR_{N-2}R_{N-3} where the substitution on phenyl is 1,3-.

77. A method of treatment according to claim 76 where R_{N-2} and R_{N-3} are the same and are C_3 alkyl.
- 5 78. A method of treatment according to claim 54 where R_N is
 $R_{N-1}-X_N-$ where X_N is -CO-, where R_{N-1} is R_{N-aryl} where R_{N-aryl} is phenyl substituted with one C_1 alkyl and with one -CO-NR_{N-2}R_{N-3} where the substitution on the phenyl is 1,3,5-.
- 10 79. A method of treatment according to claim 78 where R_{N-2} and R_{N-3} are the same and are C_3 alkyl.
80. A method of treatment according to claim 54 where R_N is
 $R_{N-1}-X_N-$ where X_N is -CO-, where R_{N-1} is $R_{N-heteroaryl}$ where $R_{N-heteroaryl}$ is
15 substituted with one -CO-NR_{N-2}R_{N-3}.
81. A method of treatment according to claim 80 where R_{N-2} and R_{N-3} are the same and are - C_3 alkyl.
- 20 82. A method of treatment according to claim 54, where R_A is:
 $-(CR_{A-x}R_{A-y})_{0-4}-R_{A-aryl}$ where R_{A-aryl} is phenyl,
 $-(CR_{A-x}R_{A-y})_{0-4}-R_{A-heteroaryl}$,
 $-\text{cyclopentyl}$ or $-\text{cyclohexyl}$ ring fused to a R_{A-aryl} or $R_{A-heteroaryl}$ or R_{A-}
heterocycle.
- 25 83. A method of treatment according to claim 82, where R_A is:
 $-(CR_{A-x}R_{A-y})_{0-4}-R_{A-aryl}$ where R_{A-aryl} is phenyl.
84. A method of treatment according to claim 83, where phenyl is substituted in the
30 3-position or 3,5-positions.
85. A method of treatment according to claim 82, where R_A is
 $-(CH_2)-R_{A-heteroaryl}$.



86. A method of treatment according to claim 82, where R_A is:



87. A method of treatment according to claim 86, where R_A is:


5 -cyclohexyl ring fused to a phenyl ring.

88. A method of treatment according to claim 54, where R_B is:



10 -cyclopentyl or -cyclohexyl ring fused to a R_{B-aryl} or $R_{B-heteroaryl}$ or $R_{B-heterocycle}$.


89. A method of treatment according to claim 88, where R_B is:




15

90. A method of treatment according to claim 89 where phenyl is substituted in the 3-position or 3,5-positions.

91. A method of treatment according to claim 88 where R_B is:



92. A method of treatment according to claim 88 where R_B is:



25 93. A method of treatment according to claim 88 where R_B is:

 -cyclohexyl ring fused to a phenyl ring.


94. A method of treatment according to claim 54 where the pharmaceutically

acceptable salt is selected from the group consisting of salts of the following acids

30 acetic, aspartic, benzenesulfonic, benzoic, bicarbonic, bisulfuric, bitartaric, butyric, calcium edetate, camsyllic, carbonic, chlorobenzoic, citric, edetic, edisyllic, estolic, esyl, esylic, formic, fumaric, gluceptic, gluconic, glutamic, glycolylarsanilic, hexamic, hexylresorcinoic, hydrabamic, hydrobromic, hydrochloric, hydroiodic, hydroxynaphthoic, isethionic, lactic, lactobionic, maleic, malic, malonic, mandelic,

methanesulfonic, methylnitric, methylsulfuric, mucic, muconic, napsylic, nitric, oxalic, p-nitromethanesulfonic, pamoic, pantothenic, phosphoric, monohydrogen phosphoric, dihydrogen phosphoric, phthalic, polygalactouronic, propionic, salicylic, stearic, succinic, succinic, sulfamic, sulfanilic, sulfonic, sulfuric, tannic, tartaric, teoclic and toluenesulfonic.

95. A method for inhibiting beta-secretase activity, comprising exposing said beta-secretase to an effective inhibitory amount of a compound of the formula XV



where R_1 is:

(I) $\text{C}_1\text{-C}_6$ alkyl, optionally substituted with one, two or three substituents selected from the group consisting of $\text{C}_1\text{-C}_3$ alkyl, $\text{C}_1\text{-C}_7$ alkyl (optionally substituted with $\text{C}_1\text{-C}_3$ alkyl and $\text{C}_1\text{-C}_3$ alkoxy), $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, $-\text{OH}$, $-\text{SH}$, $-\text{C}\equiv\text{N}$, $-\text{CF}_3$, $\text{C}_1\text{-C}_3$ alkoxy, $-\text{NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are $-\text{H}$ or $\text{C}_1\text{-C}_6$ alkyl, and $-\text{OC}=\text{O NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

(II) $-\text{CH}_2\text{-S(O)}_{0-2}\text{-(C}_1\text{-C}_6\text{ alkyl)}$,

(III) $-\text{CH}_2\text{-CH}_2\text{-S(O)}_{0-2}\text{-(C}_1\text{-C}_6\text{ alkyl)}$,

(IV) $\text{C}_2\text{-C}_6$ alkenyl with one or two double bonds, optionally substituted with one, two or three substituents selected from the group consisting of $-\text{F}$, $-\text{Cl}$, $-\text{OH}$, $-\text{SH}$, $-\text{C}\equiv\text{N}$, $-\text{CF}_3$, $\text{C}_1\text{-C}_3$ alkoxy, and $-\text{NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are $-\text{H}$ or $\text{C}_1\text{-C}_6$ alkyl,

(V) $\text{C}_2\text{-C}_6$ alkynyl with one or two triple bonds, optionally substituted with one, two or three substituents selected from the group consisting of $-\text{F}$, $-\text{Cl}$, $-\text{OH}$, $-\text{SH}$, $-\text{C}\equiv\text{N}$, $-\text{CF}_3$, $\text{C}_1\text{-C}_3$ alkoxy, and $-\text{NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are $-\text{H}$ or $\text{C}_1\text{-C}_6$ alkyl,

(VI) $-(\text{CH}_2)_{n1}\text{-(R}_{1\text{-aryl}})$ where n_1 is zero or one and where $\text{R}_{1\text{-aryl}}$ is phenyl, 1-naphthyl, 2-naphthyl and indanyl, indenyl, dihydronaphthalyl, or tetralinyl optionally substituted with one, two, three or four of the following substituents on the aryl ring:

(A) C₁-C₆ alkyl optionally substituted with one, two or three substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, and C₁-C₃ alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

5 (B) C₂-C₆ alkenyl with one or two double bonds, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, -C≡N, -CF₃, C₁-C₃ alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are -H or C₁-C₆ alkyl,

(C) C₂-C₆ alkynyl with one or two triple bonds, optionally
10 substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, -C≡N, -CF₃, C₁-C₃ alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are -H or C₁-C₆ alkyl,

(D) -F, Cl, -Br or -I,

(F) -C₁-C₆ alkoxy optionally substituted with one, two or
15 three of -F,

(G) -NR_{N-2}R_{N-3} where R_{N-2} and R_{N-3} are as defined below,

(H) -OH,

(I) -C≡N,

(J) C₃-C₇ cycloalkyl, optionally substituted with one, two or
20 three substituents selected from the group consisting of -F, -Cl, -OH, -SH, -C≡N, -CF₃, C₁-C₃ alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are -H or C₁-C₆ alkyl,

(K) -CO-(C₁-C₄ alkyl),

(L) -SO₂-NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(M) -CO-NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

25 or

(N) -SO₂-(C₁-C₄ alkyl),

(VII) -(CH₂)_{n1}-(R_{1-heteroaryl}) where n₁ is as defined above and where R_{1-heteroaryl} is selected from the group consisting of:

pyridinyl,

30 pyrimidinyl,

quinolinyl,

benzothienyl,

indolyl,

	indoliny1,
	pyridaziny1,
	pyraziny1,
	isoquinoly1,
5	quinazolinyl,
	quinoxaliny1,
	phthalaziny1,
	imidazolyl,
	isoxazolyl,
10	pyrazolyl,
	oxazolyl,
	thiazolyl,
	indoliziny1,
	indazolyl,
15	benzothiazolyl,
	benzimidazolyl,
	benzofuranyl,
	furanyl,
	thienyl,
20	pyrrolyl,
	oxadiazolyl,
	thiadiazolyl,
	triazolyl,
	tetrazolyl,
25	oxazolopyridiny1,
	imidazopyridiny1,
	isothiazolyl,
	naphthyridiny1,
	cinnoliny1,
30	carbazolyl,
	beta-carboliny1,
	isochromanyl,
	chromanyl,
	tetrahydroisoquinoliny1,

	isoindoliny1,
	isobenzotetrahydrofurany1,
	isobenzotetrahydrothieny1,
	isobenzothiény1,
5	benzoxazolyl,
	pyridopyridiny1,
	benzotetrahydrofurany1,
	benzotetrahydrothieny1,
	puriny1,
10	benzodioxolyl,
	triaziny1,
	phenoxaziny1,
	phenothiaziny1,
	pteridiny1,
15	benzothiazolyl,
	imidazopyridiny1,
	imidazothiazolyl,
	dihydrobenzisoaxaziny1,
	benzisoaxaziny1,
20	benzoxaziny1,
	dihydrobenzisothiaziny1,
	benzopyrany1,
	benzothiopyrany1,
	coumariny1,
25	isocoumariny1,
	chromony1,
	chromanony1,
	pyridiny1-N-oxide,
	tetrahydroquinoliny1
30	dihydroquinoliny1
	dihydroquinolinony1
	dihydroisoquinolinony1
	dihydrocoumariny1
	dihydroisocoumariny1

isoindolinonyl
benzodioxanyl
benzoxazolinonyl
pyrrolyl N-oxide,
5 pyrimidinyl N-oxide,
pyridazinyl N-oxide,
pyrazinyl N-oxide,
quinolinyl N-oxide,
indolyl N-oxide,
10 indolinyl N-oxide,
isoquinolyl N-oxide,
quinazolinyl N-oxide,
quinoxalinyl N-oxide,
phthalazinyl N-oxide,
15 imidazolyl N-oxide,
isoxazolyl N-oxide,
oxazolyl N-oxide,
thiazolyl N-oxide,
indolizinyll N-oxide,
20 indazolyl N-oxide,
benzothiazolyl N-oxide,
benzimidazolyl N-oxide,
pyrrolyl N-oxide,
oxadiazolyl N-oxide,
25 thiadiazolyl N-oxide,
triazolyl N-oxide,
tetrazolyl N-oxide,
benzothiopyranyl S-oxide, and
benzothiopyranyl S,S-dioxide,

30 where the R_{1-heteroaryl} group is bonded to $-(CH_2)_{n1}-$ by any ring atom of the parent R_{1-heteroaryl} group substituted by hydrogen such that the new bond to the R_{1-heteroaryl} group replaces the hydrogen atom and its bond, where heteroaryl is optionally substituted with one, two, three or four of:

- (1) C₁-C₆ alkyl optionally substituted with one, two or three substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₃ alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,
- 5 (2) C₂-C₆ alkenyl with one or two double bonds, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, -C≡N, -CF₃, C₁-C₃ alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are -H or C₁-C₆ alkyl,
- 10 (3) C₂-C₆ alkynyl with one or two triple bonds, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, -C≡N, -CF₃, C₁-C₃ alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are -H or C₁-C₆ alkyl,
- (4) -F, Cl, -Br or -I,
- (6) -C₁-C₆ alkoxy optionally substituted with one, two,
- 15 or three of -F,
- (7) -NR_{N-2}R_{N-3} where R_{N-2} and R_{N-3} are as defined below,
- (8) -OH,
- (9) -C≡N,
- 20 (10) C₃-C₇ cycloalkyl, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, -C≡N, -CF₃, C₁-C₃ alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are -H or C₁-C₆ alkyl,
- (11) -CO-(C₁-C₄ alkyl),
- (12) -SO₂-NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined
- 25 above,
- (13) -CO-NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above, or
- (14) -SO₂-(C₁-C₄ alkyl), with the proviso that when n₁ is zero R_{1-heteroaryl} is not bonded to the carbon chain by nitrogen; or
- 30 (VIII) -(CH₂)_{n1}-(R_{1-heterocycle}) where n₁ is as defined above and R_{1-heterocycle} is selected from the group consisting of:
- morpholinyl,
- thiomorpholinyl,

thiomorpholinyl S-oxide,
 thiomorpholinyl S,S-dioxide,
 piperazinyl,
 homopiperazinyl,
 5 pyrrolidinyl,
 pyrrolinyl,
 tetrahydropyranyl,
 piperidinyl,
 tetrahydrofuranyl,
 10 tetrahydrothienyl,
 homopiperidinyl,
 homomorpholinyl,
 homothiomorpholinyl,
 homothiomorpholinyl S,S-dioxide,
 15 oxazolidinonyl,
 dihydropyrazolyl,
 dihydropyrrolyl,
 dihydropyrazinyl,
 dihydropyridinyl,
 20 dihydropyrimidinyl,
 dihydrofuryl,
 dihydropyranyl,
 tetrahydrothienyl S-oxide,
 tetrahydrothienyl S,S-dioxide, and
 25 homothiomorpholinyl S-oxide,

where the R_{1-heterocycle} group is bonded by any atom of the parent R_{1-heterocycle} group substituted by hydrogen such that the new bond to the R_{1-heterocycle} group replaces the hydrogen atom and its bond, where heterocycle is optionally substituted with one, two, three or four:

30 (1) C₁-C₆ alkyl optionally substituted with one, two or three substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₃ alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

- (2) C₂-C₆ alkenyl with one or two double bonds, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, -C≡N, -CF₃, C₁-C₃ alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are -H or C₁-C₆ alkyl,
- 5 (3) C₂-C₆ alkynyl with one or two triple bonds, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, -C≡N, -CF₃, C₁-C₃ alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are -H or C₁-C₆ alkyl,
- (4) -F, Cl, -Br or -I,
- 10 (5) C₁-C₆ alkoxy,
- (6) -C₁-C₆ alkoxy optionally substituted with one, two, or three -F,
- (7) -NR_{N-2}R_{N-3} where R_{N-2} and R_{N-3} are as defined below,
- 15 (8) -OH,
- (9) -C≡N,
- (10) C₃-C₇ cycloalkyl, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, -C≡N, -CF₃, C₁-C₃ alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are -H or C₁-C₆ alkyl,
- 20 (11) -CO-(C₁-C₄ alkyl),
- (12) -SO₂-NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,
- (13) -CO-NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,
- 25 (14) -SO₂-(C₁-C₄ alkyl), or
- (15) =O, with the proviso that when n₁ is zero R_{1-heterocycle} is not bonded to the carbon chain by nitrogen;

where R₂ is:

- 30 (I)-H,
- (II) C₁-C₆ alkyl, optionally substituted with one, two or three substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH,

-SH, -C≡N, -CF₃, C₁-C₃ alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(III) -(CH₂)₀₋₄-R₂₋₁ where R₂₋₁ is R_{1-aryl} or R_{1-heteroaryl} where R_{1-aryl} and R_{1-heteroaryl} are as defined above;

5 (IV) C₂-C₆ alkenyl with one or two double bonds, optionally substituted with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, -C≡N, -CF₃, C₁-C₃ alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are -H or C₁-C₆ alkyl,

(V) C₂-C₆ alkynyl with one or two triple bonds, optionally substituted
10 with one, two or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, -C≡N, -CF₃, C₁-C₃ alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are -H or C₁-C₆ alkyl, or

(VI) -(CH₂)₀₋₄-C₃-C₇ cycloalkyl, optionally substituted with one, two
or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, -C≡N,
15 -CF₃, C₁-C₃ alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are -H or C₁-C₆ alkyl;

where R₃ is:

(I)-H,

(II) C₁-C₆ alkyl, optionally substituted with one, two or three
20 substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₃ alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(III) -(CH₂)₀₋₄-R₂₋₁ where R₂₋₁ is R_{1-aryl} or R_{1-heteroaryl} where R_{1-aryl} and R_{1-heteroaryl} are as defined above

25 (IV) C₂-C₆ alkenyl with one or two double bonds,

(V) C₂-C₆ alkynyl with one or two triple bonds; or

(VI) -(CH₂)₀₋₄-C₃-C₇ cycloalkyl, optionally substituted with one, two
or three substituents selected from the group consisting of -F, -Cl, -OH, -SH, -C≡N,
-CF₃, C₁-C₃ alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are -H or C₁-C₆ alkyl,

30 and where R₂ and R₃ are taken together with the carbon to which they are attached to form a carbocycle of three, four, five, six, or seven carbon atoms, optionally where one carbon atom is replaced by a heteroatom selected from the

group consisting of $-O-$, $-S-$, $-SO_2-$, and $-NR_{N-2}-$, where R_{N-2} is selected from the group consisting of:

- (a) $-H$,
- (b) $-C_1-C_6$ alkyl optionally substituted with one
- 5 substituent selected from the group consisting of:
 - (i) $-OH$, and
 - (ii) $-NH_2$,
 - (c) $-C_1-C_6$ alkyl optionally substituted with one, two, or three $-F$, $-Cl$, $-Br$, or $-I$,
 - 10 (d) $-C_3-C_7$ cycloalkyl,
 - (e) $-(C_1-C_2 \text{ alkyl})-(C_3-C_7 \text{ cycloalkyl})$,
 - (f) $-(C_1-C_6 \text{ alkyl})-O-(C_1-C_3 \text{ alkyl})$,
 - (g) $-C_2-C_6$ alkenyl with one or two double bonds,
 - 15 (h) $-C_2-C_6$ alkynyl with one or two triple bonds,
 - (i) $-C_1-C_6$ alkyl chain with one double bond and one triple bond,
 - (j) $-R_{1-aryl}$ where R_{1-aryl} is as defined above, and
 - 20 (k) $-R_{1-heteroaryl}$ where $R_{1-heteroaryl}$ is as defined above;

where R_N is:

- (I) $R_{N-1}-X_N-$ where X_N is selected from the group consisting of:
 - 25 (A) $-CO-$,
 - (B) $-SO_2-$,
 - (C) $-(CR'R'')_{1-6}$ where R' and R'' are the same or different and are $-H$ and C_1-C_4 alkyl,
 - (D) $-CO-(CR'R'')_{1-6}-X_{N-1}$ where X_{N-1} is selected from the
 - 30 group consisting of $-O-$, $-S-$ and $-NR'-$ and where R' and R'' are as defined above, and
 - (E) a single bond;

where R_{N-1} is selected from the group consisting of:

(A) $R_{N\text{-aryl}}$ where $R_{N\text{-aryl}}$ is phenyl, 1-naphthyl, 2-naphthyl, tetralinyl, indanyl, 6,7,8,9-tetrahydro-5H-benzo[a]cycloheptenyl, or dihydronaphthyl optionally substituted with one, two or three of the following substituents which can be the same or different and are:

5 (1) $C_1\text{-}C_6$ alkyl, optionally substituted with one, two or three substituents selected from the group consisting of $C_1\text{-}C_3$ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C \equiv N, -CF₃, $C_1\text{-}C_3$ alkoxy, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(2) -OH,

10 (3) -NO₂,

(4) -F, -Cl, -Br, -I,

(5) -CO-OH,

(6) -C \equiv N,

(7) -(CH₂)₀₋₄-CO-NR_{N-2}R_{N-3} where R_{N-2} and R_{N-3} are

15 the same or different and are selected from the group consisting of:

(a) -H,

(b) $C_1\text{-}C_6$ alkyl optionally substituted with one substituent selected from the group consisting of:

(i) -OH, and

20 (ii) -NH₂,

(c) $C_1\text{-}C_6$ alkyl optionally substituted with one, two, or three -F, -Cl, -Br, or -I,

(d) $C_3\text{-}C_7$ cycloalkyl,

(e) -($C_1\text{-}C_2$ alkyl)-($C_3\text{-}C_7$ cycloalkyl),

25 (f) -($C_1\text{-}C_6$ alkyl)-O-($C_1\text{-}C_3$ alkyl),

(g) $C_2\text{-}C_6$ alkenyl with one or two double bonds,

(h) $C_2\text{-}C_6$ alkynyl with one or two triple bonds,

30 (i) $C_1\text{-}C_6$ alkyl chain with one double bond and one triple bond,

(j) -R_{1-aryl} where R_{1-aryl} is as defined above, and

(k) $-R_{1\text{-heteroaryl}}$ where $R_{1\text{-heteroaryl}}$ is as defined

above,

(8) $-(CH_2)_{0-4}-CO-(C_1-C_{12} \text{ alkyl})$,

(9) $-(CH_2)_{0-4}-CO-(C_2-C_{12} \text{ alkenyl with one, two or}$

5 three double bonds),

(10) $-(CH_2)_{0-4}-CO-(C_2-C_{12} \text{ alkynyl with one, two or}$

three triple bonds),

(11) $-(CH_2)_{0-4}-CO-(C_3-C_7 \text{ cycloalkyl})$,

(12) $-(CH_2)_{0-4}-CO-R_{1\text{-aryl}}$ where $R_{1\text{-aryl}}$ is as defined

10 above,

(13) $-(CH_2)_{0-4}-CO-R_{1\text{-heteroaryl}}$ where $R_{1\text{-heteroaryl}}$ is as

defined above,

(14) $-(CH_2)_{0-4}-CO-R_{1\text{-heterocycle}}$ where $R_{1\text{-heterocycle}}$ is as

defined above,

15 (15) $-(CH_2)_{0-4}-CO-R_{N-4}$ where R_{N-4} is selected from the group consisting of morpholinyl, thiomorpholinyl, piperazinyl, piperidinyl, homomorpholinyl, homothiomorpholinyl, homothiomorpholinyl S-oxide, homothiomorpholinyl S,S-dioxide, pyrrolinyl and pyrrolidinyl where each group is optionally substituted with one, two, three, or four of C_1-C_6 alkyl,

20 (16) $-(CH_2)_{0-4}-CO-O-R_{N-5}$ where R_{N-5} is selected from the group consisting of:

(a) C_1-C_6 alkyl,

(b) $-(CH_2)_{0-2}-(R_{1\text{-aryl}})$ where $R_{1\text{-aryl}}$ is as defined

above,

25 (c) C_2-C_6 alkenyl containing one or two double bonds,

(d) C_2-C_6 alkynyl containing one or two triple

bonds,

(e) C_3-C_7 cycloalkyl, and

30 (f) $-(CH_2)_{0-2}-(R_{1\text{-heteroaryl}})$ where $R_{1\text{-heteroaryl}}$ is as defined above,

(17) $-(CH_2)_{0-4}-SO_2-NR_{N-2}R_{N-3}$ where R_{N-2} and R_{N-3} are

as defined above,

(18) $-(CH_2)_{0-4}-SO-(C_1-C_8 \text{ alkyl})$,

(19) $-(\text{CH}_2)_{0-4}-\text{SO}_2-(\text{C}_1-\text{C}_{12} \text{ alkyl}),$

(20) $-(\text{CH}_2)_{0-4}-\text{SO}_2-(\text{C}_3-\text{C}_7 \text{ cycloalkyl}),$

(21) $-(\text{CH}_2)_{0-4}-\text{N}(\text{H or } \text{R}_{\text{N-5}})-\text{CO}-\text{O}-\text{R}_{\text{N-5}}$ where $\text{R}_{\text{N-5}}$

can be the same or different and is as defined above,

5 (22) $-(\text{CH}_2)_{0-4}-\text{N}(\text{H or } \text{R}_{\text{N-5}})-\text{CO}-\text{N}(\text{R}_{\text{N-5}})_2,$ where $\text{R}_{\text{N-5}}$

can be the same or different and is as defined above,

(23) $-(\text{CH}_2)_{0-4}-\text{N}-\text{CS}-\text{N}(\text{R}_{\text{N-5}})_2,$ where $\text{R}_{\text{N-5}}$ can be the

same or different and is as defined above,

(24) $-(\text{CH}_2)_{0-4}-\text{N}(-\text{H or } \text{R}_{\text{N-5}})-\text{CO}-\text{R}_{\text{N-2}}$ where $\text{R}_{\text{N-5}}$ and

10 $\text{R}_{\text{N-2}}$ can be the same or different and are as defined above,

(25) $-(\text{CH}_2)_{0-4}-\text{NR}_{\text{N-2}}\text{R}_{\text{N-3}}$ where $\text{R}_{\text{N-2}}$ and $\text{R}_{\text{N-3}}$ can be

the same or different and are as defined above,

(26) $-(\text{CH}_2)_{0-4}-\text{R}_{\text{N-4}}$ where $\text{R}_{\text{N-4}}$ is as defined above,

(27) $-(\text{CH}_2)_{0-4}-\text{O}-\text{CO}-(\text{C}_1-\text{C}_6 \text{ alkyl}),$

15 (28) $-(\text{CH}_2)_{0-4}-\text{O}-\text{P}(\text{O})-(\text{OR}_{\text{N-aryl-1}})_2$ where $\text{R}_{\text{N-aryl-1}}$ is –

H or $\text{C}_1-\text{C}_4 \text{ alkyl},$

(29) $-(\text{CH}_2)_{0-4}-\text{O}-\text{CO}-\text{N}(\text{R}_{\text{N-5}})_2$ where $\text{R}_{\text{N-5}}$ is as

defined above,

(30) $-(\text{CH}_2)_{0-4}-\text{O}-\text{CS}-\text{N}(\text{R}_{\text{N-5}})_2$ where $\text{R}_{\text{N-5}}$ is as defined

20 above,

(31) $-(\text{CH}_2)_{0-4}-\text{O}-(\text{R}_{\text{N-5}})_2$ where $\text{R}_{\text{N-5}}$ is as defined

above,

(32) $-(\text{CH}_2)_{0-4}-\text{O}-(\text{R}_{\text{N-5}})_2-\text{COOH}$ where $\text{R}_{\text{N-5}}$ is as

defined above,

25 (33) $-(\text{CH}_2)_{0-4}-\text{S}-(\text{R}_{\text{N-5}})_2$ where $\text{R}_{\text{N-5}}$ is as defined

above,

(34) $-(\text{CH}_2)_{0-4}-\text{O}-(\text{C}_1-\text{C}_6 \text{ alkyl optionally substituted}$

with one, two, three, four, or five of –F),

(35) $\text{C}_3-\text{C}_7 \text{ cycloalkyl},$

30 (36) $\text{C}_2-\text{C}_6 \text{ alkenyl}$ with one or two double bonds

optionally substituted with $\text{C}_1-\text{C}_3 \text{ alkyl}, -\text{F}, -\text{Cl}, -\text{Br}, -\text{I}, -\text{OH}, -\text{SH}, -\text{C}\equiv\text{N}, -\text{CF}_3, \text{C}_1-$

$\text{C}_3 \text{ alkoxy},$ or $-\text{NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

(37) C₂-C₆ alkynyl with one or two triple bonds
optionally substituted with C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-
C₃ alkoxy, or -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(38) -(CH₂)₀₋₄-N(-H or R_{N-5})-SO₂-R_{N-2} where R_{N-5} and
5 R_{N-2} can be the same or different and are as described above, or

(39) -(CH₂)₀₋₄-C₃-C₇ cycloalkyl,

(B) -R_{N-heteroaryl} where R_{N-heteroaryl} is selected from the group
consisting of:

pyridinyl,
10 pyrimidinyl,
quinolinyl,
benzothienyl,
indolyl,
indolinyl,
15 pyridazinyl,
pyrazinyl,
isoindolyl,
isoquinolyl,
quinazolinyl,
20 quinoxalinyl,
phthalazinyl,
imidazolyl,
isoxazolyl,
pyrazolyl,
25 oxazolyl,
thiazolyl,
indolizinyll,
indazolyl,
benzothiazolyl,
30 benzimidazolyl,
benzofuranyl,
furanyl,
thienyl,

	pyrrolyl,
	oxadiazolyl,
	thiadiazolyl,
	triazolyl,
5	tetrazolyl,
	oxazolopyridinyl,
	imidazopyridinyl,
	isothiazolyl,
	naphthyridinyl,
10	cinnolyl,
	carbazolyl,
	beta-carbolinyl,
	isochromanyl,
	chromanyl,
15	tetrahydroisoquinolyl,
	isoindolyl,
	isobenzotetrahydrofuranyl,
	isobenzotetrahydrothienyl,
	isobenzothienyl,
20	benzoxazolyl,
	pyridopyridinyl,
	benzotetrahydrofuranyl,
	benzotetrahydrothienyl,
	purinyl,
25	benzodioxolyl,
	triazinyl,
	phenoxazinyl,
	phenothiazinyl,
	pteridinyl,
30	benzothiazolyl,
	imidazopyridinyl,
	imidazothiazolyl,
	dihydrobenzisoxazinyl,
	benzisoxazinyl,

	benzoxaziny1,
	dihydrobenzisothiaziny1,
	benzopyrany1,
	benzothiopyrany1,
5	coumariny1,
	isocoumariny1,
	chromony1,
	chromanony1,
	pyridiny1-N-oxide,
10	tetrahydroquinoliny1,
	dihydroquinoliny1,
	dihydroquinolinony1,
	dihydroisoquinolinony1,
	dihydrocoumariny1,
15	dihydroisocoumariny1,
	isoindolinony1,
	benzodioxany1,
	benzoxazolinony1,
	pyrroly1 N-oxide,
20	pyrimidiny1 N-oxide,
	pyridaziny1 N-oxide,
	pyraziny1 N-oxide,
	quinoliny1 N-oxide,
	indoly1 N-oxide,
25	indoliny1 N-oxide,
	isoquinolyl N-oxide,
	quinazolinyl N-oxide,
	quinoxaliny1 N-oxide,
	phthalaziny1 N-oxide,
30	imidazolyl N-oxide,
	isoxazolyl N-oxide,
	oxazolyl N-oxide,
	thiazolyl N-oxide,
	indoliziny1 N-oxide,

indazolyl N-oxide,
benzothiazolyl N-oxide,
benzimidazolyl N-oxide,
pyrrolyl N-oxide,
5 oxadiazolyl N-oxide,
thiadiazolyl N-oxide,
triazolyl N-oxide,
tetrazolyl N-oxide,
benzothiopyranyl S-oxide, and
10 benzothiopyranyl S,S-dioxide

where the $R_{N\text{-heteroaryl}}$ group is bonded by any atom of the parent $R_{N\text{-heteroaryl}}$ group substituted by hydrogen such that the new bond to the $R_{N\text{-heteroaryl}}$ group replaces the hydrogen atom and its bond, where heteroaryl is optionally substituted with one, two, three, or four of:

15 (1) $C_1\text{-}C_6$ alkyl, optionally substituted with one, two or three substituents selected from the group consisting of $C_1\text{-}C_3$ alkyl, $-F$, $-Cl$, $-Br$, $-I$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, $C_1\text{-}C_3$ alkoxy, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

(2) $-OH$,
20 (3) $-NO_2$,
(4) $-F$, $-Cl$, $-Br$, or $-I$,
(5) $-CO-OH$,
(6) $-C\equiv N$,
(7) $-(CH_2)_{0-4}-CO-NR_{N-2}R_{N-3}$ where R_{N-2} and R_{N-3} are

25 the same or different and are selected from the group consisting of:

(a) $-H$,
(b) $-C_1\text{-}C_6$ alkyl optionally substituted with one
substituent selected from the group consisting of:
(i) $-OH$, and
30 (ii) $-NH_2$,
(c) $-C_1\text{-}C_6$ alkyl optionally substituted with
one, two, or three $-F$, $-Cl$, $-Br$, $-I$,
(d) $-C_3\text{-}C_7$ cycloalkyl,

- (e) $-(C_1-C_2 \text{ alkyl})-(C_3-C_7 \text{ cycloalkyl})$,
 (f) $-(C_1-C_6 \text{ alkyl})-O-(C_1-C_3 \text{ alkyl})$,
 (g) $-C_2-C_6 \text{ alkenyl}$ with one or two double
 bonds,
 5 (h) $-C_2-C_6 \text{ alkynyl}$ with one or two triple
 bonds,
 (i) $-C_1-C_6 \text{ alkyl chain}$ with one double bond
 and one triple bond,
 (j) $-R_{1-aryl}$ where R_{1-aryl} is as defined above,
 10 (k) $-R_{1-heteroaryl}$ where $R_{1-heteroaryl}$ is as defined
 above,
 (8) $-(CH_2)_{0-4}-CO-(C_1-C_{12} \text{ alkyl})$,
 (9) $-(CH_2)_{0-4}-CO-(C_2-C_{12} \text{ alkenyl}$ with one, two or
 three double bonds),
 15 (10) $-(CH_2)_{0-4}-CO-(C_2-C_{12} \text{ alkynyl}$ with one, two or
 three triple bonds),
 (11) $-(CH_2)_{0-4}-CO-(C_3-C_7 \text{ cycloalkyl})$,
 (12) $-(CH_2)_{0-4}-CO-R_{1-aryl}$ where R_{1-aryl} is as defined
 above,
 20 (13) $-(CH_2)_{0-4}-CO-R_{1-heteroaryl}$ where $R_{1-heteroaryl}$ is as
 defined above,
 (14) $-(CH_2)_{0-4}-CO-R_{1-heterocycle}$ where $R_{1-heterocycle}$ is as
 defined above,
 (15) $-(CH_2)_{0-4}-CO-R_{N-4}$ where R_{N-4} is selected from
 25 the group consisting of morpholinyl, thiomorpholinyl, piperazinyl, piperidinyl,
 homomorpholinyl, homothiomorpholinyl, homothiomorpholinyl S-oxide,
 homothiomorpholinyl S,S-dioxide, pyrrolinyl and pyrrolidinyl where each group is
 optionally substituted with one, two, three, or four of $C_1-C_6 \text{ alkyl}$,
 (16) $-(CH_2)_{0-4}-CO-O-R_{N-5}$ where R_{N-5} is selected from
 30 the group consisting of:
 (a) $C_1-C_6 \text{ alkyl}$,
 (b) $-(CH_2)_{0-2}-(R_{1-aryl})$ where R_{1-aryl} is as defined
 above,

- bonds,
- (c) C₂-C₆ alkenyl containing one or two double bonds,
- bonds,
- (d) C₂-C₆ alkynyl containing one or two triple bonds,
- 5 (e) C₃-C₇ cycloalkyl, and
- (f) -(CH₂)₀₋₂-(R₁-heteroaryl) where R₁-heteroaryl is as defined above,
- (17) -(CH₂)₀₋₄-SO₂-NR_{N-2}R_{N-3} where R_{N-2} and R_{N-3} are as defined above,
- 10 (18) -(CH₂)₀₋₄-SO-(C₁-C₈ alkyl),
- (19) -(CH₂)₀₋₄-SO₂-(C₁-C₁₂ alkyl),
- (20) -(CH₂)₀₋₄-SO₂-(C₃-C₇ cycloalkyl),
- (21) -(CH₂)₀₋₄-N(H or R_{N-5})-CO-O-R_{N-5} where R_{N-5} can be the same or different and is as defined above,
- 15 (22) -(CH₂)₀₋₄-N(H or R_{N-5})-CO-N(R_{N-5})₂, where R_{N-5} can be the same or different and is as defined above,
- (23) -(CH₂)₀₋₄-N-CS-N(R_{N-5})₂, where R_{N-5} can be the same or different and is as defined above,
- (24) -(CH₂)₀₋₄-N(-H or R_{N-5})-CO-R_{N-2} where R_{N-5} and R_{N-2} can be the same or different and are as defined above,
- 20 (25) -(CH₂)₀₋₄-NR_{N-2}R_{N-3} where R_{N-2} and R_{N-3} can be the same or different and are as defined above,
- (26) -(CH₂)₀₋₄-R_{N-4} where R_{N-4} is as defined above,
- (27) -(CH₂)₀₋₄-O-CO-(C₁-C₆ alkyl),
- 25 (28) -(CH₂)₀₋₄-O-P(O)-(OR_{N-aryl-1})₂ where R_{N-aryl-1} is -H or C₁-C₄ alkyl,
- (29) -(CH₂)₀₋₄-O-CO-N(R_{N-5})₂ where R_{N-5} is as defined above,
- (30) -(CH₂)₀₋₄-O-CS-N(R_{N-5})₂ where R_{N-5} is as defined above,
- 30 (31) -(CH₂)₀₋₄-O-(R_{N-5})₂ where R_{N-5} is as defined above,
- (32) -(CH₂)₀₋₄-O-(R_{N-5})₂-COOH where R_{N-5} is as defined above,

(33) $-(\text{CH}_2)_{0-4}-\text{S}-(\text{R}_{\text{N}-5})_2$ where $\text{R}_{\text{N}-5}$ is as defined

above,

(34) $-(\text{CH}_2)_{0-4}-\text{O}-(\text{C}_1-\text{C}_6 \text{ alkyl optionally substituted with one, two, three, four, or five of } -\text{F}),$

5

(35) $\text{C}_3-\text{C}_7 \text{ cycloalkyl},$

(36) $\text{C}_2-\text{C}_6 \text{ alkenyl with one or two double bonds}$

optionally substituted with $\text{C}_1-\text{C}_3 \text{ alkyl}, -\text{F}, -\text{Cl}, -\text{Br}, -\text{I}, -\text{OH}, -\text{SH}, -\text{C}\equiv\text{N}, -\text{CF}_3, \text{C}_1-\text{C}_3 \text{ alkoxy},$ or $-\text{NR}_{1-\text{a}}\text{R}_{1-\text{b}}$ where $\text{R}_{1-\text{a}}$ and $\text{R}_{1-\text{b}}$ are as defined above,

(37) $\text{C}_2-\text{C}_6 \text{ alkynyl with one or two triple bonds}$

10 optionally substituted with $\text{C}_1-\text{C}_3 \text{ alkyl}, -\text{F}, -\text{Cl}, -\text{Br}, -\text{I}, -\text{OH}, -\text{SH}, -\text{C}\equiv\text{N}, -\text{CF}_3, \text{C}_1-\text{C}_3 \text{ alkoxy},$ or $-\text{NR}_{1-\text{a}}\text{R}_{1-\text{b}}$ where $\text{R}_{1-\text{a}}$ and $\text{R}_{1-\text{b}}$ are as defined above,

(38) $-(\text{CH}_2)_{0-4}-\text{N}(-\text{H or } \text{R}_{\text{N}-5})-\text{SO}_2-\text{R}_{\text{N}-2}$ where $\text{R}_{\text{N}-5}$ and

$\text{R}_{\text{N}-2}$ can be the same or different and are as defined above, or

(39) $-(\text{CH}_2)_{0-4}-\text{C}_3-\text{C}_7 \text{ cycloalkyl},$

15

(C) $\text{R}_{\text{N-aryl}}-\text{W}-\text{R}_{\text{N-aryl}},$ where $\text{R}_{\text{N-aryl}}$ can be the same or

different,

(D) $\text{R}_{\text{N-aryl}}-\text{W}-\text{R}_{\text{N-heteroaryl}},$

(E) $\text{R}_{\text{N-aryl}}-\text{W}-\text{R}_{\text{N-1-heterocycle}},$ wherein $\text{R}_{\text{N-1-heterocycle}}$ is the same

as $\text{R}_{1-\text{heterocycle}},$ and $\text{R}_{1-\text{heterocycle}}$ is as defined above

20

(F) $\text{R}_{\text{N-heteroaryl}}-\text{W}-\text{R}_{\text{N-aryl}},$

(G) $\text{R}_{\text{N-heteroaryl}}-\text{W}-\text{R}_{\text{N-heteroaryl}},$

(H) $\text{R}_{\text{N-heteroaryl}}-\text{W}-\text{R}_{\text{N-1-heterocycle}},$

(I) $\text{R}_{\text{N-heterocycle}}-\text{W}-\text{R}_{\text{N-aryl}},$ wherein $\text{R}_{\text{N-heterocycle}}$ is the same as

$\text{R}_{1-\text{heterocycle}},$ and $\text{R}_{1-\text{heterocycle}}$ is as defined above, and $\text{R}_{\text{N-aryl}}$ is as defined above,

25

(J) $\text{R}_{\text{N-heterocycle}}-\text{W}-\text{R}_{\text{N-heteroaryl}},$ and

(K) $\text{R}_{\text{N-heterocycle}}-\text{W}-\text{R}_{\text{N-1-heterocycle}},$

where W is

(29) $-(\text{CH}_2)_{0-4}-,$

(30) $-\text{O}-,$

30

(31) $-\text{S}(\text{O})_{0-2}-,$

(32) $-\text{N}(\text{R}_{\text{N}-5})-$ where $\text{R}_{\text{N}-5}$ is as defined

above, or

(5) $-\text{CO}-;$

(II) $-\text{CO}-(\text{C}_1\text{-C}_{10}\text{ alkyl})$ where alkyl is optionally substituted with one three substitutents selected from the group consisting of:

- (A) $-\text{OH}$,
(B) $-\text{C}_1\text{-C}_6\text{ alkoxy}$,
5 (C) $-\text{C}_1\text{-C}_6\text{ thioalkoxy}$,
(D) $-\text{CO-O-R}_{\text{N-8}}$ where $\text{R}_{\text{N-8}}$ is $-\text{H}$, $\text{C}_1\text{-C}_6\text{ alkyl}$ or $-\text{phenyl}$,
(E) $-\text{CO-NR}_{\text{N-2}}\text{R}_{\text{N-3}}$ where $\text{R}_{\text{N-2}}$ and $\text{R}_{\text{N-3}}$ are the same or different and are as defined above,
(F) $-\text{CO-R}_{\text{N-4}}$ where $\text{R}_{\text{N-4}}$ is as defined above,
10 (G) $-\text{SO}_2-(\text{C}_1\text{-C}_8\text{ alkyl})$,
(H) $-\text{SO}_2\text{-NR}_{\text{N-2}}\text{R}_{\text{N-3}}$ where $\text{R}_{\text{N-2}}$ and $\text{R}_{\text{N-3}}$ are the same or different and are as defined above,
(I) $-\text{NH-CO}-(\text{C}_1\text{-C}_6\text{ alkyl})$,
(J) $-\text{NH-CO-O-R}_{\text{N-8}}$ where $\text{R}_{\text{N-8}}$ is as defined above,
15 (K) $-\text{NR}_{\text{N-2}}\text{R}_{\text{N-3}}$ where $\text{R}_{\text{N-2}}$ and $\text{R}_{\text{N-3}}$ are the same or different and are as defined above,
(L) $-\text{R}_{\text{N-4}}$ where $\text{R}_{\text{N-4}}$ is as defined above,
(M) $-\text{O-CO}-(\text{C}_1\text{-C}_6\text{ alkyl})$,
(N) $-\text{O-CO-NR}_{\text{N-8}}\text{R}_{\text{N-8}}$ where $\text{R}_{\text{N-8}}$ are the same or different
20 and are as defined above,
(O) $-\text{O}-(\text{C}_1\text{-C}_5\text{ alkyl})\text{-COOH}$,
(P) $-\text{O}-(\text{C}_1\text{-C}_6\text{ alkyl})$ optionally substituted with one, two, or three of $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, or $-\text{I}$,
(Q) $-\text{NH-SO}_2-(\text{C}_1\text{-C}_6\text{ alkyl})$, and
25 (R) $-\text{F}$, or $-\text{Cl}$,
(III) $-\text{CO}-(\text{C}_1\text{-C}_6\text{ alkyl})\text{-O}-(\text{C}_1\text{-C}_6\text{ alkyl})$ where alkyl is optionally substituted with one, two, or three substitutents selected from the group consisting of:

- (A) $-\text{OH}$,
30 (B) $-\text{C}_1\text{-C}_6\text{ alkoxy}$,
(C) $-\text{C}_1\text{-C}_6\text{ thioalkoxy}$,
(D) $-\text{CO-O-R}_{\text{N-8}}$ where $\text{R}_{\text{N-8}}$ is $-\text{H}$, $\text{C}_1\text{-C}_6\text{ alkyl}$ or $-\text{phenyl}$,

- (E) $-\text{CO}-\text{NR}_{\text{N-2}}\text{R}_{\text{N-3}}$ where $\text{R}_{\text{N-2}}$ and $\text{R}_{\text{N-3}}$ are the same or different and are as defined above,
- (F) $-\text{CO}-\text{R}_{\text{N-4}}$ where $\text{R}_{\text{N-4}}$ is as defined above,
- (G) $-\text{SO}_2-(\text{C}_1-\text{C}_8 \text{ alkyl})$,
- 5 (H) $-\text{SO}_2-\text{NR}_{\text{N-2}}\text{R}_{\text{N-3}}$ where $\text{R}_{\text{N-2}}$ and $\text{R}_{\text{N-3}}$ are the same or different and are as defined above,
- (I) $-\text{NH}-\text{CO}-(\text{C}_1-\text{C}_6 \text{ alkyl})$,
- (J) $-\text{NH}-\text{CO}-\text{O}-\text{R}_{\text{N-8}}$ where $\text{R}_{\text{N-8}}$ is as defined above,
- (K) $-\text{NR}_{\text{N-2}}\text{R}_{\text{N-3}}$ where $\text{R}_{\text{N-2}}$ and $\text{R}_{\text{N-3}}$ are the same or different
- 10 and are as defined above,
- (L) $-\text{R}_{\text{N-4}}$ where $\text{R}_{\text{N-4}}$ is as defined above,
- (M) $-\text{O}-\text{CO}-(\text{C}_1-\text{C}_6 \text{ alkyl})$,
- (N) $-\text{O}-\text{CO}-\text{NR}_{\text{N-8}}\text{R}_{\text{N-8}}$ where $\text{R}_{\text{N-8}}$ are the same or different and are as defined above,
- 15 (O) $-\text{O}-(\text{C}_1-\text{C}_5 \text{ alkyl})-\text{COOH}$,
- (P) $-\text{O}-(\text{C}_1-\text{C}_6 \text{ alkyl})$ optionally substituted with one, two, or three of $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, or $-\text{I}$,
- (Q) $-\text{NH}-\text{SO}_2-(\text{C}_1-\text{C}_6 \text{ alkyl})$, and
- (R) $-\text{F}$, or $-\text{Cl}$,
- 20 (IV) $-\text{CO}-(\text{C}_1-\text{C}_6 \text{ alkyl})-\text{S}-(\text{C}_1-\text{C}_6 \text{ alkyl})$ where alkyl is optionally substituted with one, two, or three of substituents selected from the group consisting of:
- (A) $-\text{OH}$,
- (B) $-\text{C}_1-\text{C}_6 \text{ alkoxy}$,
- 25 (C) $-\text{C}_1-\text{C}_6 \text{ thioalkoxy}$,
- (D) $-\text{CO}-\text{O}-\text{R}_{\text{N-8}}$ where $\text{R}_{\text{N-8}}$ is as defined above,
- (E) $-\text{CO}-\text{NR}_{\text{N-2}}\text{R}_{\text{N-3}}$ where $\text{R}_{\text{N-2}}$ and $\text{R}_{\text{N-3}}$ are the same or different and are as defined above,
- (F) $-\text{CO}-\text{R}_{\text{N-4}}$ where $\text{R}_{\text{N-4}}$ is as defined above,
- 30 (G) $-\text{SO}_2-(\text{C}_1-\text{C}_8 \text{ alkyl})$,
- (H) $-\text{SO}_2-\text{NR}_{\text{N-2}}\text{R}_{\text{N-3}}$ where $\text{R}_{\text{N-2}}$ and $\text{R}_{\text{N-3}}$ are the same or different and are as defined above,
- (I) $-\text{NH}-\text{CO}-(\text{C}_1-\text{C}_6 \text{ alkyl})$,

(J) $-\text{NH}-\text{CO}-\text{O}-\text{R}_{\text{N}-8}$ where $\text{R}_{\text{N}-8}$ is as defined above,

(K) $-\text{NR}_{\text{N}-2}\text{R}_{\text{N}-3}$ where $\text{R}_{\text{N}-2}$ and $\text{R}_{\text{N}-3}$ are the same or different

and are as defined above,

(L) $-\text{R}_{\text{N}-4}$ where $\text{R}_{\text{N}-4}$ is as defined above,

5 (M) $-\text{O}-\text{CO}-(\text{C}_1-\text{C}_6 \text{ alkyl})$,

(N) $-\text{O}-\text{CO}-\text{NR}_{\text{N}-8}\text{R}_{\text{N}-8}$ where $\text{R}_{\text{N}-8}$ are the same or different

and are as defined above,

(O) $-\text{O}-(\text{C}_1-\text{C}_5 \text{ alkyl})-\text{COOH}$,

(P) $-\text{O}-(\text{C}_1-\text{C}_6 \text{ alkyl optionally substituted with one, two, or$

10 three of $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, or $-\text{I}$),

(Q) $-\text{NH}-\text{SO}_2-(\text{C}_1-\text{C}_6 \text{ alkyl})$, and

(R) $-\text{F}$, or $-\text{Cl}$,

(V) $-\text{CO}-\text{CH}(-(\text{CH}_2)_{0-2}-\text{O}-\text{R}_{\text{N}-10})-(\text{CH}_2)_{0-2}-\text{R}_{\text{N-aryl}}/\text{R}_{\text{N-heteroaryl}}$ where

$\text{R}_{\text{N-aryl}}$ and $\text{R}_{\text{N-heteroaryl}}$ are as defined above, where $\text{R}_{\text{N}-10}$ is selected from the group

15 consisting of:

(A) $-\text{H}$,

(B) $\text{C}_1-\text{C}_6 \text{ alkyl}$,

(C) $\text{C}_3-\text{C}_7 \text{ cycloalkyl}$,

(D) $\text{C}_2-\text{C}_6 \text{ alkenyl}$ with one double bond,

20 (E) $\text{C}_2-\text{C}_6 \text{ alkynyl}$ with one triple bond,

(F) $\text{R}_{1-\text{aryl}}$ where $\text{R}_{1-\text{aryl}}$ is as defined above, and

(G) $\text{R}_{\text{N-heteroaryl}}$ where $\text{R}_{\text{N-heteroaryl}}$ is as defined above, or

(VI) $-\text{CO}-(\text{C}_3-\text{C}_8 \text{ cycloalkyl})$ where alkyl is optionally substituted

with one or two substituents selected from the group consisting of:

25 (A) $-(\text{CH}_2)_{0-4}-\text{OH}$,

(B) $-(\text{CH}_2)_{0-4}-\text{C}_1-\text{C}_6 \text{ alkoxy}$,

(C) $-(\text{CH}_2)_{0-4}-\text{C}_1-\text{C}_6 \text{ thioalkoxy}$,

(D) $-(\text{CH}_2)_{0-4}-\text{CO}-\text{O}-\text{R}_{\text{N}-8}$ where $\text{R}_{\text{N}-8}$ is $-\text{H}$, $\text{C}_1-\text{C}_6 \text{ alkyl}$ or

phenyl,

30 (E) $-(\text{CH}_2)_{0-4}-\text{CO}-\text{NR}_{\text{N}-2}\text{R}_{\text{N}-3}$ where $\text{R}_{\text{N}-2}$ and $\text{R}_{\text{N}-3}$ are the same or different and are as defined above,

(F) $-(\text{CH}_2)_{0-4}-\text{CO}-\text{R}_{\text{N}-4}$ where $\text{R}_{\text{N}-4}$ is as defined above,

(G) $-(\text{CH}_2)_{0-4}-\text{SO}_2-(\text{C}_1-\text{C}_8 \text{ alkyl})$,

(H) $-(\text{CH}_2)_{0-4}-\text{SO}_2-\text{NR}_{\text{N}-2}\text{R}_{\text{N}-3}$ where $\text{R}_{\text{N}-2}$ and $\text{R}_{\text{N}-3}$ are the same or different and are as defined above,

(I) $-(\text{CH}_2)_{0-4}-\text{NH}-\text{CO}-(\text{C}_1-\text{C}_6 \text{ alkyl})$,

(J) $-\text{NH}-\text{CO}-\text{O}-\text{R}_{\text{N}-8}$ where $\text{R}_{\text{N}-8}$ is as defined above,

5 (K) $-(\text{CH}_2)_{0-4}-\text{NR}_{\text{N}-2}\text{R}_{\text{N}-3}$ where $\text{R}_{\text{N}-2}$ and $\text{R}_{\text{N}-3}$ are the same or different and are as defined above,

(L) $-(\text{CH}_2)_{0-4}-\text{R}_{\text{N}-4}$ where $\text{R}_{\text{N}-4}$ is as defined above,

(M) $-\text{O}-\text{CO}-(\text{C}_1-\text{C}_6 \text{ alkyl})$,

(N) $-\text{O}-\text{CO}-\text{NR}_{\text{N}-8}\text{R}_{\text{N}-8}$ where $\text{R}_{\text{N}-8}$ are the same or different
10 and are as defined above,

(O) $-\text{O}-(\text{C}_1-\text{C}_5 \text{ alkyl})-\text{COOH}$,

(P) $-\text{O}-(\text{C}_1-\text{C}_6 \text{ alkyl})$ optionally substituted with one, two, or three of $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, or $-\text{I}$,

(Q) $-\text{NH}-\text{SO}_2-(\text{C}_1-\text{C}_6 \text{ alkyl})$, and

15 (R) $-\text{F}$, or $-\text{Cl}$;

where R_A is:

(I) C_1-C_{10} alkyl optionally substituted with one, two or three substituents selected from the group consisting of C_1-C_3 alkyl, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, $-\text{OH}$,
20 $-\text{SH}$, $-\text{C}\equiv\text{N}$, $-\text{CF}_3$, C_1-C_6 alkoxy, $-\text{O}-\text{phenyl}$, $-\text{NR}_{1-\text{a}}\text{R}_{1-\text{b}}$ where $\text{R}_{1-\text{a}}$ and $\text{R}_{1-\text{b}}$ are as defined above, $-\text{OC}=\text{O NR}_{1-\text{a}}\text{R}_{1-\text{b}}$ where $\text{R}_{1-\text{a}}$ and $\text{R}_{1-\text{b}}$ are as defined above, $-\text{S}(=\text{O})_{0-2}\text{R}_{1-\text{a}}$ where $\text{R}_{1-\text{a}}$ is as defined above, $-\text{NR}_{1-\text{a}}\text{C}=\text{O NR}_{1-\text{a}}\text{R}_{1-\text{b}}$ where $\text{R}_{1-\text{a}}$ and $\text{R}_{1-\text{b}}$ are as defined above, $-\text{C}=\text{O NR}_{1-\text{a}}\text{R}_{1-\text{b}}$ where $\text{R}_{1-\text{a}}$ and $\text{R}_{1-\text{b}}$ are as defined above, and $-\text{S}(=\text{O})_2 \text{NR}_{1-\text{a}}\text{R}_{1-\text{b}}$ where $\text{R}_{1-\text{a}}$ and $\text{R}_{1-\text{b}}$ are as defined above,

25 (II) $-(\text{CH}_2)_{0-3}-(\text{C}_3-\text{C}_8)$ cycloalkyl where cycloalkyl can be optionally substituted with one, two or three substituents selected from the group consisting of C_1-C_3 alkyl, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, $-\text{OH}$, $-\text{SH}$, $-\text{C}\equiv\text{N}$, $-\text{CF}_3$, C_1-C_6 alkoxy, $-\text{O}-\text{phenyl}$, $-\text{CO}-\text{OH}$, $-\text{CO}-\text{O}-(\text{C}_1-\text{C}_4 \text{ alkyl})$, and $-\text{NR}_{1-\text{a}}\text{R}_{1-\text{b}}$ where $\text{R}_{1-\text{a}}$ and $\text{R}_{1-\text{b}}$ are as defined above,

30 (III) $-(\text{CR}_{\text{A}-\text{x}}\text{R}_{\text{A}-\text{y}})_{0-4}-\text{R}_{\text{A}-\text{aryl}}$ where $\text{R}_{\text{A}-\text{x}}$ and $\text{R}_{\text{A}-\text{y}}$ are

(A) $-\text{H}$,

(B) C_1-C_4 alkyl optionally substituted with one or two $-\text{OH}$,

(C) C₁-C₄ alkoxy optionally substituted with one, two, or three of -F,

(D) -(CH₂)₀₋₄-C₃-C₇ cycloalkyl,

(E) C₂-C₆ alkenyl containing one or two double bonds,

5 (F) C₂-C₆ alkynyl containing one or two triple bonds, or

(G) phenyl,

and where R_{A-x} and R_{A-y} are taken together with the carbon to which they are attached to form a carbocycle of three, four, five, six or seven carbon atoms, optionally where one carbon atom is replaced by a heteroatom selected from the

10 group consisting of -O-, -S-, -SO₂-, and -NR_{N-2}- and R_{A-aryl} is the same as R_{N-aryl},

(IV) -(CR_{A-x}R_{A-y})₀₋₄-R_{A-heteroaryl} where R_{A-heteroaryl} is the same as R_{N-heteroaryl} and R_{A-x} and R_{A-y} are as defined above,

(V) -(CR_{A-x}R_{A-y})₀₋₄-R_{A-aryl}-R_{A-aryl} where R_{A-aryl}, R_{A-x} and R_{A-y} are as defined above,

15 (VI) -(CR_{A-x}R_{A-y})₀₋₄-R_{A-aryl}-R_{A-heteroaryl} where R_{A-aryl}, R_{A-heteroaryl}, R_{A-x} and R_{A-y} are as defined above,

(VII) -(CR_{A-x}R_{A-y})₀₋₄-R_{A-heteroaryl}-R_{A-aryl} where R_{A-heteroaryl}, R_{A-aryl}, R_{A-x} and R_{A-y} are as defined above,

20 (VIII) -(CR_{A-x}R_{A-y})₀₋₄-R_{A-heteroaryl}-R_{A-heteroaryl} where R_{A-heteroaryl}, R_{A-x} and R_{A-y} are as defined above,

(IX) -(CR_{A-x}R_{A-y})₀₋₄-R_{A-aryl}-R_{A-heterocycle} where R_{A-heterocycle} is defined as R_{1-heterocycle}, and where R_{A-aryl}, R_{A-x} and R_{A-y} are as defined above,

(X) -(CR_{A-x}R_{A-y})₀₋₄-R_{A-heteroaryl}-R_{A-heterocycle} where R_{A-heteroaryl}, R_{A-heterocycle}, R_{A-x} and R_{A-y} are as defined above,

25 (XI) -(CR_{A-x}R_{A-y})₀₋₄-R_{A-heterocycle}-R_{A-aryl} where R_{A-heterocycle}, R_{A-aryl}, R_{A-x} and R_{A-y} are as defined above,

(XII) -(CR_{A-x}R_{A-y})₀₋₄-R_{A-heterocycle}-R_{A-heteroaryl} where R_{A-heterocycle}, R_{A-heteroaryl}, R_{A-x} and R_{A-y} are as defined above,

30 (XIII) -(CR_{A-x}R_{A-y})₀₋₄-R_{A-heterocycle}-R_{A-heterocycle} where R_{A-heterocycle}, R_{A-x} and R_{A-y} are as defined above,

(XIV) -(CR_{A-x}R_{A-y})₀₋₄-R_{A-heterocycle} where R_{A-heterocycle}, R_{A-x} and R_{A-y} are as defined above,

(XV) -[C(R_{A-1})(R_{A-2})]₁₋₃-CO-N-(R_{A-3})₂ where R_{A-1} and R_{A-2} are the same or different and are selected from the group consisting of:

(A) -H,

(B) -C₁-C₆ alkyl, optionally substituted with one, two or three substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(C) C₂-C₆ alkenyl with one or two double bonds, optionally substituted with one, two or three substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(D) C₂-C₆ alkynyl with one or two triple bonds, optionally substituted with one, two or three substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(E) -(CH₂)₁₋₂-S(O)₀₋₂-(C₁-C₆ alkyl),

(F) -(CH₂)₀₋₄-C₃-C₇ cycloalkyl, optionally substituted with one, two or three substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(G) -(C₁-C₄ alkyl)-R_{A'-aryl} where R_{A'-aryl} is as defined for R_{1-aryl},

(H) -(C₁-C₄ alkyl)-R_{A-heteroaryl} where R_{A-heteroaryl} is as defined above,

(I) -(C₁-C₄ alkyl)-R_{A-heterocycle} where R_{A-heterocycle} is as defined above,

(J) -R_{A-heteroaryl} where R_{A-heteroaryl} is as defined above,

(K) -R_{A-heterocycle} where R_{A-heterocycle} is as defined above,

(M) -(CH₂)₁₋₄-R_{A-4}-(CH₂)₀₋₄-R_{A'-aryl} where R_{A-4} is -O-, -S- or -NR_{A-5}- where R_{A-5} is C₁-C₆ alkyl, and where R_{A'-aryl} is defined above,

(N) -(CH₂)₁₋₄-R_{A-4}-(CH₂)₀₋₄-R_{A-heteroaryl} where R_{A-4} and R_{A-heteroaryl} are as defined above, and

(O) -R_{A'-aryl} where R_{A'-aryl} is as defined above,

and where R_{A-3} is the same or different and is:

(A) -H,

(B) $\text{C}_1\text{-C}_6$ alkyl optionally substituted with one, two or three substituents selected from the group consisting of $\text{C}_1\text{-C}_3$ alkyl, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, $-\text{OH}$, $-\text{SH}$, $-\text{C}\equiv\text{N}$, $-\text{CF}_3$, $\text{C}_1\text{-C}_6$ alkoxy, $-\text{O-phenyl}$, $-\text{NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

5 (C) $\text{C}_2\text{-C}_6$ alkenyl with one or two double bonds, optionally substituted with one, two or three substituents selected from the group consisting of $\text{C}_1\text{-C}_3$ alkyl, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, $-\text{OH}$, $-\text{SH}$, $-\text{C}\equiv\text{N}$, $-\text{CF}_3$, $\text{C}_1\text{-C}_6$ alkoxy, $-\text{O-phenyl}$, and $-\text{NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

(D) $\text{C}_2\text{-C}_6$ alkynyl with one or two triple bonds, optionally
10 substituted with one, two or three substituents selected from the group consisting of $\text{C}_1\text{-C}_3$ alkyl, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, $-\text{OH}$, $-\text{SH}$, $-\text{C}\equiv\text{N}$, $-\text{CF}_3$, $\text{C}_1\text{-C}_6$ alkoxy, $-\text{O-phenyl}$, and $-\text{NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

(E) $-(\text{CH}_2)_{0-4}\text{-C}_3\text{-C}_7$ cycloalkyl, optionally substituted with one, two or three substituents selected from the group consisting of $\text{C}_1\text{-C}_3$ alkyl, $-\text{F}$,
15 $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, $-\text{OH}$, $-\text{SH}$, $-\text{C}\equiv\text{N}$, $-\text{CF}_3$, $\text{C}_1\text{-C}_6$ alkoxy, $-\text{O-phenyl}$, and $-\text{NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

(F) $-\text{R}_{\text{A-aryl}}$ where $\text{R}_{\text{A-aryl}}$ is as defined above,

(G) $-\text{R}_{\text{A-heteroaryl}}$ where $\text{R}_{\text{A-heteroaryl}}$ is as defined above,

(H) $-\text{R}_{\text{A-heterocycle}}$ where $\text{R}_{\text{A-heterocycle}}$ is as defined above,

20 (I) $-(\text{C}_1\text{-C}_4 \text{ alkyl})\text{-R}_{\text{A-aryl}}$ where $\text{R}_{\text{A-aryl}}$ is as defined above,

(J) $-(\text{C}_1\text{-C}_4 \text{ alkyl})\text{-R}_{\text{A-heteroaryl}}$ where $\text{R}_{\text{A-heteroaryl}}$ is as defined above,

(K) $-(\text{C}_1\text{-C}_4 \text{ alkyl})\text{-R}_{\text{A-heterocycle}}$ where $\text{R}_{\text{A-heterocycle}}$ is as defined
25 above, or

(XVI) $-\text{CH}(\text{R}_{\text{A-aryl}})_2$ where $\text{R}_{\text{A-aryl}}$ are the same or different and are as defined above,

(XVII) $-\text{CH}(\text{R}_{\text{A-heteroaryl}})_2$ where $\text{R}_{\text{A-heteroaryl}}$ are the same or different and are as defined above,

30 (XVIII) $-\text{CH}(\text{R}_{\text{A-aryl}})(\text{R}_{\text{A-heteroaryl}})$ where $\text{R}_{\text{A-aryl}}$ and $\text{R}_{\text{A-heteroaryl}}$ are as defined above,

(XIX) $-\text{cyclopentyl}$, $-\text{cyclohexyl}$, or $-\text{cycloheptyl}$ ring fused to $\text{R}_{\text{A-aryl}}$, $\text{R}_{\text{A-heteroaryl}}$, $\text{R}_{\text{A-heterocycle}}$ where $\text{R}_{\text{A-aryl}}$ or $\text{R}_{\text{A-heteroaryl}}$ or $\text{R}_{\text{A-heterocycle}}$ are as defined above

where one carbon of cyclopentyl, cyclohexyl, or -cycloheptyl is optionally replaced with NH, $\text{NR}_{\text{N-5}}$, O, or $\text{S}(=\text{O})_{0-2}$, and where cyclopentyl, cyclohexyl, or -cycloheptyl can be optionally substituted with one or two $\text{-C}_1\text{-C}_3$ alkyl, -F , -OH , -SH , $\text{-C}\equiv\text{N}$, -CF_3 , $\text{C}_1\text{-C}_6$ alkoxy, $=\text{O}$, or $\text{-NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

5 (XX) $\text{C}_2\text{-C}_{10}$ alkenyl containing one or two double bonds optionally substituted with one, two or three substituents selected from the group consisting of $\text{C}_1\text{-C}_3$ alkyl, -F , -Cl , -Br , -I , -OH , -SH , $\text{-C}\equiv\text{N}$, -CF_3 , $\text{C}_1\text{-C}_6$ alkoxy, -O-phenyl , and $\text{-NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

(XXI) $\text{C}_2\text{-C}_{10}$ alkynyl containing one or two triple bonds optionally substituted with one, two or three substituents selected from the group consisting of $\text{C}_1\text{-C}_3$ alkyl, -F , -Cl , -Br , -I , -OH , -SH , $\text{-C}\equiv\text{N}$, -CF_3 , $\text{C}_1\text{-C}_6$ alkoxy, -O-phenyl , $\text{-NR}_{1-a}\text{R}_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

(XXI) $\text{-(CH}_2\text{)}_{0-1}\text{-CHR}_{\text{A-6}}\text{-(CH}_2\text{)}_{0-1}\text{-R}_{\text{A-aryl}}$ where $\text{R}_{\text{A-aryl}}$ is as defined above and $\text{R}_{\text{A-6}}$ is $\text{-(CH}_2\text{)}_{0-6}\text{-OH}$,

15 (XXII) $\text{-(CH}_2\text{)}_{0-1}\text{-CHR}_{\text{A-6}}\text{-(CH}_2\text{)}_{0-1}\text{-R}_{\text{A-heteroaryl}}$ where $\text{R}_{\text{A-heteroaryl}}$ and $\text{R}_{\text{A-6}}$ is as defined above,

(XXIII) $\text{-CH(R}_{\text{A-aryl}}\text{ or R}_{\text{A-heteroaryl}}\text{)-CO-O(C}_1\text{-C}_4\text{ alkyl)}$ where $\text{R}_{\text{A-aryl}}$ and $\text{R}_{\text{A-heteroaryl}}$ are as defined above,

(XXIV) $\text{-CH(CH}_2\text{-OH)-CH(OH)-micro-NO}_2$,

20 (XXV) $\text{(C}_1\text{-C}_6\text{ alkyl)-O-(C}_1\text{-C}_6\text{ alkyl)-OH}$,

(XXVII) $\text{-CH}_2\text{-NH-CH}_2\text{-CH(O-CH}_2\text{-CH}_3\text{)}_2$,

(XXVIII) -H ,

(XXIX) $\text{-(CH}_2\text{)}_{0-6}\text{-C(=NR}_{1-a}\text{)(NR}_{1-a}\text{R}_{1-b})$ where R_{1-a} and R_{1-b} are as defined above; or

25 (XXX)

$\text{-C=OC(HR}_6\text{)NHR}_7$, where R_6 and R_7 are as defined below,

-C=OR_7 , where R_7 is as defined below,

-C=OOR_7 , where R_7 is as defined below, or

-SOOR_7 where R_7 is as defined below,

30 wherein R_6 is:

hydrogen,

$\text{C}_1\text{-C}_3$ alkyl,

phenyl,

thioalkoxyalkyl,
alkyl substituted aryl,
cycloalkyl,
cycloalkylalkyl,
5 hydroxyalkyl,
alkoxyalkyl,
aryloxyalkyl,
haloalkyl,
carboxyalkyl,
10 alkoxycarbonylalkyl,
aminoalkyl,
(N-protected)aminoalkyl,
alkylaminoalkyl,
((N-protected)(alkyl)amino)alkyl,
15 dialkylaminoalkyl,
guanidinoalkyl,
lower alkenyl,
heterocyclic,
(heterocyclic)alkyl),
20 arylthioalkyl,
arylsulfonylalkyl,
(heterocyclic)thioalkyl,
(heterocyclic)sulfonylalkyl,
(heterocyclic)oxyalkyl,
25 arylalkoxyalkyl,
arylthioalkoxyalkyl,
arylalkylsulfonylalkyl,
(heterocyclic))alkoxyalkyl,
(heterocyclic)thioalkoxyalkyl,
30 (heterocyclic)alkylsulfonylalkyl,
cycloalkyloxyalkyl,
cycloalkylthioalkyl,
cycloalkylsulfonylalkyl,
cycloalkylalkoxyalkyl,

- 5 cycloalkylthioalkoxyalkyl,
cycloalkylalkylsulfonylalkyl,
aminocarbonyl,
alkylaminocarbonyl,
dialkylaminocarbonyl,
aroalkyl,
(heterocyclic)carbonylalkyl,
polyhydroxyalkyl,
aminocarbonylalkyl,
10 alkylaminocarbonylalkyl,
dialkylaminocarbonylalkyl,
aryloxyalkyl, or
alkylsulfonylalkyl,
wherein heterocyclic is pyridyl, thiazolyl, isothiazolyl,
15 oxazolyl, isoxazolyl, furanyl, thienyl, tetrahydrofuranyl, tetrahydrothienyl and
tetrahydro[2H]pyranyl and wherein the heterocycle is unsubstituted or substituted
with one to three substituents independently selected from hydroxy, halo, amino,
alkylamino, dialkylamino, alkoxy, polyalkoxy, haloalkyl, cycloalkyl,
cycloalkylalkyl, aryl, arylalkyl, COOH, -SO₃H, lower alkenyl or lower alkyl;
20 wherein R₇ is:
C₁ - C₃ alkyl,
phenyl,
thioalkoxyalkyl,
(aryl)alkyl,
25 cycloalkyl,
cycloalkylalkyl,
hydroxyalkyl,
alkoxyalkyl,
aryloxyalkyl,
30 haloalkyl,
carboxyalkyl,
alkoxycarbonylalkyl,
aminoalkyl,
(N-protected)aminocalkyl,

alkylaminoalkyl,
((N-protected)(alkyl)amino)alkyl,
dialkylaminoalkyl,
guanidinoalkyl,
5 lower alkenyl,
heterocyclic,
(heterocyclic)alkyl),
arylthioalkyl,
arylsulfonylalkyl,
10 (heterocyclic)thioalkyl,
(heterocyclic)sulfonylalkyl,
(heterocyclic)oxyalkyl,
arylalkoxyalkyl,
arylthioalkoxyalkyl,
15 arylalkylsulfonylalkyl,
(heterocyclic))alkoxyalkyl,
(heterocyclic)thioalkoxyalkyl,
(heterocyclic)alkylsulfonylalkyl,
cycloalkyloxyalkyl,
20 cycloalkylthioalkyl,
cycloalkylsulfonylalkyl,
cycloalkylalkoxyalkyl,
cycloalkylthioalkoxyalkyl,
cycloalkylalkylsulfonylalkyl,
25 aminocarbonyl,
alkylaminocarbonyl,
dialkylaminocarbonyl,
aroalkyl,
(heterocyclic)carbonylalkyl,
30 polyhydroxyalkyl,
aminocarbonylalkyl,
alkylaminocarbonylalkyl,
dialkylaminocarbonylalkyl,
aryloxyalkyl, or

alkylsulfonylalkyl,

wherein heterocyclic is pyridyl, thiazolyl, isothiazolyl, oxazolyl, isoxazolyl, furanyl, thienyl, tetrahydrofuranyl, tetrahydrothienyl and tetrahydro[2H]pyranyl and wherein the heterocycle is unsubstituted or substituted
 5 with one to three substituents independently selected from hydroxy, halo, amino, alkylamino, dialkylamino, alkoxy, polyalkoxy, haloalkyl, cycloalkyl, cycloalkylalkyl, aryl, arylalkyl, COOH, -SO₃H, lower alkenyl or lower alkyl;

where X is -N, or -O, with the proviso that when X is O, R_B is absent;

10 and when X is N,

R_B is:

(I)-C₁-C₁₀ alkyl optionally substituted with one, two or three substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as
 15 defined above, -OC=O NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above, -S(=O)₀₋₂ R_{1-a} where R_{1-a} is as defined above, -NR_{1-a}C=O NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above, -C=O NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above, and -S(=O)₂ NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(II) -(CH₂)₀₋₃-(C₃-C₈) cycloalkyl where cycloalkyl can be optionally
 20 substituted with one, two or three substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, -CO-OH, -CO-O-(C₁-C₄ alkyl), and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(III) -(CR_{B-x}R_{B-y})₀₋₄-R_{B-aryl} where R_{B-x} and R_{B-y} are

(A) -H,
 25 (B) C₁-C₄ alkyl optionally substituted with one or two -OH,
 (C) C₁-C₄ alkoxy optionally substituted with one, two, or three of -F,

(D) -(CH₂)₀₋₄-C₃-C₇ cycloalkyl,
 (E) C₂-C₆ alkenyl containing one or two double bonds,
 30 (F) C₂-C₆ alkynyl containing one or two triple bonds, or
 (G) phenyl,

and where R_{B-x} and R_{B-y} are taken together with the carbon to which they are attached to form a carbocycle of three, four, five, six or seven carbon

atoms, optionally where one carbon atom is replaced by a heteroatom selected from the group consisting of -O-, -S-, -SO₂-, and -NR_{N-2} where R_{N-2} is as defined above, and R_{B-aryl} is the same as R_{N-aryl} and is defined above

(IV) -(CR_{B-x}R_{B-y})₀₋₄-R_{B-heteroaryl} where R_{B-heteroaryl} is the same as R_N.

5 heteroaryl, R_{B-x}, and R_{B-y} are as defined above,

(V) -(CR_{B-x}R_{B-y})₀₋₄-R_{B-aryl}-R_{B-aryl} where R_{B-aryl}, R_{B-x}, and R_{B-y} are as defined above,

(VI) -(CR_{B-x}R_{B-y})₀₋₄-R_{B-aryl}-R_{B-heteroaryl} where R_{B-aryl}, R_{B-heteroaryl}, R_{B-x} and R_{B-y} are as defined above,

10 (VII) -(CR_{B-x}R_{B-y})₀₋₄-R_{B-heteroaryl}-R_{B-aryl} where R_{B-heteroaryl}, R_{B-aryl}, R_{B-x} and R_{B-y} are as defined above,

(VIII) -(CR_{B-x}R_{B-y})₀₋₄-R_{B-heteroaryl}-R_{B-heteroaryl} where R_{B-heteroaryl}, R_{B-x} and R_{B-y} are as defined above,

(IX) -(CR_{B-x}R_{B-y})₀₋₄-R_{B-aryl}-R_{B-heterocycle} where R_{B-heterocycle} is defined as 15 R_{1-heterocycle}, and where R_{B-aryl}, R_{B-x} and R_{B-y} are as defined above,

(X) -(CR_{B-x}R_{B-y})₀₋₄-R_{B-heteroaryl}-R_{B-heterocycle} where R_{B-heteroaryl}, R_{B-heterocycle}, R_{B-x} and R_{B-y} are as defined above,

(XI) -(CR_{B-x}R_{B-y})₀₋₄-R_{B-heterocycle}-R_{B-aryl} where R_{B-heterocycle}, R_{B-aryl}, R_{B-x} and R_{B-y} are as defined above,

20 (XII) -(CR_{B-x}R_{B-y})₀₋₄-R_{B-heterocycle}-R_{B-heteroaryl} where R_{B-heterocycle}, R_{B-heteroaryl}, R_{B-x} and R_{B-y} are as defined above,

(XIII) -(CR_{B-x}R_{B-y})₀₋₄-R_{B-heterocycle}-R_{B-heterocycle} where R_{B-heterocycle}, R_{B-x} and R_{B-y} are as defined above,

(XIV) -(CR_{B-x}R_{B-y})₀₋₄-R_{B-heterocycle} where R_{B-heterocycle}, R_{B-x} and R_{B-y} are 25 as defined above,

(XV) -[C(R_{B-1})(R_{B-2})]₁₋₃-CO-N-(R_{B-3})₂ where R_{B-1} and R_{B-2} are the same or different and are selected from the group consisting of:

(A) -H,

(B) -C₁-C₆ alkyl, optionally substituted with one, two or three 30 substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(C) C₂-C₆ alkenyl with one or two double bonds, optionally substituted with one, two or three substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

5 (D) C₂-C₆ alkynyl with one or two triple bonds, optionally substituted with one, two or three substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(E) -(CH₂)₁₋₂-S(O)₀₋₂-(C₁-C₆ alkyl),

10 (F) -(CH₂)₀₋₄-C₃-C₇ cycloalkyl, optionally substituted with one, two or three substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(G) -(C₁-C₄ alkyl)-R_{B'-aryl} where R_{B'-aryl} is as defined above for

15 R_{1-aryl},

(H) -(C₁-C₄ alkyl)-R_{B-heteroaryl} where R_{B-heteroaryl} is as defined above,

(I) -(C₁-C₄ alkyl)-R_{B-heterocycle} where R_{B-heterocycle} is as defined above,

20 (J) -R_{B-heteroaryl} where R_{B-heteroaryl} is as defined above,

(K) -R_{B-heterocycle} where R_{B-heterocycle} is as defined above,

(M) -(CH₂)₁₋₄-R_{B-4}-(CH₂)₀₋₄-R_{B'-aryl} where R_{B-4} is -O-, -S- or -NR_{B-5}- where R_{B-5} is C₁-C₆ alkyl, and where R_{B'-aryl} is defined above,

(N) -(CH₂)₁₋₄-R_{B-4}-(CH₂)₀₋₄-R_{B-heteroaryl} where R_{B-4} and R_{B-heteroaryl} are as defined above, and

(O) -R_{B'-aryl} where R_{B'-aryl} is as defined above,

and where R_{B-3} is the same or different and is:

(A) -H,

(B) -C₁-C₆ alkyl optionally substituted with one, two or three
30 substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(C) C₂-C₆ alkenyl with one or two double bonds, optionally substituted with one, two or three substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

5 (D) C₂-C₆ alkynyl with one or two triple bonds, optionally substituted with one, two or three substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(E) -(CH₂)₀₋₄-C₃-C₇ cycloalkyl, optionally substituted with
10 one, two or three substituents selected from the group consisting of C₁-C₃ alkyl, -F, -Cl, -Br, -I, -OH, -SH, -C≡N, -CF₃, C₁-C₆ alkoxy, -O-phenyl, and -NR_{1-a}R_{1-b} where R_{1-a} and R_{1-b} are as defined above,

(F) -R_{B'-aryl} where R_{B'-aryl} is as defined above,

(G) -R_{B-heteroaryl} where R_{B-heteroaryl} is as defined above,

15 (H) -R_{B-heterocycle} where R_{B-heterocycle} is as defined above,

(I) -(C₁-C₄ alkyl)-R_{B'-aryl} where R_{B'-aryl} is as defined above,

(J) -(C₁-C₄ alkyl)-R_{B-heteroaryl} where R_{B-heteroaryl} is as defined above,

20 (K) -(C₁-C₄ alkyl)-R_{B-heterocycle} where R_{B-heterocycle} is as defined above, or

(XVI) -CH(R_{B-aryl})₂ where R_{B-aryl} are the same or different and are as defined above,

(XVII) -CH(R_{B-heteroaryl})₂ where R_{B-heteroaryl} are the same or different
25 and are as defined above,

(XVIII) -CH(R_{B-aryl})(R_{B-heteroaryl}) where R_{B-aryl} and R_{B-heteroaryl} are as defined above,

(XIX) -cyclopentyl, -cyclohexyl, or -cycloheptyl ring fused to R_{B-aryl} or R_{B-heteroaryl} or R_{B-heterocycle} where R_{B-aryl} or R_{B-heteroaryl} or R_{B-heterocycle} are as defined
30 above where one carbon of cyclopentyl, cyclohexyl, or -cycloheptyl is optionally replaced with NH, NR_{N-5}, O, or S(=O)₀₋₂, and where cyclopentyl, cyclohexyl, or -cycloheptyl can be optionally substituted with one or two -C₁-C₃ alkyl, -F, -OH, -

SH, $-C\equiv N$, $-CF_3$, C_1-C_6 alkoxy, $=O$, or $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

(XX) C_2-C_{10} alkenyl containing one or two double bonds optionally substituted with one, two or three substituents selected from the group consisting of
 5 C_1-C_3 alkyl, $-F$, $-Cl$, $-Br$, $-I$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, C_1-C_6 alkoxy, $-O$ -phenyl, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

(XXI) C_2-C_{10} alkynyl containing one or two triple bonds optionally substituted with one, two or three substituents selected from the group consisting of
 10 C_1-C_3 alkyl, $-F$, $-Cl$, $-Br$, $-I$, $-OH$, $-SH$, $-C\equiv N$, $-CF_3$, C_1-C_6 alkoxy, $-O$ -phenyl, and $-NR_{1-a}R_{1-b}$ where R_{1-a} and R_{1-b} are as defined above,

(XXI) $-(CH_2)_{0-1}-CHR_{C-6}-(CH_2)_{0-1}-R_{B-aryl}$ where R_{B-aryl} is as defined above and R_{C-6} is $-(CH_2)_{0-6}-OH$,

(XXII) $-(CH_2)_{0-1}-CHR_{B-6}-(CH_2)_{0-1}-R_{B-heteroaryl}$ where $R_{B-heteroaryl}$ and R_{C-6} is as defined above,

15 (XXIII) $-\text{CH}(-R_{B-aryl} \text{ or } R_{B-heteroaryl})-\text{CO}-O(C_1-C_4 \text{ alkyl})$ where R_{B-aryl} and $R_{B-heteroaryl}$ are as defined above,

(XXIV) $-\text{CH}(-CH_2-OH)-\text{CH}(-OH)-\text{micro-NO}_2$,

(XXV) $(C_1-C_6 \text{ alkyl})-O-(C_1-C_6 \text{ alkyl})-OH$,

(XXVII) $-\text{CH}_2-\text{NH}-\text{CH}_2-\text{CH}(-O-\text{CH}_2-\text{CH}_3)_2$,

20 (XXVIII) $-H$, or

(XXIX) $-(CH_2)_{0-6}-C(=NR_{1-a})(NR_{1-a}R_{1-b})$ where R_{1-a} and R_{1-b} are as defined above.

96. A method of treatment according to claim 95,

25 where R_1 is:

$-(CH_2)_{0-1}-(R_{1-aryl})$, or

$-(CH_2)_{n1}-(R_{1-heteroaryl})$;

where R_N is:

$R_{N-1}-X_N$, where X_N is selected from the group consisting of:

30 $-\text{CO}-$, and

$-\text{SO}_2-$,

where R_{N-1} is selected from the group consisting of:

$-R_{N-aryl}$, and

-R_{N-heteroaryl}, or

-CO-CH(-(CH₂)₀₋₂-O-R_{N-10})-(CH₂)₀₋₂-R_{N-aryl}/R_{N-heteroaryl});

where R_A is:

-C₁-C₈ alkyl,

5 -(CH₂)₀₋₃-(C₃-C₇) cycloalkyl,

-(CR_{A-x}R_{A-y})₀₋₄-R_{A-aryl},

-(CR_{A-x}R_{A-y})₀₋₄-R_{A-heteroaryl},

-(CR_{A-x}R_{A-y})₀₋₄-R_{A-heterocycle},

-cyclopentyl or -cyclohexyl ring fused to R_{A-aryl} or R_{A-heteroaryl} or R_{A-}

10 heterocycle;

where X is -N or -O, with the proviso that when X is O, R_B is absent;

and when X is N,

R_B is:

-C₁-C₈ alkyl,

15 -(CH₂)₀₋₃-(C₃-C₇) cycloalkyl,

-(CR_{A-x}R_{A-y})₀₋₄-R_{A-aryl},

-(CR_{A-x}R_{A-y})₀₋₄-R_{A-heteroaryl},

-(CR_{A-x}R_{A-y})₀₋₄-R_{A-heterocycle},

-cyclopentyl or -cyclohexyl ring fused to R_{A-aryl} or R_{A-heteroaryl} or R_{A-}

20 heterocycle.

97. A method of treatment according to claim 96,

where R₁ is:

-(CH₂)-(R_{1-aryl}), or

25 -(CH₂)-(R_{1-heteroaryl});

where R₂ is -H;

where R₃ is -H;

where R_N is:

R_{N-1}-X_N where X_N is:

30 -CO-,

where R_{N-1} is selected from the group consisting of:

-R_{N-aryl}, and

-R_{N-heteroaryl};

where R_A is:

- 5 -C₁-C₈ alkyl,
 -(CH₂)₀₋₃-(C₃-C₇) cycloalkyl,
 -(CR_{A-x}R_{A-y})₀₋₄-R_{A-aryl},
 -(CR_{A-x}R_{A-y})₀₋₄-R_{A-heteroaryl},
 -(CR_{A-x}R_{A-y})₀₋₄-R_{A-heterocycle},
 -cyclopentyl or -cyclohexyl ring fused to R_{A-aryl} or R_{A-heteroaryl} or R_{A-}
 heterocycle;

where X is -N or -O, with the proviso that when X is O, R_B is absent;
 and when X is N,

- 10 R_B is:
 -C₁-C₈ alkyl,
 -(CH₂)₀₋₃-(C₃-C₇) cycloalkyl,
 -(CR_{B-x}R_{B-y})₀₋₄-R_{B-aryl},
 -(CR_{B-x}R_{B-y})₀₋₄-R_{B-heteroaryl},
 15 -(CR_{B-x}R_{B-y})₀₋₄-R_{B-heterocycle},
 -cyclopentyl or -cyclohexyl ring fused to R_{B-aryl} or R_{B-heteroaryl} or R_{B-}
 heterocycle.

98. A method of treatment according to claim 97

- 20 where R_A is:
 -(CR_{A-x}R_{A-y})₀₋₄-R_{A-aryl},
 -(CR_{A-x}R_{A-y})₀₋₄-R_{A-heteroaryl}, or
 -cyclopentyl or -cyclohexyl ring fused to a R_{A-aryl} or R_{A-heteroaryl} or R_{A-}
 heterocycle; and

- 25 where R_B is:
 -(CR_{B-x}R_{B-y})₀₋₄-R_{B-aryl},
 -(CR_{B-x}R_{B-y})₀₋₄-R_{B-heteroaryl}, or
 -cyclopentyl or -cyclohexyl ring fused to R_{B-aryl} or R_{B-heteroaryl} or R_{B-}
 heterocycle.

30

99. A method of treatment according to claim 95 where R₁ is

-(CH₂)-(R_{1-aryl}) where R_{1-aryl} is phenyl.

100. A method of treatment according to claim 95, where R₁ is

$-(\text{CH}_2)-(\text{R}_{1\text{-aryl}})$ where $\text{R}_{1\text{-aryl}}$ is phenyl substituted with two $-\text{F}$.

101. A method of treatment according to claim 100 where the $-\text{F}$ substitutions are at the 3- and 5- positions.

5

102. A method of treatment according to claim 95 where R_2 is $-\text{H}$.

103. A method of treatment according to claim 95 where R_3 is $-\text{H}$.

10 104. A method of treatment according to claim 95 where R_N is

$\text{R}_{\text{N-1}}-\text{X}_\text{N}-$, where X_N is $-\text{CO}-$, where $\text{R}_{\text{N-1}}$ is $\text{R}_{\text{N-aryl}}$ where $\text{R}_{\text{N-aryl}}$ is phenyl substituted with one $-\text{CO}-\text{NR}_{\text{N-2}}\text{R}_{\text{N-3}}$ where the substitution on phenyl is 1,3-.

15 105. A method of treatment according to claim 104 where $\text{R}_{\text{N-2}}$ and $\text{R}_{\text{N-3}}$ are the same and are C_3 alkyl.

106. A method of treatment according to claim 95 where R_N is

$\text{R}_{\text{N-1}}-\text{X}_\text{N}-$ where X_N is $-\text{CO}-$, and where $\text{R}_{\text{N-1}}$ is $\text{R}_{\text{N-aryl}}$ where $\text{R}_{\text{N-aryl}}$ is phenyl substituted with one C_1 alkyl and with one $-\text{CO}-\text{NR}_{\text{N-2}}\text{R}_{\text{N-3}}$ where the substitution on the phenyl is 1,3,5-.

20

107. A method of treatment according to claim 106 where $\text{R}_{\text{N-2}}$ and $\text{R}_{\text{N-3}}$ are the same and are C_3 alkyl.

25 108. A method of treatment according to claim 95 where R_N is


$\text{R}_{\text{N-1}}-\text{X}_\text{N}-$, where X_N is $-\text{CO}-$, and where $\text{R}_{\text{N-1}}$ is $\text{R}_{\text{N-heteroaryl}}$ where $\text{R}_{\text{N-heteroaryl}}$ is substituted with one $-\text{CO}-\text{NR}_{\text{N-2}}\text{R}_{\text{N-3}}$.

109. A method of treatment according to claim 108 where $\text{R}_{\text{N-2}}$ and $\text{R}_{\text{N-3}}$ are the same and are $-\text{C}_3$ alkyl.

30

110. A method of treatment according to claim 95 where R_A and R_B are each independently:

$-(\text{CR}_{\text{A-x}}\text{R}_{\text{A-y}})_{0-4}-\text{R}_{\text{A-aryl}}$ where $\text{R}_{\text{A-aryl}}$ is phenyl,



-cyclopentyl or -cyclohexyl ring fused to a $\text{R}_{\text{A-aryl}}$ or $\text{R}_{\text{A-heteroaryl}}$ or $\text{R}_{\text{A-heterocycle}}$.

5 111. The method of claim 95, wherein said beta-secretase is exposed to said compound *in vitro*.


112. The method of claim 95, wherein said beta-secretase is exposed to said compound in a cell.

10

113. The method of claim 95, wherein said cell is in an animal.

114. The method of claim 113, wherein said animal is a human.

15 115. A method for inhibiting cleavage of amyloid precursor protein (APP), in a reaction mixture, at a site between Met596 and Asp597, numbered for the APP-695 amino acid isotype; or at a corresponding site of an isotype or mutant thereof, comprising exposing said reaction mixture to an effective inhibitory amount of a compound of formula XV



20

where R_1 , R_2 , R_3 , R_{N} , R_{A} , R_{B} , and X are as defined in claim 1.

116. The method of claim 115, wherein said cleavage site is between Met652 and Asp653, numbered for the APP-751 isotype; between Met 671 and Asp 672, numbered for the APP-770 isotype; between Leu596 and Asp597 of the APP-695 Swedish Mutation; between Leu652 and Asp653 of the APP-751 Swedish Mutation; or between Leu671 and Asp672 of the APP-770 Swedish Mutation.

117. The method of claim 115, wherein said reaction mixture is exposed *in vitro*.


30

118. The method of claim 115, wherein said reaction mixture is exposed in a cell.

119. The method of claim 118, wherein said cell is an animal cell.

5 120. The method of claim 119, wherein said cell is a human cell.

121. A method for inhibiting production of amyloid beta peptide (A beta) in a cell, comprising administering to said cell an effective inhibitory amount of a compound of the formula XV




where R_1 , R_2 , R_3 , R_N , R_A , R_B , and X are as defined in claim 1.

122. The method of claim 121, wherein said administering is to an animal.

15 123. The method of claim 122, wherein said administering is to a human.


124. A method for inhibiting the production of beta-amyloid plaque in an animal, comprising administering to said animal an effective inhibitory amount of a compound of the formula XV



where R_1 , R_2 , R_3 , R_N , R_A , R_B , and X are as defined in claim 1.

125. The method of claim 124, wherein said animal is a human.

126. A method for treating or preventing a disease characterized by beta-amyloid deposits in the brain comprising administering to a patient an effective therapeutic amount of a hydroxyethylene compound of the formula XV



5 where $R_1, R_2, R_3, R_N, R_A, R_B$, and X are as defined in claim 1.

127. The method of claim 126, wherein said therapeutic amount is in the range of from about 0.1 to about 1000 mg/day.

10 128. The method of claim 126, wherein said therapeutic amount is in the range of
from about 15 to about 1500 mg/day.

129. The method of claim 128, wherein said therapeutic amount is in the range of from about 1 to about 100 mg/day.

15


130. The method of claim 129, wherein said therapeutic amount is in the range of from about 5 to about 50 mg/day.

131. The method of claim 126, wherein said disease is Alzheimer's disease.

20


132. The method of claim 126, wherein said disease is Mild Cognitive Impairment, Down's Syndrome, or Hereditary Cerebral Hemorrhage with Amyloidosis of the Dutch Type.

25 133. A composition comprising beta-secretase complexed with a compound of the
formula XV



where R_1 , R_2 , R_3 , R_N , R_A , R_B , and X are as defined in claim 1.

134. A method for producing a beta-secretase complex comprising: exposing beta-
5 secretase to a compound of the formula XV




where R_1 , R_2 , R_3 , R_N , R_A , R_B , and X are as defined in claim 1, or a pharmaceutically acceptable salt thereof in a reaction mixture under conditions suitable for the production of said complex.

10

135. The method of claim 134, where said exposing is *in vitro*.

136. The method of claim 133, wherein said reaction mixture is a cell.

- 15 137. A kit comprising component parts capable of being assembled,
wherein at least one component part comprises a compound of formula XV




where R_1 , R_2 , R_3 , R_N , R_A , R_B , and X are as defined in claim 1, enclosed in a container.

20

138. The kit of claim 137, wherein said compound is lyophilized and at least one further component part comprises a diluent.

139. A kit comprising a plurality of containers, each container comprising one or more unit dose of a compound of formula XV



where R_1 , R_2 , R_3 , R_N , R_A , R_B , and X are as defined in claim 1.


5

140. The kit of claim 139, wherein each container is adapted for oral delivery and comprises a tablet, gel, or capsule.

141. The kit of claim 140, wherein each container is adapted for parenteral
10 delivery and comprises a depot product, syringe, ampoule, or vial.

142. The kit of claim 141, wherein each container is adapted for topical delivery and comprises a patch, medipad, ointment, or cream.


15 143. A kit comprising a compound of formula XV



where R_1 , R_2 , R_3 , R_N , R_A , R_B , and X are as defined in claim 1;

and one or more therapeutic agent selected from the group consisting of an antioxidant, an anti-inflammatory, a gamma secretase inhibitor, a neurotrophic agent,
20 an acetylcholinesterase inhibitor, a statin, an A beta peptide, and an anti-A beta antibody.


144. A composition comprising
a compound of formula XV



where R_1 , R_2 , R_3 , R_N , R_A , R_B , and X are as defined in claim 1;
and an inert diluent or edible carrier.


5 145. The composition of claim 144, wherein said carrier is an oil.

146. A composition comprising
a compound of formula XV




10 where R_1 , R_2 , R_3 , R_N , R_A , R_B , and X are as defined in claim 1; and
and a binder, excipient, disintegrating agent, lubricant, or gildant.

147. A composition comprising
a compound of formula XV



15 where R_1 , R_2 , R_3 , R_N , R_A , R_B , and X are as defined in claim 1, disposed in a cream,
ointment, or patch.

148. Use of a substituted amine of formula (XV)



where R_1 , R_2 , R_3 , R_N , R_A , R_B , and X are as defined in claim 1;

or pharmaceutically acceptable salts thereof for the manufacture of a

- 5 medicament for use in treating a patient who has, or in preventing a patient from getting, a disease or condition selected from the group consisting of Alzheimer's disease, for helping prevent or delay the onset of Alzheimer's disease, for treating patients with mild cognitive impairment (MCI) and preventing or delaying the onset of Alzheimer's disease in those who would progress from MCI to AD, for treating
- 10 Down's syndrome, for treating humans who have Hereditary Cerebral Hemorrhage with Amyloidosis of the Dutch-Type, for treating cerebral amyloid angiopathy and preventing its potential consequences, i.e. single and recurrent lobar hemorrhages, for treating other degenerative dementias, including dementias of mixed vascular and degenerative origin, dementia associated with Parkinson's disease, dementia
- 15 associated with progressive supranuclear palsy, dementia associated with cortical basal degeneration, diffuse Lewy body type of Alzheimer's disease.

149. Use of a substituted amine of formula (XV) according to claim 148 where the disease is Alzheimer's disease.

20

150. Use of a substituted amine of formula (XV) according to claim 148 where the method is helping prevent or delay the onset of Alzheimer's disease,

151. Use of a substituted amine of formula (XV) according to claim 148 where the

25 disease is mild cognitive impairment.

152. Use of a substituted amine of formula (XV) according to claim 148 where the disease is Down's syndrome.

153. Use of a substituted amine of formula (XV) according to claim 148 where the disease is Hereditary Cerebral Hemorrhage with Amyloidosis of the Dutch-Type.

154. Use of a substituted amine of formula (XV) according to claim 148 where the disease is cerebral amyloid angiopathy.

155. Use of a substituted amine of formula (XV) according to claim 148 where the disease is degenerative dementias.

156. Use of a substituted amine of formula (XV) according to claim 148 where the disease is diffuse Lewy body type of Alzheimer's disease.

157. Use of a substituted amine of formula (XV) according to claim 148:

where R_1 is:

15 $-(CH_2)_{0-1}-(R_1\text{-aryl})$
 $-(CH_2)_{n1}-(R_1\text{-heteroaryl});$

where R_N is:

$R_{N-1}-X_N$, where X_N is selected from the group consisting of:

20 $-\text{CO}-$, and
 $-\text{SO}_2-$,

where R_{N-1} is selected from the group consisting of:

$-R_{N\text{-aryl}}$, and
 $-R_{N\text{-heteroaryl}}$, or
 $-\text{CO}-\text{CH}(-(CH_2)_{0-2}-\text{O}-R_{N-10})-(CH_2)_{0-2}-R_{N\text{-aryl}}/R_{N\text{-heteroaryl}});$ and

25 where R_A and R_B are each independently:

$-\text{C}_1-\text{C}_8$ alkyl,
 $-(CH_2)_{0-3}-(\text{C}_3-\text{C}_7)$ cycloalkyl,
 $-(\text{C}R_{A-x}R_{A-y})_{0-4}-R_{A\text{-aryl}}$,
 $-(\text{C}R_{A-x}R_{A-y})_{0-4}-R_{A\text{-heteroaryl}}$,
30 $-(\text{C}R_{A-x}R_{A-y})_{0-4}-R_{A\text{-heterocycle}}$,
 $-\text{cyclopentyl}$ or $-\text{cyclohexyl}$ ring fused to $R_{A\text{-aryl}}$ or $R_{A\text{-heteroaryl}}$ or $R_{A\text{-heterocycle}}$; and

where X is:

$-\text{N}$, or

-O, with the proviso that if X is O, R_B is absent.

158. Use of a substituted amine of formula (XV) according to claim 148:

where R₁ is:

5 -(CH₂)-(R_{1-aryl}), or
 -(CH₂)-(R_{1-heteroaryl});

where R₂ is -H;

where R₃ is -H;

where R_N is:

10 R_{N-1}-X_N- where X_N is:

-CO-,

where R_{N-1} is selected from the group consisting of:

-R_{N-aryl}, and

-R_{N-heteroaryl};

15 where R_A and R_B are each independently:

-(CH₂)₀₋₃-(C₃-C₇) cycloalkyl,

-(CR_{A-x}R_{A-y})₀₋₄-R_{A-aryl},

-(CR_{A-x}R_{A-y})₀₋₄-R_{A-heteroaryl},

-(CR_{A-x}R_{A-y})₀₋₄-R_{A-heterocycle}, or

20 -cyclopentyl or -cyclohexyl ring fused to a R_{A-aryl} or R_{A-heteroaryl} or R_{A-}

heterocycle; and

where X is:

-N, or

-O with the proviso that if X is O, R_B is absent.

25 159. Use of a substituted amine of formula (XV) according to claim 171 where R_A
 and R_B are each independently:

-(CR_{C-x}R_{C-y})₀₋₄-R_{C-aryl},

-(CR_{C-x}R_{C-y})₀₋₄-R_{C-heteroaryl},

-cyclopentyl or -cyclohexyl ring fused to a R_{C-aryl} or R_{C-heteroaryl} or R_{C-}

30 heterocycle.

160. Use of a substituted amine of formula (XV) according to claim 148 where R₁
 is:

-(CH₂)-(R_{1-aryl}) where R_{1-aryl} is phenyl.

161. Use of a substituted amine of formula (XV) according to claim 160 where R_1 is:

$-(CH_2)-(R_{1-aryl})$ where R_{1-aryl} is phenyl substituted with two -F.

5

162. Use of a substituted amine of formula (XV) according to claim 161 where the -F substitution is 3,5-difluorobenzyl.

163. Use of a substituted amine of formula (XV) according to claim 148 where R_2 is
10 -H.

164. Use of a substituted amine of formula (XV) according to claim 148 where R_3 is -H.

15 165. Use of a substituted amine of formula (XV) according to claim 148 where R_N is

$R_{N-1}-X_N-$ where X_N is -CO-, where R_{N-1} is R_{N-aryl} where R_{N-aryl} is phenyl substituted with one -CO-NR_{N-2}R_{N-3} where the substitution on phenyl is 1,3-.

20 166. Use of a substituted amine of formula (XV) according to claim 165 where R_{N-2} and R_{N-3} are the same and are C₃ alkyl.

167. Use of a substituted amine of formula (XV) according to claim 148 where R_N is

25 $R_{N-1}-X_N-$ where X_N is -CO-, where R_{N-1} is R_{N-aryl} where R_{N-aryl} is phenyl substituted with one C₁ alkyl and with one -CO-NR_{N-2}R_{N-3} where the substitution on the phenyl is 1,3,5-.

30 168. Use of a substituted amine of formula (XV) according to claim 167 where R_{N-2} and R_{N-3} are the same and are C₃ alkyl.

169. Use of a substituted amine of formula (XV) according to claim 148 where R_N is

$R_{N-1}-X_N$ where X_N is $-\text{CO}-$, where R_{N-1} is $R_{N\text{-heteroaryl}}$ where $R_{N\text{-heteroaryl}}$ is substituted with one $-\text{CO}-\text{NR}_{N-2}\text{R}_{N-3}$.

170. Use of a substituted amine of formula (XV) according to claim 169 where R_{N-2} and R_{N-3} are the same and are $-\text{C}_3$ alkyl.

171. Use of a substituted amine of formula (XV) according to claim 148, where R_A is:

$-(\text{CR}_{A-x}\text{R}_{A-y})_{0-4}-\text{R}_{A\text{-aryl}}$ where $\text{R}_{A\text{-aryl}}$ is phenyl,
 $-(\text{CR}_{A-x}\text{R}_{A-y})_{0-4}-\text{R}_{A\text{-heteroaryl}}$,
 $-\text{cyclopentyl}$ or $-\text{cyclohexyl}$ ring fused to a $\text{R}_{A\text{-aryl}}$ or $\text{R}_{A\text{-heteroaryl}}$ or $\text{R}_{A\text{-heterocycle}}$.

172. Use of a substituted amine of formula (XV) according to claim 171, where R_A is:

$-(\text{CR}_{A-x}\text{R}_{A-y})_{0-4}-\text{R}_{A\text{-aryl}}$ where $\text{R}_{A\text{-aryl}}$ is phenyl.

173. Use of a substituted amine of formula (XV) according to claim 172, where phenyl is substituted in the 3-position or 3,5-positions.

174. Use of a substituted amine of formula (XV) according to claim 171, where R_A is

$-(\text{CH}_2)-\text{R}_{A\text{-heteroaryl}}$.

175. Use of a substituted amine of formula (XV) according to claim 171, where R_A is:

$-(\text{CH}_2)-\text{R}_{A\text{-heterocycle}}$.

176. Use of a substituted amine of formula (XV) according to claim 175, where R_A is:

$-\text{cyclohexyl}$ ring fused to a phenyl ring.

177. Use of a substituted amine of formula (XV) according to claim 148, where R_B is:

$-(\text{CR}_{\text{B-x}}\text{R}_{\text{B-y}})_{0-4}-\text{R}_{\text{B-aryl}}$ where $\text{R}_{\text{B-aryl}}$ is phenyl,
 $-(\text{CR}_{\text{B-x}}\text{R}_{\text{B-y}})_{0-4}-\text{R}_{\text{B-heteroaryl}}$,
 -cyclopentyl or -cyclohexyl ring fused to a $\text{R}_{\text{B-aryl}}$ or $\text{R}_{\text{B-heteroaryl}}$ or $\text{R}_{\text{B-heterocycle}}$.

heterocycle-

5

178. Use of a substituted amine of formula (XV) according to claim 177, where R_{B} is:

$-(\text{CR}_{\text{B-x}}\text{R}_{\text{B-y}})_{0-4}-\text{R}_{\text{B-aryl}}$ where $\text{R}_{\text{B-aryl}}$ is phenyl.

10 179. Use of a substituted amine of formula (XV) according to claim 178 where phenyl is substituted in the 3-position or 3,5-positions.

180. Use of a substituted amine of formula (XV) according to claim 179 where R_{B} is:

15 $-(\text{CH}_2)-\text{R}_{\text{B-heteroaryl}}$.

181. Use of a substituted amine of formula (XV) according to claim 177 where R_{B} is:

$-(\text{CH}_2)-\text{R}_{\text{B-heterocycle}}$.

20

182. Use of a substituted amine of formula (XV) according to claim 177 where R_{B} is:

-cyclohexyl ring fused to a phenyl ring.

25 183. Use of a substituted amine of formula (XV) according to claim 148 where the pharmaceutically acceptable salt is selected from the group consisting of salts of the following acids acetic, aspartic, benzenesulfonic, benzoic, bicarbonic, bisulfuric, bitartaric, butyric, calcium edetate, camrylic, carbonic, chlorobenzoic, citric, edetic, edisylic, estolic, esyl, esylic, formic, fumaric, gluceptic, gluconic, glutamic,
 30 glycolylarsanilic, hexamic, hexylresorcinoic, hydrabamic, hydrobromic, hydrochloric, hydroiodic, hydroxynaphthoic, isethionic, lactic, lactobionic, maleic, malic, malonic, mandelic, methanesulfonic, methylnitric, methylsulfuric, mucic, muconic, napsylic, nitric, oxalic, p-nitromethanesulfonic, pamoic, pantothenic, phosphoric, monohydrogen phosphoric, dihydrogen phosphoric, phthalic,

polygalactouronic, propionic, salicylic, stearic, succinic, sulfamic, sulfanilic, sulfonic, sulfuric, tannic, tartaric, teoclic and toluenesulfonic.

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
10 January 2002 (10.01.2002)

PCT

(10) International Publication Number
WO 02/002506 A3

(51) International Patent Classification⁷: **C07C 243/22**,
243/28, 239/20, A61K 31/165, A61P 25/28

(21) International Application Number: PCT/US01/20930

(22) International Filing Date: 29 June 2001 (29.06.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/215,323 30 June 2000 (30.06.2000) US

(71) Applicant: **ELAN PHARMACEUTICALS, INC.**
[US/US]; 800 Gateway Boulevard, So. San Francisco, CA
94080 (US).

(72) Inventors: **FANG, Lawrence, Y.**; 1193 Beach Park Boule-
vard, Foster City, CA 94404 (US). **JOHN, Varghese**; 1772
18th Avenue, San Francisco, CA 94122 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT (util-
ity model), AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH,
CN, CO, CR, CU, CZ (utility model), DE (utility model),

DK (utility model), DM, DZ, EC, EE (utility model), ES,
FI (utility model), GB, GD, GE, GH, GM, HR, HU, ID, IL,
IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU,
LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL,
PT, RO, RU, SD, SE, SG, SI, SK (utility model), SL, TJ,
TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments


(88) Date of publication of the international search report:
29 August 2002

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.



WO 02/002506 A3

(54) Title: COMPOUNDS TO TREAT ALZHEIMER'S DISEASE



(57) Abstract: The present invention is substi-
tuted amines of formula (XV) useful in treating
Alzheimer's disease and other similar diseases.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 01/20930

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07C243/22 C07C243/28 C07C239/20 A61K31/165 A61P25/28

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07C A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

CHEM ABS Data, BEILSTEIN Data, EPO-Internal, WPI Data, PAJ

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 99 65870 A (VERTEX PHARMACEUTICALS INC) 23 December 1999 (1999-12-23) page 74, lines 11-12; example 54, step 1; example 80, step 1; example 112, step 1; example 153, steps 2-3; example 161, steps 1-2 ---	31, 34, 46-49, 51-53
A	WO 98 33795 A (THE REGENTS OF THE UNIVERSITY OF CALIFORNIA) 6 August 1998 (1998-08-06) cited in the application page 4, lines 8-26; page 7, lines 21-26; table 1; page 30, line 1 - page 32, line 28; claims --- -/-	1, 54, 55, 65, 95, 115, 121, 124, 126, 131, 133, 134, 144-149

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

° Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

31 May 2002

Date of mailing of the international search report

10. 6. 02.*

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Van Amsterdam, L

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 01/20930

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	N. CHEVALLIER ET AL: BRAIN RESEARCH, vol. 750, no. 1/2, 1997, pages 11-19, XP000921314 -----	
A	EP 0 652 009 A (ELI LILLY AND CO ET AL) 10 May 1995 (1995-05-10) cited in the application -----	

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US 01/20930

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
see FURTHER INFORMATION sheet PCT/ISA/210
2. ☒ Claims Nos.:
—
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
see FURTHER INFORMATION sheet PCT/ISA/210
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see additional sheet

1. ☒ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☒ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International Application No. PCT/US 01 /20930

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.1

Although claims 54-94, 113-114, 122-132 and, in as far as they relate to a treatment in vivo, claims 95-110, 112, 115-116, 118-121, 134 and 136 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.

Continuation of Box I.2

Formula XV of claim 1 and formulae II, III, IV, XI and XII of claims 31, 36, 41, 46 and 51, respectively, all relate to an extremely large number of possible compounds. In fact, the claims contain so many options and possible permutations that a lack of clarity and conciseness within the meaning of Article 6 PCT arises to such an extent as to render a meaningful search of the claims impossible. The search has been guided by those parts of the application which do appear to be sufficiently clear and concise, namely examples 1-3 and CHARTS A, B and C' and has been carried out for compounds of formulae XV, II, III, IV, XI and XII, wherein, where appropriate,

R1 = (VI) with n1 = 1 and R1-aryl = phenyl, optionally substituted with 1-4 the same or different (A)-(N);

R2 = R3 = H;

RN = (I) with XN = (A) and RN-1 = (A) = phenyl, optionally substituted with 1-4 the same or different (1)-(30)/(34)-(39);

RA = (I), (II), (III) with RA-x = RA-y = H and RA-aryl = RN-aryl = phenyl, or (XXVIII);

X = -N or -O;

RB = (I), (II), (III) with RB-x = RB-y = H and RB-aryl = RN-aryl = phenyl, or (XXVIII);

PROTECTING GROUP = an oxycarbonyl containing group as defined in claim 31, page 198, lines 9-27.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

INTERNATIONAL SEARCH REPORT

International Application No. **PCT/US 01 /20930**

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. Claims: 1-45, 51-183

Compounds of formula XV and their use in the treatment of Alzheimer's disease; intermediates of formulae II, III, IV and XII.

2. Claims: 46-50

Intermediates of formula XI.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 01/20930

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9965870	A	23-12-1999	AU 4576099 A	05-01-2000
			BR 9912169 A	10-04-2001
			EP 1086076 A1	28-03-2001
			NO 20006405 A	19-02-2001
			WO 9965870 A2	23-12-1999
			US 2002049201 A1	25-04-2002
WO 9833795	A	06-08-1998	AU 6268698 A	25-08-1998
			EP 0958293 A1	24-11-1999
			JP 2001510474 T	31-07-2001
			US 6150416 A	21-11-2000
			WO 9833795 A1	06-08-1998
EP 652009	A	10-05-1995	AU 6897094 A	16-02-1995
			CA 2129689 A1	10-02-1995
			CN 1120040 A	10-04-1996
			CZ 9401841 A3	15-03-1995
			EP 0652009 A1	10-05-1995
			JP 7165606 A	27-06-1995
			NO 942883 A	10-02-1995
			NZ 264143 A	26-11-1996
			ZA 9405719 A	01-02-1996